

Source Habitats for Terrestrial Vertebrates of Focus in the Interior Columbia Basin: Broad-Scale Trends and Management Implications

Volume 2 - Group Level Results

Michael J. Wisdom, Richard S. Holthausen, Barbara C. Wales, Christina D. Hargis, Victoria A. Saab, Danny C. Lee, Wendel J. Hann, Terrell D. Rich, Mary M. Rowland, Wally J. Murphy, and Michelle R. Eames

NOTE: This is a compact disc version of PNW-GTR-485 in draft form as it was submitted by the authors to the Pacific Northwest Research Station for layout and production of galley proofs. Minor differences in style and content are expected between this compact disc version and the final publication of this GTR in hard-copy format. Differences are due to layout and final editing. The substance, however, will remain the same between this compact disc version and the final, hard-copy publication.

AUTHORS

MICHAEL J. WISDOM is a terrestrial ecologist, and BARBARA C. WALES is a wildlife biologist, U.S. Department of Agriculture (USDA), Forest Service, Pacific Northwest Research Station, Forestry and Range Sciences Laboratory, 1401 Gekeler Lane, La Grande, OR 97850; RICHARD S. HOLTHAUSEN is the national wildlife ecologist, and CHRISTINA D. HARGIS is the associate national wildlife ecologist, USDA Forest Service, Forestry Sciences Laboratory, 2500 S. Pine Knoll, Flagstaff, AZ 86001; DANNY C. LEE was a research biologist, and VICTORIA A. SAAB is a research wildlife biologist, USDA Forest Service, Forestry Sciences Laboratory, 316 E. Myrtle St., Boise, ID 83702; WENDEL J. HANN is the national landscape ecologist, National Interagency Fire Center, Boise, ID 83705; TERRELL D. RICH is the senior avian ecologist, U.S. Department of the Interior (USDI), Bureau of Land Management, 1387 S. Vinnell Way, Boise, ID 83709; MARY M. ROWLAND is a wildlife biologist, USDI Bureau of Land Management, Forestry and Range Sciences Laboratory, 1401 Gekeler Lane, La Grande, OR 97850; WALLY J. MURPHY was a natural resources staff officer, USDA Forest Service, Umatilla National Forest, 2517 S. W. Hailey, Pendleton, OR 97801; and MICHELLE R. EAMES is a wildlife biologist, USDI Fish and Wildlife Service, 11103 E. Montgomery Drive, Suite 2, Spokane, WA 99206.

CONTRIBUTORS

David A. Newhouse
Fred B. Samson
Alan Sands
Nancy Warren

Source Habitats for Terrestrial Vertebrates of Focus in the Interior Columbia Basin: Broad-Scale Trends and Management Implications

Volume 2--Group Level Results

Michael J. Wisdom, Richard S. Holthausen, Barbara C. Wales, Christina D. Hargis, Victoria A. Saab, Danny C. Lee, Wendel J. Hann, Terrell D. Rich, Mary M. Rowland, Wally J. Murphy, and Michelle R. Eames

Interior Columbia Basin Ecosystem Management Project: Scientific Assessment

Thomas M. Quigley, Editor

Volume 2 contains pages 1 through 222.

U.S. Department of Agriculture
Forest Service
Pacific Northwest Research Station
Portland, Oregon
General Technical Report PNW-GTR-485
2000

ABSTRACT

Wisdom, Michael J.; Holthausen, Richard S.; Wales, Barbara C.; Hargis, Christina D.; Saab, Victoria A.; Lee, Danny C.; Hann, Wendel J.; Rich, Terrell D.; Rowland, Mary M.; Murphy, Wally J.; Eames, Michelle R. 2000. Source habitats for terrestrial vertebrates of focus in the interior Columbia basin: broad-scale trends and management implications. Volume 2—group level results. Gen. Tech. Rep. PNW-GTR-485. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 3 vol. (Quigley, Thomas M., tech. ed.; Interior Columbia Basin Ecosystem Management Project: scientific assessment).

We defined habitat requirements (source habitats) and assessed trends in these habitats for 91 species of terrestrial vertebrates on 58 million ha (145 million acres) of public and private lands within the interior Columbia basin (basin). We also summarized knowledge about species-road relations for each species and mapped source habitats in relation to road densities for four species of terrestrial carnivores. Our assessment was conducted as part of the Interior Columbia Basin Ecosystem Management Project (ICBEMP), a multiresource, multidisciplinary effort by the USDA Forest Service (FS) and the USDI Bureau of Land Management (BLM) to develop an ecosystem-based strategy for managing FS and BLM lands within the basin. Our assessment was designed to provide technical support for the ICBEMP and was done in five steps. First, we identified species of terrestrial vertebrates for which there was ongoing concern about population or habitat status (species of focus), and for which habitats could be estimated reliably by using a large mapping unit (pixel size) of 100 ha (247 acres) and broad-scale methods of spatial analysis. Second, we evaluated change in source habitats from early European settlement (historical, circa 1850 to 1890) to current (circa 1985 to 1995) conditions, for each species and for hierarchically nested groups of species and families of groups, at the spatial scales of the watershed (5th hydrologic unit code [HUC]), subbasin (4th HUC), ecological reporting unit (ERU), and basin. Third, we summarized the effects of roads and road-associated factors on populations and habitats for each of the 91 species and described the results in relation to broad-scale patterns of road density. Fourth, we mapped classes of the current abundance of source habitats for four species of terrestrial carnivores in relation to classes of road density across the 164 subbasins and used the maps to identify areas having high potential to support persistent populations. And fifth, we used our results, along with results from other studies, to describe broad-scale implications for managing habitats deemed to have undergone long-term decline and for managing species negatively affected by roads or road-associated factors.

Our results indicated that habitats for species, groups, and families associated with old-forest structural stages, with native grasslands, or with native shrublands have undergone strong, widespread decline. Implications of these results for managing old-forest structural stages include consideration of (1) conservation of habitats in subbasins and watersheds where decline in old forests has been strongest; (2) silvicultural manipulations of mid-seral forests to accelerate development of late-seral stages; and (3) long-term silvicultural manipulations and long-term accommodation of fire and other disturbance regimes in all forested structural stages to hasten development and improvement in the amount, quality, and distribution of old-forest stages. Implications of our results for managing rangelands include the potential to (1) conserve native grasslands and shrublands that have not undergone large-scale reduction in composition of native plants; (2) control or eradicate exotic plants on native grasslands and shrublands where invasion potential or spread of exotics is highest; and (3) restore native plant communities by using intensive range practices where potential for restoration is highest.

Our analysis also indicated that >70 percent of the 91 species are affected negatively by one or more factors associated with roads. Moreover, maps of the abundance of source habitats in relation to classes of road density suggested that road-associated factors hypothetically may reduce the potential to support persistent populations of terrestrial carnivores in many subbasins. Management implications of our summarized road effects include the potential to mitigate a diverse set of negative factors associated with

roads. Comprehensive mitigation of road-associated factors would require a substantial reduction in the density of existing roads as well as effective control of road access in relation to management of livestock, timber, recreation, hunting, trapping, mineral development, and other human activities.

A major assumption of our work was that validation research will be conducted by agency scientists and other researchers to corroborate our findings. As a preliminary step in the process of validation, we found high agreement between trends in source habitats and prior trends in habitat outcomes that were estimated as part of the viability analysis for terrestrial species within the basin. Results of our assessment also were assumed to lead to finer scale evaluations of habitats for some species, groups, or families as part of implementation procedures. Implementation procedures are necessary to relate our findings to local conditions; this would enable managers to effectively apply local conservation and restoration practices to support broad-scale conservation and restoration strategies that may evolve from our findings.

Keywords: Cluster analysis, conservation, forest management, habitat, habitat condition, habitat management, habitat trend, interior Columbia basin, Interior Columbia Basin Ecosystem Management Project, landscape ecology, landscape analysis, population viability, rangeland management, terrestrial vertebrates, spatial analysis, species of focus, sink, sink environment, source, source environment, source habitat, source habitats, restoration, species groups, monitoring, validation research, viability, wildlife, wildlife-habitat relations.

FOREWORD

This publication consists of three volumes. Use of three volumes was necessary so that our findings--which consist of hundreds of tables, figures, pages of text, and supporting citations--could be presented in a manner most usable to resource managers, biologists, and the public. That goal in mind, volume 1 is designed as an overview of objectives, methods, key results, and management implications. Volumes 2 and 3 contain increasingly detailed results that support and complement results in volume 1. We believe that resource managers may find sufficient detail in the generalized results and implications presented in volume 1, but that management biologists and other users of the results and supporting data will want to refer to all three volumes. Results, management implications, and supporting citations provided in volume 2 are especially important to consider as part of step-down implementation procedures and related management conducted by field units within the interior Columbia basin. By contrast, information in volume 1 may be particularly useful in serving broad-scale planning issues, objectives, and strategies for the interior Columbia basin as a whole. Regardless of application, all three volumes are intended to function together as a comprehensive assessment of habitat trends and a summary of other environmental factors affecting terrestrial vertebrates whose population or habitat status is of ongoing concern to resource managers.

Contents

ABSTRACT	4
FOREWORD	6
LIST OF FIGURES	10
INTRODUCTION	24
GROUP 1--PYGMY NUTHATCH, WHITE-BREASTED NUTHATCH, AND WHITE-HEADED WOODPECKER	26
GROUP 2 - LEWIS' WOODPECKER (MIGRANT POPULATION)	30
GROUP 3--WESTERN GRAY SQUIRREL	34
GROUP 4--BLUE GROUSE (WINTER)	37
GROUP 5--NORTHERN GOSHAWK (SUMMER), FLAMMULATED OWL, AMERICAN MARTEN, AND FISHER	39
GROUP 6 - VAUX'S SWIFT, WILLIAMSON'S SAPSUCKER, PILEATED WOODPECKER, HAMMOND'S FLYCATCHER, CHESTNUT-BACKED CHICKADEE, BROWN CREEPER, WINTER WREN, GOLDEN-CROWNED KINGLET, VARIED THRUSH, SILVER-HAIRED BAT, AND HOARY BAT ..	46
GROUP 7--BOREAL OWL	50
GROUP 8--GREAT GRAY OWL	54
GROUP 9--BLACK-BACKED WOODPECKER	57
GROUP 10--OLIVE-SIDED FLYCATCHER	62
GROUP 11 - THREE-TOED WOODPECKER AND WHITE-WINGED CROSSBILL	64
GROUP 12--WOODLAND CARIBOU	67
GROUP 13 --NORTHERN FLYING SQUIRREL	70
GROUP 14--HERMIT WARBLER	73
GROUP 15--PYGMY SHREW AND WOLVERINE	74
GROUP 16--LYNX	78
GROUP 17--BLUE GROUSE (SUMMER) AND MOUNTAIN QUAIL (SUMMER)	81
GROUP 18--LAZULI BUNTING	85
GROUP 19--GRAY WOLF AND GRIZZLY BEAR	88

GROUP 20--MOUNTAIN GOAT	94
GROUP 21--LONG-EARED OWL	97
GROUP 22--CALIFORNIA BIGHORN SHEEP AND ROCKY MOUNTAIN BIGHORN SHEEP	100
GROUP 23--RUFIOUS HUMMINGBIRD AND BROAD-TAILED HUMMINGBIRD	105
GROUP 24--SHARPTAIL SNAKE, CALIFORNIA MOUNTAIN KINGSSNAKE, AND BLACK-CHINNED HUMMINGBIRD	108
GROUP 25--NORTHERN GOSHAWK (WINTER)	111
GROUP 26--YUMA MYOTIS, LONG-EARED MYOTIS, FRINGED MYOTIS, AND LONG-LEGGED MYOTIS	114
GROUP 27--PINE SISKIN AND TOWNSEND'S BIG-EARED BAT	119
GROUP 28--SPOTTED BAT, PALLID BAT, AND WESTERN SMALL-FOOTED MYOTIS	124
GROUP 29--WESTERN BLUEBIRD	127
GROUP 30--ASH-THROATED FLYCATCHER AND BUSHTIT	131
GROUP 31--FERRUGINOUS HAWK, BURROWING OWL, SHORT-EARED OWL, VESPER SPARROW, LARK SPARROW, WESTERN MEADOWLARK, and PRONGHORN	133
GROUP 32--PREBLE'S SHREW, UINTA GROUND SQUIRREL, WHITE-TAILED ANTELOPE SQUIRREL, WYOMING GROUND SQUIRREL, WASHINGTON GROUND SQUIRREL, STRIPED WHIPSNAKE, LONGNOSE SNAKE, GROUND SNAKE, MOJAVE BLACK-COLLARED LIZARD, AND LONGNOSE LEOPARD LIZARD	140
GROUP 33--BREWER'S SPARROW, LARK BUNTING, SAGE SPARROW, SAGE THRASHER, SAGE GROUSE, PYGMY RABBIT, AND SAGEBRUSH VOLE	147
GROUP 34--KIT FOX AND BLACK-THROATED SPARROW	153
GROUP 35--LOGGERHEAD SHRIKE	156
GROUP 36 --COLUMBIAN SHARP-TAILED GROUSE (SUMMER)	161
GROUP 37--GRASSHOPPER SPARROW, CLAY-COLORED SPARROW, AND IDAHO GROUND SQUIRREL	166
GROUP 38--BLACK ROSY FINCH AND GRAY-CROWNED ROSY FINCH	173
GROUP 39--LEWIS' WOODPECKER (RESIDENT POPULATION)	175
GROUP 40--BROWN-HEADED COWBIRD	179
ABBREVIATIONS	182

REFERENCES 183

FOOTNOTES 221

LIST OF FIGURES

Figure 1--Assessment boundaries of the Interior Columbia Basin Ecosystem Management Project and the 13 ecological reporting units.

Figure 2--Land ownership within the Interior Columbia basin Ecosystem Management Project science assessment area.

Figure 3--Ranges of species in group 1 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 4--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 1 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 5--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 1, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by n .

Figure 6--Ranges of species in group 2 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 7--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 2 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 8--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 2, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by n .

Figure 9--Ranges of species in group 3 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 10--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 3

within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 11--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 3, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 12--Ranges of species in group 4 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 13--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 4 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 14--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 4, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 15--Ranges of species in group 5 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 16--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 5 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 17--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 5, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 18--Ranges of species in group 6 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 19--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 6 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 20--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 6, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 21--Ranges of species in group 7 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 22--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 7 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 23--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 7, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 24--Ranges of species in group 8 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 25--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 8 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 26--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 8, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 27--Ranges of species in group 9 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and

current ranges; otherwise, the current range map also denotes the historical range.

Figure 28--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 9 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 29--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 9, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 30--Ranges of species in group 10 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 31--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 10 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 32--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 10, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 33--Ranges of species in group 11 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 34--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 11 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 35--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 11, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 36--Ranges of species in group 12 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 37--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 12 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 38--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 12, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 39--Ranges of species in group 13 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 40--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 13 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 41--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 13, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 42--Ranges of species in group 14 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 43--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 14 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 44--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 14, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of

watersheds from which estimates were derived is denoted by \underline{n} .

Figure 45--Ranges of species in group 15 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 46--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 15 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 47--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 15, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 48--Ranges of species in group 16 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 49--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 16 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 50--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 16, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 51--Ranges of species in group 17 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 52--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 17 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 53--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 17, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase

of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 54--Ranges of species in group 18 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 55--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 18 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 56--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 18, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 57--Ranges of species in group 19 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 58--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 19 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 59--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 19, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 60--Ranges of species in group 20 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 61--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 20 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 62--Percentage of watersheds within five trend categories of relative change in source habitats

from historical to current periods for group 20, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 63--Ranges of species in group 21 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 64--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 21 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 65--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 21, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 66--Ranges of species in group 22 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 67--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 22 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 68--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 22, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 69--Ranges of species in group 23 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 70--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 23 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 71--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 23, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 72--Ranges of species in group 24 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 73--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 24 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 74--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 24, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 75--Ranges of species in group 25 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 76--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 25 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 77--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 25, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 78--Ranges of species in group 26 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 79--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 26 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent;

and 2 = an increase of ≥ 60 percent.

Figure 80--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 26, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 81--Ranges of species in group 27 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 82--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 27 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 83--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 27, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 84--Ranges of species in group 28 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 85--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 28 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 86--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 28, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 87--Ranges of species in group 29 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 88--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 29 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of

five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 89--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 29, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 90--Ranges of species in group 30 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 91--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 30 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 92--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 30, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 93--Ranges of species in group 31 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 94--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 31 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 95--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 31, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 96--Ranges of species in group 32 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 97--Percentage of area identified as source habitats, historically (A) and currently (B), and the

relative change in percentage of area of source habitats from historical to current periods (C), for group 32 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 98--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 32, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 99--Ranges of species in group 33 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 100--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 33 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 101--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 33, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 102--Ranges of species in group 34 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 103--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 34 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 104--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 34, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 105--Ranges of species in group 35 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 106--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 35 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 107--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 35, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 108--Ranges of species in group 36 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 109--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 36 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 110--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 36, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 111--Ranges of species in group 37 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 112--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 37 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 113--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 37, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 114--Ranges of species in group 38 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for

historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 115--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 38 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 116--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 38, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 117--Ranges of species in group 39 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 118--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 39 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 119--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 39, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

Figure 120--Ranges of species in group 40 within the basin (from Marcot and others, in prep.). For species whose ranges shifted significantly from historical conditions, separate maps are shown for historical and current ranges; otherwise, the current range map also denotes the historical range.

Figure 121--Percentage of area identified as source habitats, historically (A) and currently (B), and the relative change in percentage of area of source habitats from historical to current periods (C), for group 40 within each of 2,562 watersheds in the basin. Relative change for each watershed is shown as one of five trend categories, where -2 = a decrease of ≥ 60 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; 1 = an increase of ≥ 20 percent but < 60 percent; and 2 = an increase of ≥ 60 percent.

Figure 122--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 40, basin-wide and by ecological reporting units. Trend categories correspond to the following relative changes from historical to current periods: 2 = an increase of ≥ 60 percent; 1 = an increase of ≥ 20 percent but < 60 percent; 0 = an increase or decrease of < 20 percent; -1 = a decrease of ≥ 20 percent but < 60 percent; and -2 = a decrease of ≥ 60 percent. Number of watersheds from which estimates were derived is denoted by \underline{n} .

INTRODUCTION

This volume is the second in a three-volume publication that defines and assesses trends in source habitats for 91 terrestrial vertebrate species within the interior Columbia River basin (hereafter referred to as "basin") (See "Glossary," vol. 3, for terms used in this paper). This assessment was conducted as part of the Interior Columbia Basin Ecosystem Management Project (ICBEMP), a multiresource, multidisciplinary effort by the USDA Forest Service (FS) and the USDI Bureau of Land Management (BLM) to develop an ecosystem-based strategy for managing lands within the basin administered by the FS and BLM. The assessment area extends over 58 million ha¹ (145 million acres) in eastern Washington, eastern Oregon, Idaho, western Montana, and small portions of Nevada, California, Wyoming, and Utah (figs. 1 and 2). The purpose of this publication is to provide technical support to the ICBEMP regarding trends in the aerial extent of wildlife habitats in the basin, as well as management implications regarding those trends. Additionally, it can be used to provide a broad-scale view of how wildlife habitats have changed in the basin since early European settlement and factors that have contributed to those changes.

Vol. 2, Figure 1--Assessment boundaries of the Interior Columbia Basin Ecosystem Management Project and the 13 ecological reporting units.

Vol. 2, Figure 2--Land ownership within the Interior Columbia basin Ecosystem Management Project science assessment area. pdf version (6.9mb) lo-res version (71kb)

This publication is focused on source habitats rather than all habitats in which a species is known to occur. Source habitats are those characteristics of macrovegetation that contribute to stationary or positive population growth for a species in a specified area and time. Source habitats contribute to source environments (Pulliam 1988, Pulliam and Danielson 1991), which represent the composite of all environmental conditions that results in stationary or positive population growth for a species in a specified area and time. The distinction between source habitats and source environments is important for understanding our evaluation and its limitations. For example, source habitats for a bird species during the breeding season would include those characteristics of vegetation that contribute to successful nesting and rearing of young, but would not include nonvegetative factors, such as the effects of pesticides on thinning of eggshells, which also affect production of young. Consequently, we have attempted to identify all factors that affect population performance of each species as a complement to our explicit analysis of source habitats. As the foundation for our analysis, we relied on published literature and guidance from species experts to identify source habitats and additional factors that presumably affect population performance.

The 91 species in our analysis are organized into 40 groups, 37 of which are then organized into 12 families. Groups are composed of one or more species that share common source habitats, as defined by vegetation cover types and structural stages. Similar groups are also clustered into families whose source habitats generally fall into similar terrestrial community groups, a broader classification that includes several cover types. Group size ranges from 1 to 17 species, and family size ranges from one to nine groups.

Volume 1 describes methods used to select species for analysis, place them in groups and families, estimate source habitats, and analyze habitat trends. That volume also includes general analyses of source habitat trends at all three levels--species, group, and family--including a correlation analysis that evaluates how well species-level trends in source habitats are reflected in the higher level group- and family-level trends. Volume 1 also identifies causes for the observed trends and ecological processes important for maintaining source habitats as part of the family-level results. Additionally, volume 1 provides a special section on species and groups that are negatively affected by road-related human activities. In volume 2, we present more detailed results on the analysis of source habitat trends at the group level in support of the more generalized results presented in volume 1. The appendices in volume 3 provide further data and results in support of both volumes 1 and 2.

For each of the 40 groups discussed in volume 2, we specifically present results on source habitat trends, interpret those results, and discuss management implications. In the results section, we list the species included in each group, display range maps for each of the species, and describe source habitats and special habitat features for each species. Source habitats and special habitat features with each species in each group and family are listed in volume 3, appendix 1, tables 1 and 2.

In the results section of volume 2, we specifically display maps that compare the historical and current distribution of source habitats within the basin for each group, and describe changes in areal extent that were projected to have occurred since the historical period. These changes are analyzed at the watershed level, a unit of land that whose mean size is about 22 500 ha (56,000 acres). The watershed results are summarized by ecological reporting units (ERUs), which represent 13 broad geographical regions within the basin (fig. 1) that differ significantly in biophysical characteristics (Hann and others 1997).

The section on interpretation of results in volume 2 consists of four components. First, we provide a description of the vegetation changes that underlie source habitat changes. Ecological processes and management actions that caused the vegetation changes are described in volume 1, and that information is not repeated here. Second, changes from historical to current in the condition of special habitat features are disclosed for those features for which information is available. Third, factors other than habitat that have a significant effect on species in the group are discussed, with these discussions frequently focusing on the effects of specific management activities and other human disturbances. Finally, any available data on population status and trends for any species in the group are presented. We have not attempted any correlations or discussion of anecdotal similarity between habitat trends and population trends because our habitat analysis addresses different time frames and different geographic areas than population trend data available for most species.

The final section of volume 2 discusses management implications based on both the findings of this analysis and on published literature for each group of species. Management implications are presented in three parts. First, issues relevant to species in the group are discussed. These include issues related to broad-scale source habitats, special habitat features, and other factors that significantly influence the group. Broad strategies that could be used to resolve these issues are presented, and geographic priorities for the strategies are offered where appropriate. The third part of the management implications section consists of specific on-the-ground management practices that could be used in the implementation of the strategies. In all cases, the discussion of strategies and practices is intended to be addressed within the context of broader ecosystem-based objectives. Implementation of the strategies and practices for any single group without consideration of other ecosystem elements would not be appropriate.

The list of strategies and practices outlined for each group of species in volume 2 should be considered a menu of possible approaches that could be adopted by managers to help achieve their objectives for conservation and restoration of habitats. Before any of these approaches are adopted, they should be

analyzed to determine their effectiveness, their compatibility with overall ecosystem management objectives, and their applicability to specific situations. Testing and validation should continue through all the geographic scales of implementation.

In summary, the strategies presented at the family level in volume 1 represent a synthesis of similar group strategies developed in volume 2. Volume 1 therefore provides a broader, more generalized perspective of source habitat trends in the basin, whereas volume 2 offers a more specific, in-depth coverage of the same analysis. Thus, users of our publication can refer to volume 1 for an overview of results and implications, refer to volume 2 for detailed results that support the overview, and refer to volume 3 for the most specific results and information in support of both volumes 1 and 2.

GROUP 1--PYGMY NUTHATCH, WHITE-BREASTED NUTHATCH, AND WHITE-HEADED WOODPECKER

Results

Species ranges, source habitats, and special habitat features--Group 1 consists of the pygmy nuthatch, white-breasted nuthatch, and white-headed woodpecker, all of which are year-round residents within the basin.² The pygmy nuthatch is widespread except for the Columbia Plateau and southern portions of the basin, and the white-breasted nuthatch occurs throughout most of the basin (fig. 3). The white-headed woodpecker has the most restricted range, occurring in the eastern slope of the Cascade Range, the Blue Mountains, the Okanogan Mountains, and mountains of Idaho. Source habitats for group 1 are found in old forests of Sierra Nevada mixed-conifer and ponderosa pine cover types. The white-breasted nuthatch also breeds in old forests of aspen and cottonwood-willow, in Oregon white oak, and in unmanaged young forests of interior ponderosa pine (vol. 3, appendix 1, table 1).

Vol. 2, Figure 3--Ranges of species in group 1 within the basin.

A special habitat feature for group 1 is large-diameter snags for nesting and foraging (vol. 3, appendix 1, table 2). Both nuthatches are secondary cavity nesters and can use various nesting structures (McEllin 1979), whereas the white-headed woodpecker is a primary cavity excavator of soft snags and is therefore more limited by the degree of wood decay suitable for nest hole excavation (Garrett and others 1996). White-headed woodpeckers typically nest in snags and leaning logs, and occasionally nest in the dead tops of live trees (Garrett and others 1996, Milne and Hejl 1989). White-breasted nuthatches nest in natural cavities of live ponderosa pine more often than in snags (Brawn and Balda 1988, McEllin 1979). Suitable nest sites for all three species usually are found within the upper diameter classes of trees and snags. Average diameters reported for nest trees are 57.93 ± 3.65 cm (22.80 ± 1.43 in [$\bar{x} \pm SE$]) for pygmy nuthatch (McEllin 1979), 53.77 ± 1.56 cm (21.16 ± 0.61 in [$\bar{x} \pm SE$]) for white-breasted nuthatch (McEllin 1979), and 80 ± 65 cm (31 ± 25 in [$\bar{x} \pm SE$]) for white-headed woodpecker (Garrett and others 1996, Milne and Hejl 1989).

All three species forage primarily on live trees. White-breasted nuthatches glean insects from tree trunks and were observed in Colorado to spend nearly 75 percent of foraging time on ponderosa pine trunks (Bock 1969). In the same study, pygmy nuthatches foraged more generally in live ponderosa pine, dividing their foraging time fairly equally among needles, branches, and trunks. In Oregon, 80 percent of white-headed woodpecker foraging time was on live trees, and a preference was shown for trees with

diameters >25 cm (10 in) (Bull and others 1986a).

Broad-scale changes in source habitats--Source habitats for group 1 likely occurred throughout the mountainous areas of the basin historically, and were most extensive throughout the Cascade Range, the Okanogan Mountains, and in central Oregon (fig. 4A). Currently, source habitats cover roughly the same geographical extent, but habitat patches appear more disjunct (fig. 4B). The Upper Klamath ERU continues to provide extensive source habitats, but elsewhere, <25 percent of most watersheds within the distribution of these species currently contains source habitats.

Basin-wide, >50 percent of watersheds had strong negative declines in the availability of source habitats (fig. 5). This basin-wide trend was mirrored within six ERUs that also had strong negative declines in more than 50 percent of the watersheds within the individual ERU boundaries: the Northern Cascades, Northern Glaciated Mountains, Lower Clark Fork, Upper Clark Fork, Upper Snake, and Snake Headwaters ERUs (fig. 5). Source habitats in the Upper Snake and Snake Headwaters ERUs were less than 2 percent of either ERU, both historically and currently (vol. 3, appendix 1, table 3). The extent of coverage in the Northern Cascades, Northern Glaciated Mountains, and Lower Clark Fork, however, was substantial historically, accounting for 19 to 24 percent of the total area of these ERUs (vol. 3, appendix 1, table 3). In general, areas predominated by declining trends were in the northern basin, whereas the central and southwestern parts of the basin had mixed trends (fig. 4C).

Vol. 2, Figure 4--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 1.

Vol. 2, Figure 5--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 1, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Most projected declines in source habitats were due to losses, particularly in the northern part of the basin, of late-seral forests that today are in early- and mid-seral stages (vol. 3, appendix 1, table 4). Throughout the basin, mid-seral shade-tolerant forests seem to be at nearly twice their historical levels (Hann and others 1997). A widespread change has been the transition of Pacific and interior ponderosa pine old forests to mid-seral stands of interior Douglas-fir and grand fir-white fir.

Managed young-forest structural stages of ponderosa pine, used as source habitats for the white-breasted nuthatch, generally had strongly increasing trends corresponding to the decline in old-forest structural stages. In contrast, unmanaged young forests, characterized by higher snag densities than managed forests, experienced strong declines throughout the range of group 1 (vol. 3, appendix 1, table 4).

Within the cottonwood-willow cover type, old forests had strongly declining trends throughout the basin (vol. 3, appendix 1, table 4), and generally remain only in stands smaller than the 1-km² (0.4-mi²)

mapping unit used in this analysis. These losses occurred from changes in historical hydrologic regimes. Flooding by reservoirs eliminated many cottonwood-willow stands, and reservoirs also reduced periodic flooding, a disturbance that is frequently needed for cottonwood seed establishment (Merigliano 1996, Rood and Heinze-Milne 1989). The declines in riparian woodlands and old-forest ponderosa pine documented for the basin are part of a larger picture of similar declines throughout the Western United States (Noss and others 1995).

Condition of special habitat features--Large-diameter ponderosa pine snags are a special habitat feature for group 1. In roaded areas with a history of timber sales, large-diameter snags >53 cm (21 in) have been reduced basin-wide (Hann and others 1997; Hessburg and others 1999; Quigley and others 1996). Nesting and foraging substrates for group 1 have therefore been reduced.

Other factors affecting the group--Roads indirectly affect group 1, because roaded areas in the basin have fewer snags than unroaded areas (Hann and others 1997). Roads enable snags to be cut, either in conjunction with timber sales, or by individuals seeking firewood. The additional loss of snags in areas where snags are already in low density could limit populations of group 1 species.

Population status and trends--Population trends were estimated for all three species by using Breeding Bird Survey (BBS) route data from 1966 to 1995 (Sauer and others 1996). These data have not been summarized for the basin, but summaries for various state, USDI Fish and Wildlife Service regions, and BBS physiographic regions are available. Pygmy nuthatch numbers were stable within all summary geographic areas of relevance to the basin, which were physiographic region 64 (Central Rocky Mountains), USDI Fish and Wildlife Service Region 1 (5 western states), and the Western United States (11 western states) (Sauer and others 1996). White-breasted nuthatch numbers were stable in physiographic region 64 but increased 3.6 percent annually ($\bar{n} = 149$, $P < 0.01$) in USDI Fish and Wildlife Service Region 1 and about the same throughout the Western United States. White-headed woodpecker numbers were not summarized for physiographic region 64 but increased 3.3 percent annually ($\bar{n} = 45$, $P < 0.10$) in USDI Fish and Wildlife Service Region 1 and similarly throughout the 11 western states (Sauer and others 1996).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 1 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The results of our habitat trend analysis suggest the following issues are of high priority for group 1:

1. Basin-wide decline in late-seral interior and Pacific ponderosa pine.
2. Basin-wide loss of large-diameter snags (>53 cm [21 in]).
3. High risk of additional loss of ponderosa pine habitat through stand-replacing fires.
4. Decline in old forests of aspen and cottonwood-willow.

Potential strategies--The following strategies could be used to reverse broad-scale declines in source habitats:

1. (To address issue no. 1) Retain stands of interior and Pacific ponderosa pine where old-forest conditions are present, and actively manage to promote their long-term sustainability. The white-

headed woodpecker has the most restricted distribution of all group members, and therefore the retention of existing old forests is particularly important within the range of this species where declines in old forests have been most pronounced: watersheds within the Northern Cascades, Northern Glaciated Mountains, Upper Clark Fork, Lower Clark Fork, and Blue Mountains ERUs.

2. (To address issue no. 1) Restore dominance of ponderosa pine to sites where transition to other cover types has occurred.
3. (To address issues nos. 1 and 2) Accelerate development of late-seral conditions, including snag recruitment, within stands that are currently in mid-seral stages. Areas for emphasis are the same as those listed for Strategy no. 1.
4. (To address issue no. 2) Include provisions for snag retention and snag recruitment where needed in all management plans involving forests used as source habitats for group 1.
5. (To address issue no. 3) Reduce risk of stand-replacing fires in late-seral ponderosa pine.
6. (To address issue no. 4) Within all ERUs with cottonwood-willow stands, maintain existing old forests and identify younger stands for eventual development of old-forest structural conditions. Return natural hydrologic regimes to large river systems, particularly in the Central Idaho Mountains, Upper Snake, and Snake Headwaters ERUs where large riparian cottonwood woodlands still remain.

Practices that support strategies---The following practices would be effective in implementing the strategies listed above:

1. (In support of strategies no. 1-4) Use understory thinning and prescribed burns to enhance development of ponderosa pine old forests and to reduce fuel loads. Refer to Blair and others (1995) for specific recommendations regarding live tree densities for the old-forest structural stage.
2. (In support of strategy no. 4) Retain existing snags, particularly if >53 cm (21 in), and provide measures for snag replacement. Review existing or develop new snag guidelines that reflect local ecological conditions and that address snag numbers, diameter, height, decay class, species, and distribution.
3. (In support of strategy no. 4) Reduce road densities in managed forests where ponderosa pines snags are currently in low abundance. Close roads after timber harvests and other management activities, and minimize the period when such roads are open, to minimize removal of snags along roads. In addition or as an alternative to road management, actively enforce fuelwood regulations to minimize removal of snags.
4. (In support of strategy no. 4) Restrict fuel wood permits to disallow snag cutting where ponderosa pine snags are in low abundance, and particularly where existing roads cannot be closed. Blair and others (1995) recommend that public fuel wood harvest should be limited to trees <38 cm (15 in) diameter at breast height (d.b.h.).

GROUP 2 - LEWIS' WOODPECKER (MIGRANT POPULATION)

Results

Species ranges, source habitats, and special habitat features--group 2 consists of populations of Lewis' woodpecker that breed but do not overwinter in the basin. Breeding occurs in portions of all ERUs except the Upper Klamath and Northern Great Basin (fig. 6).

Vol. 2, Figure 6--Ranges of species in group 2 within the basin.

Source habitats of Lewis' woodpecker include old-forest, single-storied structural stages of ponderosa pine and multi-storied stages of Douglas-fir, western larch, and riparian cottonwood woodlands (vol. 3, appendix 1, table 1). Unlike most woodpecker species, the Lewis' woodpecker is an aerial insectivore and requires openings for foraging maneuvers. Their breeding distribution is strongly associated with the distribution of ponderosa pine in western North America (see Diem and Zeveloff 1980). This species is often classified as a specialist in burned pine forest habitat, although suitability of burned areas as habitat may differ with postfire age, size, and intensity of burn, and geographic region (Block and Brennan 1987, Bock 1970, Linder 1994, Raphael and White 1984, Saab and Dudley 1998). Burned ponderosa pine forests created by stand-replacing fires seem to be highly productive source habitats compared to unburned pine or cottonwood riparian forest (see Tobalski 1997). Burned-unburned stand condition was not included in the analysis of source habitat extent but is addressed in regards to source habitat quality.

Among nine cavity-nesting species, Lewis' woodpecker was a highly successful nester and the most abundant species nesting in a large (100 000 ha [250,000 acres]), recently burned pine forest in western Idaho (Saab and Dudley 1998). Openings in partially logged, burned forests likely provide greater opportunities for aerial foraging. Within the large burned forests in western Idaho, Lewis' woodpecker nested (1) almost exclusively in salvage-logged units (1.1 nests per km [1.7 per mi] surveyed), compared to unlogged units (0.05 nests per km [0.08 per mi] surveyed); (2) in sites where snags were distributed in clumps; (3) in areas with densities of snags >23 cm (9 in) d.b.h. averaging 59.3 snags per ha (24 snags per acre); and (4) in areas with snag densities for trees >53 cm d.b.h. (21 in) averaging 15.6 snags per ha (6.3 snags per acre) (Saab and Dudley 1998). Nest sites generally are associated with an abundance of flying insects, open-canopy forest or tree clumps, snags, and dense ground cover in the form of shrubs, downed material, and grasses (Bock 1970, Saab and Dudley 1998, Tashiro-Vierling 1994, Tobalski 1997, Vierling 1997). In burned habitats in Wyoming (Linder 1994) and California (Block and Brennan 1987), the percentage of shrub canopy in breeding areas was 13 to 16 percent.

Snags are a special habitat feature for this species (vol. 3, appendix 1, table 2). Lewis' woodpeckers require large snags in an advanced state of decay or trees with soft sapwood for ease of cavity excavation (Bock 1970, Raphael and White 1984, Saab and Dudley 1995). Additionally, Lewis' woodpeckers usurp occupied cavities (Saab and Dudley 1995), reuse old cavities created by strong excavators (for example, hairy woodpecker, black-backed woodpecker, or northern flicker), or nest in natural cavities of trees (Bock 1970, Saab and Dudley 1995, Tashiro-Vierling 1994, Vierling 1997). Reuse of old nests and excavation of highly decayed wood probably are associated with their weak excavation morphology compared to that of other woodpeckers (see Tobalski 1997). Nest tree species are typically ponderosa pine and cottonwood, and less commonly aspen, lodgepole pine, juniper, willow species, and paper birch (Tobalski 1997). Snags and trees used for nesting are generally larger in diameter and more heavily decayed than that expected based on availability of such snags. In burned

ponderosa pine forests of western Idaho, nest trees were large ($\bar{x} \pm SD = 44.5 \pm 1.8$ cm d.b.h. [17.5 \pm 0.7 in]) and were of heavier decay than were trees measured at random ($n = 206$ nests; Saab and Dudley 1998). In Colorado, cottonwood nest trees had a larger d.b.h. (112.6 ± 38.8 cm [44.3 \pm 15.3 in]) than random trees ($n = 47$ nests; Tashiro-Vierling 1994, Vierling 1997). In burned pine-fir forests of the Sierras, nest height averaged 7.3 m (24.0 ft), tree height 11.4 m (37.4 ft), tree d.b.h. 66.5 cm (26.2 in), and tree diameter at cavity 52.2 cm (20.6 in) ($n = 37$ nests; Raphael and White 1984).

Broad-scale changes in source habitats--Historically, the greatest concentrations of Lewis' woodpecker source habitats (excluding burned coniferous forest and riparian habitat that were not considered at the scale of this analysis) were in the Northern Glaciated Mountains, Lower Clark Fork, and Blue Mountains ERUs (fig. 7A). Up to 50 percent of several watersheds within these ERUs are thought to have provided source habitats, whereas lesser amounts of source habitats likely occurred in most watersheds of the Columbia Plateau, Southern Cascades, Upper Clark Fork, Central Idaho Mountains, and Snake Headwaters ERUs (fig. 7A).

The current amount of source habitat is strongly reduced from historical levels in all 11 ERUs that provide source habitat (fig. 7B). The Central Idaho Mountains currently provide the most contiguous habitats, yet these comprise <25 percent of most watersheds (fig. 7B).

Dramatic declines in source habitats seem widespread, based on strong negative trends in 85 percent of the watersheds throughout the basin (figs. 7C and 8). Strong negative trends were particularly evident in the northern watersheds of the basin (Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork ERUs, where more than 95 percent of the watersheds experienced declines (fig. 8). Relative change in extent of source habitats for the Lewis' woodpecker was the greatest (that is, most negative) of any species analyzed in this report (vol. 1, table 7).

Vol. 2, Figure 7--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 2.

Vol. 2, Figure 8--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 2, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Declines in areal extent of source habitats were due primarily to a basin-wide alteration of old-forest ponderosa pine to mid-seral structural stages (Hann and others 1997). The current extent of mid-seral dry forest types is nearly twice the historical level (Hann and others 1997). In the northern and central ERUs, less than 10 percent of the historical extent of interior ponderosa pine in the old-forest single-story structural stage remains (vol. 3, appendix 1, table 4). Late-seral western larch also underwent immense declines and is nearly absent at the broad scale in all ERUs in which it historically occurred (vol. 3, appendix 1, table 4).

Within the cottonwood-willow cover type, old forests have strongly declining trends throughout the basin (see vol. 3, appendix 1, table 4) and generally remain only in stands smaller than the 1-km² (0.4-mi²) mapping unit used in this analysis. These losses occurred from changes in historical hydrologic regimes. Flooding by reservoirs eliminated many cottonwood-willow stands, and reservoirs also reduced periodic flooding, a disturbance that is frequently needed for cottonwood seed establishment (Merigliano 1996, Rood and Heinze-Milne 1989). The declines in riparian woodlands, old-forest ponderosa pine, and western larch documented for the basin are part of a larger picture of similar declines throughout the Western United States (Noss and others 1995).

Condition of special habitat features--Abundance of large (>53 cm [21 in]), heavily decayed snags for nesting have been reduced basin-wide because of changes in vegetation structure from old-forest single stratum to mid-seral structures as well as snag removal by woodcutters (Hann and others 1997; Hessburg and others 1999; Quigley and others 1996). Reductions in the amount of old-forest single stratum and stand initiation structures have reduced forest patch openings that allow foraging maneuvers. In the central and southern regions of the basin, increases in closed-canopy, multi-storied forests have reduced understory shrubs and presumably reduced the abundance of associated arthropods on which Lewis' woodpecker feed.

Other factors affecting the group--Road densities have significantly increased throughout the basin (Hann and others 1997), thereby allowing greater human access into forested regions and greater potential for snag removal along roads. Prolonged human presence at or near nest sites may cause abandonment (Bock 1970), although stable populations coexist with park development and heavy tourist use during the breeding season in British Columbia (Siddle and Davidson 1991). Chlorinated hydrocarbons (such as DDT, a pesticide formerly used in fruit orchards and gardens) could have potential negative effects on Lewis' woodpeckers (Tobalski 1997) because they sometimes nest in agricultural settings (Sorensen 1986, Tashiro-Vierling 1994). Elevated energetic costs and stress may be associated with high rates of territorial encounters with European starlings, which could reduce reproductive success even if Lewis' woodpecker dominates the interaction (Siddle and Davidson 1991).

Population status and trends--Breeding Bird Surveys indicate that population trends have been stable within the basin from 1968 to 1994 (Saab and Rich 1997). Saab and Rich (1997), however, included the Lewis' woodpecker as one of 15 Neotropical migrants in the basin that are of high concern to management under all future management themes for the basin, due to the species close association with old forest stages of ponderosa pine. Populations may have declined by about 60 percent within the Western United States since the 1960s, on the basis of BBS data (1966 to 1995, -4.0 percent per yr, $\bar{n} = 61$, $\underline{P} < 0.01$; Sauer and others 1996). Also, Christmas Bird Counts (CBC) showed a decline in Lewis' woodpecker observations across the entire range of the species, from an average of 10 birds per 1,000 observation hours in 1960 to about four birds per 1,000 observation hours in 1989 ($\bar{n} = 20$, $\underline{P} < 0.05$; Tashiro-Vierling 1994).

Trend data generated by the BBS and CBC may not be adequate for monitoring populations of Lewis' woodpecker (Saab and Rich 1997, Tobalski 1997) because of their sporadic distribution (Bock 1970) and relatively uncommon status (DeSante and Pyle 1986). Dramatic cycles of abundance may be related to local changes in habitat (Bock 1970) and to nomadic behavior of Lewis' woodpeckers in search of burned forests for nesting habitat.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 2 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The following issues were identified from results of our analysis in combination with relevant vegetation dynamics documented by Hann and others (1997):

1. Declines in shrub understories of montane and lower montane forests.
2. Basin-wide decline in old forests of interior and Pacific ponderosa pine and interior western larch.
3. Basin-wide decline in old forests of cottonwood woodlands.
4. Decline in availability of large snags and trees for foraging and nesting.
5. Potential for negative impacts from agricultural pesticides.

Potential strategies--The issues identified above suggest the following broad-scale strategies for the long-term persistence of Lewis' woodpecker.

1. (To address issue no. 1) Rejuvenate and enhance shrub understory of lower montane community groups (old-forest ponderosa pine) and montane community groups that include interior Douglas-fir and western larch in the Northern Glaciated Mountains, Lower Clark Fork, Upper Clark Fork, and Blue Mountains ERUs.
2. (To address issue no. 2) Restore degraded stands and maintain high-quality existing stands of old-forest interior and Pacific ponderosa pine, interior Douglas-fir, western larch, and cottonwood-willow. Protection and restoration of existing old forests is especially important within the range of this species where declines in old forests have been most pronounced. Areas of emphasis include Blue Mountains, Northern Glaciated Mountains, Upper Clark Fork, Lower Clark Fork, and Central Idaho Mountains ERUs. Within these same ERUs, accelerate development of old forests within stands that are currently mid-seral structural stages.
3. (To address issue no. 3) Within all ERUs with cottonwood-willow stands, maintain existing old forests, and identify younger stands for eventual development of old-forest structural conditions. Return natural hydrologic regimes to large river systems, particularly in the Central Idaho Mountains, Upper Snake and Snake Headwaters ERUs where large cottonwood riparian woodlands still remain.
4. (To address issue no. 4) Retain all large-diameter (>53 cm d.b.h.) ponderosa pine, cottonwood, Douglas-fir, and western larch snags within the basin, preferably in clumps, and provide opportunities for snag recruitment.
5. (To address issue no. 5) Reduce exposure to pesticides during nesting season. Avoid use of toxic chlorinated hydrocarbons and organophosphorus insecticides near Lewis' woodpecker nesting sites.

Practices that support strategies---The following practices would be effective in implementing the strategies listed above:

1. (In support of strategies nos. 1 and 2) Use prescribed burns and understory thinning of small-diameter trees (≤ 25 cm d.b.h. [10 in]) to maintain existing old-forest ponderosa pine stands and to accelerate development of midsuccessional stages to old-forest conditions. These practices also can be used to enhance and develop shrub understories (>13 percent shrub canopy) to attract arthropod prey.
2. (In support of strategies nos. 1 and 2) Allow stand-replacing wildfires to burn in lower montane wilderness and other lands managed with a reserve emphasis (for example, designated wilderness, research natural areas, and areas of critical environmental concern). Such opportunities can be found

particularly in the Central Idaho Mountains, Blue Mountains, and Snake Headwaters ERUs, and in western Montana.

3. (In support of strategy no. 4) Develop measures for snag recruitment in unburned forests. Management for snag recruitment (particularly broken-topped snags) in unburned forests with high risks of stand-replacing fires will provide nest trees during the first few years after wildfire when other trees are not easily excavated.
4. (In support of strategy no. 4) In salvage-logged, postfire ponderosa pine forests, retain snags in clumps rather than evenly spaced, leaving both hard and soft decay classes to lengthen the time that those stands are suitable for nesting by Lewis' woodpeckers. Snag densities should approximate 59 snags per ha (24 snags per acre) of d.b.h. size >23 cm [9 in], and of these, about 15 snags per ha (6 snags per acre) should be large snags (\geq 53 cm d.b.h. [21 in]) (Saab and Dudley 1998).
5. (In support of strategy no. 4) Minimize the density of roads open to motorized vehicles. Close roads after timber harvests and other management activities, and maintain short time periods during which such roads are open to minimize removal of snags along roads. In addition or as an alternative to road management, actively enforce fuel wood regulations to minimize removal of snags.
6. (In support of strategy no. 4) Restrict fuel wood permits to disallow snag cutting where ponderosa pine snags are in low abundance, and particularly where existing roads cannot be closed. Blair and others (1995) recommend for Idaho that public fuel wood harvest should be limited to trees <38 cm (15 in) d.b.h.
7. (In support of strategy no. 5) Avoid use of toxic agricultural insecticides near Lewis' woodpecker nest sites.

GROUP 3--WESTERN GRAY SQUIRREL

Results

Species ranges, source habitats, and special habitat features--Group 3 is composed of the western gray squirrel, a year-round resident of the basin. The western gray squirrel is distributed within the western portion of the basin. Its range includes the Southern Cascades, most of the Northern Cascades and Upper Klamath, and portions of the Northern Glaciated Mountains, Columbia Plateau, and Northern Great Basin ERUs (fig. 9). Currently, however, only small, disjunct areas within this range are occupied by squirrel populations (Ryan and Carey 1995).

Vol. 2, Figure 9--Ranges of species in group 3 within the basin.

Source habitats for the western gray squirrel include interior ponderosa pine and Oregon white oak woodlands (vol. 3, appendix 1, table 1). Structural stages of interior ponderosa pine that provide source habitat are old-forest single-story, old-forest multi-story, and both managed and unmanaged young forest.

Mast-producing trees are an important component of western gray squirrel habitat. Species of mast-producing trees differ throughout the range of the squirrel and include both the native Oregon white oak

and introduced English and black walnuts (Barnum 1975). The western gray squirrel uses tree cavities and stick nests as winter dens and for rearing young (Ryan and Carey 1995). The presence of a contiguous tree canopy that allows for arboreal travel around nest sites is also an important habitat feature (ICBEMP 1996c).

Broad-scale changes in source habitats--The trend in broad-scale source habitats for western gray squirrels from historical to current periods was mixed (fig. 10). Moderate or strong decreases were projected in about 30 percent of the watersheds basin-wide, with moderate to strong increases in nearly an equal number (fig. 11). In the Northern Cascades, there were negative and strongly negative trends in about 65 percent of the watersheds (fig. 11). More than half the watersheds in the Northern Great Basin had declining or strongly declining trends. In the Columbia Plateau, there were increasing or strongly increasing trends in about 65 percent of watersheds (fig. 11). Other ERUs either showed mixed trends in source habitats (Southern Cascades, Upper Klamath) or had few watersheds that fell within the range of the squirrel (Blue Mountains, Northern Glaciated Mountains).

Vol. 2, Figure 10--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 3.

Vol. 2, Figure 11--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 3, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Declines in source habitats in the Northern Cascades were due to large decreases in old-forest single-story, old-forest multi-story, and unmanaged young-forest structural stages of interior ponderosa pine (vol. 3, appendix 1, table 4). In the Northern Great Basin, most of the decline resulted from decreases in old-forest single-storied interior ponderosa pine (vol. 3, appendix 1, table 4). Increasing trends in the Columbia Plateau were mostly due to increases in the managed young-forest stage of interior ponderosa pine.

Although oak woodlands were listed as an important source habitat, there was not a measurable vegetation change in this cover type in the ERUs within the range of the species (vol. 3, appendix 1, table 4). In many cases, oak woodlands do not occur in large patches in the basin, and may not have been adequately sampled by the 1 km² (0.4 mi²) pixel size used to interpret vegetation.

Condition of special habitats features--Mast-producing trees, such as oak, likely have declined primarily because of increasing human developments (Washington Department of Wildlife 1993c). In roaded areas with a history of timber harvests, densities of large diameter trees (>53 cm [21 in] d.b.h.) have declined from historical conditions (Hann and others 1997; Hessburg and others 1999; Quigley and others 1996), thus reducing the availability of cavities.

Other factors affecting the group--Introduced eastern fox squirrels and gray squirrels (eastern) are

potential competitors in parts of the range of the western gray squirrel (Ryan and Carey 1995). Humans often shoot western gray squirrels both legally and illegally. In Washington the western gray squirrel is protected from hunting; in Oregon, however, the western gray squirrel is a game species and is regarded as a pest in nut orchards (Ryan and Carey 1995).

Local extirpations caused by mange infestations have seriously affected populations of western gray squirrels. Recovery of populations from disease outbreaks may be difficult when populations are small and widely dispersed (Ryan and Carey 1995).

Population status and trends--Although there is no specific evidence of a reduction in range of western gray squirrels from historical conditions, there is evidence that populations within the range are sparser and more scattered (Washington Department of Wildlife 1993c). This suggests a declining population trend, but there are no direct population data available to confirm the trend.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 3 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--Our results, combined with literature and other empirical information, suggest that the following issues are important for the western gray squirrel:

1. Loss of habitat because of increased human development, timber harvest, and other management activities.
2. Loss or decline of oak trees as a cover type and within other cover types.
3. Isolation of squirrel populations because of loss of habitat.
4. Interspecific competition with nonnative squirrels.
5. Direct mortality because of hunting and illegal shooting.

Potential strategies--Issues for the squirrel suggest that the following strategies may help land managers effectively address declines in habitats or populations within the range of the squirrel in the basin:

1. (To address issue nos. 1 and 2) Across the current range of the squirrel, provide source habitats composed of young- and old-forest interior ponderosa pine stands that include an oak component.
2. (To address issue no. 2) Manage for the maintenance and restoration of oak woodlands.
3. (To address issue no. 3) Provide connectivity among current squirrel populations (Ryan and Carey 1995) by increasing the areal extent of habitats where these have declined, particularly in watersheds within the Northern Cascades, Southern Cascades, and Upper Klamath ERUs.
4. (To address issue nos. 4 and 5) Coordinate with other agencies and parties on cooperative efforts to ensure that habitats and populations are maintained.

Practices that support strategies---The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Where mixed-coniferous/deciduous forest stands have the potential to support a significant oak component, manage them to provide a mixed tree species composition by (1) killing overtopping conifers to allow oaks to grow to an open form; (2) thinning dense pure oak and conifer-oak stands to reduce crowding and water stress and allowing remaining oaks to become larger, more vigorous, more productive, and more fire-resistant; (3) removing smaller conifer trees under the oak canopy that are competing with oaks for water and that will eventually overtop the oaks (Ryan and Carey 1995); and (4) retaining old and large conifers within oaks stands where these trees are widely spaced and have an open crown that intercepts little sunlight while providing good year-round shelter for wildlife and their nests (Ryan and Carey 1995).
2. (In support of strategy no. 2) Manage oak woodlands to achieve the following attributes: (1) large, live, open-form oaks; (2) nearby water; (3) adjacent intergrading stands of ponderosa pine; (4) associated deciduous trees and shrubs; (5) a second age class of closed-form oaks to replace aging oaks; (6) natural prairie plant associations to provide an open to patchy understory; and (7) corridors linking habitat fragments (Ryan and Carey 1995). Minimum size of oak stands should be 2 ha (5 acres), with a desired size of 4 ha (10 acres) (Ryan and Carey 1995).
3. (In support of strategy nos. 2 and 3) Identify and emphasize the location of mature oak stands in relevant management plans, particularly where such stands could potentially link existing populations. Include oak preservation in planning criteria (Ryan and Carey 1995). Increase public awareness of Oregon white oak and western gray squirrels (Ryan and Carey 1995).
4. (In support of strategy no. 4) Improve coordination among state agencies to design hunting seasons to target only areas of crop depredations and to avoid introductions of competitive species.

GROUP 4--BLUE GROUSE (WINTER)

Results

Species ranges and source habitats--This group consists of winter habitat for blue grouse. Blue grouse are widely distributed across the basin, occurring along the crest of the Cascade Range, in the Blue Mountains, and throughout Idaho and western Montana (fig. 12). Spring and summer habitat for blue grouse occurs at lower elevation than winter habitat, and is discussed in group 17. Specific winter source habitats for blue grouse are old-forest single-story, old-forest multi-story, and understory reinitiation stages of interior Douglas-fir, western larch, Sierra Nevada mixed conifer, Pacific ponderosa pine, and interior ponderosa pine; and mixed conifer woodlands (vol. 3, appendix 1, table 1).

Vol. 2, Figure 12--Ranges of species in group 4 within the basin.

Broad-scale changes in source habitats--Significant areas of blue grouse winter range occur in 9 of the 13 ERUs (fig. 12). Within the winter range of the blue grouse, there has been an overall decline in its winter habitat with about 70 percent of watersheds showing a moderate or strong decline (figs. 13 and 14). Moderate or strong declines occurred in source habitat in at least 50 percent of watersheds within eight ERUs that included the Northern Cascades, Southern Cascades, Blue Mountains, Northern Glaciated Mountains, Lower Clark Fork, Upper Clark Fork, Snake Headwaters, and Central Idaho Mountains (figs. 13 and 14). Moderate or strong habitat increases were projected in over 50 percent of

watersheds only in the Upper Klamath. The Northern Great Basin, Columbia Plateau, Owyhee Uplands, and Upper Snake ERUs contain only small areas of blue grouse winter habitat (fig. 13).

Vol. 2, Figure 13--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 4.

Vol. 2, Figure 14--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 4, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Many of the cover types and structural stage combinations estimated to provide source habitats for wintering blue grouse have decreased in area from historical to current periods (Hann and others 1997; vol. 3, appendix 1, table 4). Interior ponderosa pine old-forest single-story stage was the major contributor to declines in habitat in seven of the eight ERUs, with moderate or strong declines (vol. 3, appendix 1, table 4). Other habitats that declined within these ERUs were interior ponderosa pine understory reinitiation and old-forest multi-storied stages, interior Douglas-fir old-forest single- and multi-storied stages, western larch old-forest multi-storied stage, and mixed-conifer woodland (vol. 3, appendix 1, table 4). In the Upper Klamath, the only ERU for which a moderate or strong increase was projected, the largest increases were projected for interior ponderosa pine old-forest multi-storied stage and interior Douglas-fir old-forest single- and multi-storied stages (vol. 3, appendix 1, table 4).

Other factors affecting the group--Blue grouse are sedentary during winter, moving only 69 m (226 ft) per day on average (Cade and Hoffman 1993, Hines 1986). Their sedentary nature makes them vulnerable to various predators such as lynx, red fox, weasels, American marten, merlin, prairie falcon, northern goshawk, and Cooper's hawk (Zwickel 1992). There are, however, no reports of predation seriously depressing blue grouse populations.

Population status and trends--Although blue grouse still occupy most of their original range (fig. 12), accounts suggest higher historical densities in parts of their range (Zwickel 1992). There are, however, no empirical data on population trend for blue grouse within the basin.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 4 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--Our analysis indicates winter habitats for blue grouse have declined in the basin; the following issue could be addressed for this species within overall ecosystem-based strategies:

1. Reduction in the amount of montane and lower montane old forests.

Potential strategies--Blue grouse winter habitat could be improved by strategies that focus on the following:

1. (To address issue no. 1) Retention of existing interior ponderosa pine, interior Douglas-fir, and western larch old forests, with highest priority for retention in watersheds that still support substantial blue grouse winter habitat within ERUs that have shown large decreases in habitat.
2. (To address issue no. 1) Management of early- and mid-seral montane and lower montane forests to accelerate restoration of late-seral conditions of interior ponderosa pine, interior Douglas-fir, and western larch.

Practices that support strategies---The following practice would be effective in implementing the strategies listed above:

1. (In support of strategy nos. 1 and 2) Retain remnant, large trees (Pekins and others 1991) in all seral stages of montane forests. In a Colorado study, Cade and Hoffman (1990) found wintering blue grouse in late-seral Douglas-fir stands as small as 1 ha (2.5 acres). Remington and Hoffman (1996) recommended selective logging that would retain clumps of trees of that size.

GROUP 5--NORTHERN GOSHAWK (SUMMER), FLAMMULATED OWL, AMERICAN MARTEN, AND FISHER

Results

Species ranges, source habitats, and special habitat features--Group 5 consists of the northern goshawk, flammulated owl, American marten, and fisher. Only summer habitat for northern goshawks is included in this group. Goshawk winter habitat is analyzed separately as group 25 because it includes juniper habitats not used by members of this group. Flammulated owls migrate out of the basin in winter, so only their breeding habitat is represented in this group. Goshawks occur throughout forested areas of the basin (fig. 15). Flammulated owls are broadly distributed throughout the Northern Cascades, Northern Glaciated Mountains, Upper and Lower Clark Fork, Blue Mountains, Central Idaho Mountains, and Upper Klamath ERUs. The range of the American marten includes parts of the western, central eastern, and northeastern portions of the basin (fig. 15). Currently the fisher occurs in the western portion of the basin and in central and northern Idaho and western Montana (fig. 15); historically its range included more areas in the northern, central, and eastern portions of the basin (fig. 15).

Vol. 2, Figure 15a, b--Ranges of species in group 5 within the basin.

Source habitats common to all four species are late-seral stages of the montane community group; unmanaged young forests also are source habitats because this structural stage, like late-seral stages, contains sufficient large-diameter snags and logs needed for various life functions of species in the group (vol. 3, appendix 1, table 1). Managed young-forest stages do not provide source habitat because of the lack of remnant large trees and snags. Source habitats for martens extend up into these same stages of subalpine forests, whereas habitats for goshawks and flammulated owls extend down into the same

stages of lower montane forests. For goshawks, flammulated owls, and martens, source habitat also is provided by the old-forest multi-storied and unmanaged young-forest stages of aspen, whereas goshawks, flammulated owls, and fisher find source habitat in these same stages of cottonwood-willow. In addition, flammulated owls use limber pine (McCallum and Gehlbach 1988) and mixed-conifer woodlands as source habitats, and goshawks use chokecherry-serviceberry-rose as source habitats.

Goshawks nest in various forest structural conditions, from open, parklike stands of aspen (Younk and Bechard 1994), to multi-storied old forests (Reynolds 1983). Nest stands are generally characterized by large trees and the densest canopy cover available within the area (Reynolds and others 1992) but are occasionally located in small-diameter trees (Hayward and Escano 1989, Squires and Ruggiero 1996). Foraging occurs in various cover types and structural stages, and the juxtaposition of several habitats may enhance the quality of foraging habitat around nest sites (Hargis and others 1994). Home range for a nesting pair is estimated at >2400 ha (5,930 acres) (Hargis and others 1994, Kennedy and others 1994, Reynolds and others 1992).

Martens seem more sensitive to patch size than other group members, and usually avoid clearcuts dominated by grasses, forbs, and saplings, especially in winter. These areas do not provide access to the subnivean zone or offer protection from predation, and have more severe microclimatic conditions than areas with forest cover (Buskirk and Powell 1994). At the broad scale, the presence of multiple clearcuts may render the entire landscape unsuitable. In Utah, martens were rarely found in areas with >25 percent of the landscape in a combination of natural openings and clearcuts (Hargis 1996). In Maine, no adult female territories were found in landscapes with >31 percent of mature forest cover removed (Chapin 1995).

Although fishers will cross openings to access forested areas (Arthur and others 1989), a negative association with clearcuts has been documented. Fisher occurrence in California was positively associated with large stands of mature forest and distance from clearcuts (Rosenburg and Raphael 1986); fishers in Idaho avoided stands with <40 percent canopy cover (Jones 1991, Jones and Garton 1994).

Old forests consisting of ponderosa pine and Douglas-fir seem to be a key component of flammulated owl home ranges (Reynolds and Linkhart 1992). Home ranges composed of at least 75 percent old ponderosa pine/Douglas-fir forest were occupied more continuously than home ranges consisting of less than 75 percent in this forest type (Reynolds and Linkhart 1990). Variability in the structure of these old stands seems important to support life functions of flammulated owls. Roosting occurs in fairly dense stands. Goggans (1986) showed that tree densities immediately surrounding roost trees averaged 2016 per ha (816 per acre), whereas overall home ranges averaged 589 trees per ha (238 per acre). In contrast, relatively open stands seem to be selected for foraging (Linkhart 1984), and open, mature stands are selected for nest sites (McCallum 1994). In two Oregon studies, mean d.b.h. of nest trees was 56.3 cm (22.2 in) (Goggans 1986) and 72.0 cm (28.4 in) (Bull and others 1990).

Several special habitat features have been identified for this group (see vol. 3, appendix 1, table 2). Fishers and American martens use down logs. Downed woody material is likely the key component of foraging areas for marten (Coffin and others 1997), providing habitat for many of their prey, particularly southern red-backed voles, and subnivean access to prey during winter (Corn and Raphael 1992). Fishers and martens depend on down logs for resting and denning (Buskirk and Powell 1994, Raphael and Jones 1997). Snags are a special habitat feature for flammulated owls, fishers, and martens. Flammulated owls nest in cavities in both snags and large live trees (Bull and others 1990, McCallum and Gehlbach 1988). Snags provide rest sites and den sites for fishers and martens.

Broad-scale changes in source habitats--Historically, source habitats likely occurred throughout the forested portions of the basin, with some of the greatest concentrations in the western, central, and

northern portions of the basin (fig. 16A). Currently, the largest extent of source habitats is in the south-central and southwestern portions of the basin (fig. 16B). The primary change from historical to current times has been a broad shift in the geographic distribution of source habitats away from the north and towards the southwestern portion of the basin (fig. 16C).

Basin-wide, there were moderately or strongly declining habitat trends in nearly 70 percent of watersheds within the range of species in group 5, and moderately or strongly increasing trends in about 33 percent of watersheds (fig. 17).

Positive changes in source habitat occurred in more than 50 percent of watersheds in the Upper Klamath and Northern Great Basin ERUs; mixed trends in the Southern Cascades and Upper Snake ERUs; and negative trends in more than 50 percent of watersheds in all remaining ERUs (figs. 16 and 17). The most strongly negative trends were projected across the northern portion of the basin in the Northern Cascades, Northern Glaciated Mountains, Lower Clark Fork and Upper Clark Fork ERUs (figs. 16 and 17).

Vol. 2, Figure 16--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 5.

Vol. 2, Figure 17--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 5, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Interior ponderosa pine old-forest single-story stage declined in all but one of the ERUs in which source habitat declined in more than 50 percent of watersheds (vol. 3, appendix 1, table 4). Interior ponderosa pine old-forest multi-story stage declined in nearly half of these ERUs. Less consistent declines were projected for the old-forest single-story stage of interior Douglas-fir; the old-forest multi-story stages of interior Douglas-fir, lodgepole, grand fir-white fir, Engelmann spruce-subalpine fir, western larch, and western white pine; the unmanaged young forest stages of whitebark pine, Engelmann spruce-subalpine fir, western larch, and lodgepole pine; and mixed conifer woodland (vol. 3, appendix 1, table 4). In the ERUs with the most strongly negative trends, the Northern Cascades, Northern Glaciated Mountains, Lower Clark Fork and Upper Clark Fork, negative trends were projected for up to nine of these habitat types (vol. 3, appendix 1, table 4). In the Upper Klamath, the only ERU with a significant amount of source habitat for the group and a positive trend in more than 50 percent of watersheds, the increasing trend was associated with increases in the old-forest multi-story stages of interior ponderosa pine, interior Douglas-fir, lodgepole pine, and grand fir-white fir; and the old-forest single-story stage of interior Douglas-fir. In addition, riparian woodland (including aspen and cottonwood-willow) declined basin-wide, and also underwent a shift from early- and late-seral stages to mid-seral stages (Hann and others 1997).

Condition of special habitat features--Densities of large-diameter snags (>53 cm [21 in] d.b.h.) declined basin-wide from historical to current levels (Hann and others 1997; Hessburg and others 1999; Quigley

and others 1996). Trends in snag abundance ultimately affect the availability of large down logs and cavities.

Other factors affecting the group--Populations of martens and fishers can be impacted by fur harvesting if trapping is not carefully regulated (Fortin and Cantin 1994, Jones 1991, Quick 1956). Trapping also affects populations by altering the sex and age structure through the disproportionate capture of juveniles and males (Hodgman and others 1994, Quick 1956). Historically, both marten and fisher were heavily trapped in the basin. Currently, martens are still trapped in all states in the basin, but fishers are only trapped in Montana (Heinemeyer 1995).

Secondary roads in forested areas increase trapping pressures for martens and fishers, resulting in significantly higher captures in roaded versus unroaded areas (Hodgman and others 1994) and in logged versus unlogged areas, in which the difference was again attributed to higher road densities in logged stands (Thompson 1994). Secondary roads also might increase the likelihood that snags and logs will be removed for fuel wood. This could impact fisher, marten, and flammulated owls, and also could have a negative effect on the prey base for goshawk (Reynolds and others 1992).

Studies have shown that fisher, marten, and goshawk populations respond to food limitation. Fisher populations can undergo fluctuations related to prey abundance (Powell and Zielinski 1994). Marten populations also have been observed to decline after a decline in principal prey species (Thompson and Colgan 1987, Weckwerth and Hawley 1962). Some of the decline is due to lower reproductive rates in females, but evidence of starvation also has been observed (Hodgman and others 1994, Weckwerth and Hawley 1962). Several studies suggest that goshawk populations are frequently food-limited. In Alaska and the Yukon where snowshoe hare is a dominant prey item, goshawk numbers fluctuate with snowshoe hare cycles (Doyle and Smith 1994). A review of several studies by Widen (1989) suggests correlations between goshawk numbers and other prey. Maj and others (1995) suggest that heavy levels of grazing in ponderosa pine communities may degrade insect habitat and reduce prey populations for flammulated owls.

Changes in forest structure related to fire suppression seem to increase the extent of some of the cover types and structural stages judged to be source habitats for goshawks. However, such stands, which are characterized by closed canopies and dense conifer understory, may not be as valuable to goshawks as the more open habitats, which they replaced. A high density of small-diameter understory trees may be detrimental to foraging and nesting aspects of goshawk ecology in at least three ways: (1) by obstructing flight corridors used by goshawks to obtain forest-associated prey; (2) by suppressing tree growth needed to produce large-diameter trees for nest sites; and (3) by reducing the growth of an herbaceous understory that supports potential prey species (Reynolds and others 1992). Therefore, although fire suppression may have increased the extent of multi-storied closed forests within the basin, the inherent value of these stands may be less than that of more open stands maintained by fire. This supposition warrants further investigation.

Conversely, the harvest of large-diameter overstory trees can create forest structures that are more open than normally used by goshawks. A secondary effect is increased competition with raptors adapted to more open habitats (Moore and Henny 1983). Goshawk nest sites are more frequently used by red-tailed hawks, great horned owls, or long-eared owls in harvested areas than in unharvested sites (Crocker-Bedford 1990, Patla 1990).

Flammulated owls are Neotropical migrants, so their population status may be affected by conditions of their winter habitat. Their winter range is suspected to be in southern Mexico and northern Central America (McCallum 1994).

Population status and trends--Fishers may be close to extirpation in Washington (Aubrey and Houston

1992, cited in Powell and Zielinski 1994), and sightings are rare in Oregon. The last reliable reports of native fishers in Idaho and Montana were during the 1920s (Dodge 1977; Weckwerth and Wright 1968, cited in Powell and Zielinski 1994). Fisher populations were reintroduced to Idaho in the 1960s and to Montana in the 1950s and 1980s (Powell and Zielinski 1994). Projected declines in source habitats may have contributed to historical extirpations, coupled with the effects of trapping and the fragmented nature of remaining habitats.

The distribution of marten within the basin has been fairly stable since historical times, but population changes are not known, other than through trapping records, which fluctuate widely with fur prices and may not reflect actual population trends.

The BBS data for the goshawk were insufficient to determine population trends for the basin (Saab and Rich 1997) or for any state or physiographic region within the basin (Sauer and others 1996) because of low detection of goshawks under the BBS survey method. Sufficient data were available, however, for western North America to indicate a stable trend in numbers between the years 1966 and 1995 (Sauer and others 1996).

A separate trend estimate was derived from fall migration counts conducted by Hawkwatch International at four locations in Utah and New Mexico. These data indicated an average rate of decline in migrating goshawks of about 4 percent annually between 1977 and 1991 (Hoffman and others 1992). The extent to which the migration data represented local declines near the survey stations was not determined.

No population trend data were found for flammulated owls. The BBS survey method is not adequate for surveying flammulated owls because of low numbers and nocturnal behavior. Specialized monitoring would be required to determine the population trend of owls (Saab and Rich 1997).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 5 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The following issues were identified from the results of our analysis and published research:

1. Reduction in the amount of old-forests and associated structures (snags, logs, and cavities), particularly within the montane and lower montane community groups.
2. Fragmentation of habitat.
3. Low population numbers of fisher.
4. Negative effects resulting from higher road densities in source habitats. For marten throughout the basin and fishers in Montana, there is increased trapping pressure associated with roads. For all species in the group, loss of snags and logs associated with firewood collection may be higher along open roads.
5. Declines in overall extent of aspen and cottonwood-willow, and shifts from early- and late-seral to mid-seral stages of these cover types (vol. 3, appendix 1, table 4).
6. Possibly unsustainable conditions of old forests where there have been large transitions from shade-intolerant to shade-tolerant tree species. This last issue stems from the exclusion of fire from many

forested communities, which has resulted in increased susceptibility to stand-replacing fires (USDA Forest Service 1996).

7. Decline in suitable foraging areas around goshawk nest sites. On Federal lands, the immediate areas around active nests generally are protected from timber harvests, but the larger foraging areas surrounding nests frequently are managed without explicit consideration of goshawk foraging. Goshawks typically use a nest stand and nearby alternative nest stands for many years, and therefore, the long-term maintenance of suitable foraging areas is as important for successful reproduction as protection of the immediate nest stand.

Potential strategies--The following strategies could be used to reverse broad-scale declines in source habitats and populations:

1. (To address issue no. 1) Increase the representation of late-seral forests in all cover types used as source habitats, particularly in the northern half of the basin (Northern Cascades, Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork ERUs).
2. (To address issue no. 2) Increase connectivity of disjunct habitat patches and prevent further reduction of large blocks of contiguous habitat.
3. (To address issue nos. 3 and 4) Identify potential species strongholds for long-term management of marten and fisher (see practice no. 6 for criteria).
4. (To address issue no. 4) Reduce human disturbances in source habitats.
5. (To address issue no. 5) Restore aspen and cottonwood-willow forests, particularly the unmanaged young forest and late seral stages.
6. (To address issue no. 6) Reduce the risk of loss of habitat by focusing old-forest retention and restoration efforts on areas where fire regimes are either nonlethal or mixed (USDA Forest Service 1996). In ERUs where old-forest habitat has remained stable or increased from historical conditions, efforts could be focused on retaining existing habitat in areas with lower fire and insect risk while managing other areas to reduce risks of catastrophic loss of habitat.
7. (To address issue no. 7) Maintain stands with active goshawk nests in old-forest condition.
8. (To address issue no. 7) Embed the conservation of old forests within a larger, ecosystem context that considers historical fire regimes and landscape patterns and the habitat needs of species that are prey of the members of this group. For goshawks, Reynolds and others (1992) gave specific recommendations for promoting a variety of cover types and structural stages in 2430 ha (6,005 acres) of potential home range around each active nest.

Practices that support strategies---The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) In the northern basin, identify representative stands of old forests for retention and mid-successional stages for development into old-forest conditions. Priority should be given to large blocks having high interior-to-edge ratios and few large openings.
2. (In support of strategy no. 1) Actively recruit snags and logs from green trees to increase the representation of old forest structures (snags and logs) in mid-seral stands and in old forests where snags and logs are in low density or absent.

3. (In support of strategy no. 1) Retain slash piles and decks of cull logs to substitute for down logs over the short term. Raphael and Jones (1997) recommended retention of a minimum of 1.3 slash piles per ha (0.5 per acre) on a site that had been extensively harvested.
4. (In support of strategy no. 2) Where possible, use selection harvest rather than clearcutting. If clearcuts are used, aggregate cuts so that large blocks of unharvested forest are retained.
5. (In support of strategy no. 2) Adjust activities, including timber harvests, to provide links among currently isolated patches of source habitats.
6. (In support of strategy no. 3) Identify existing areas with the following desired conditions, or manage selected areas to create the following desired conditions for strongholds: existing populations of marten or fisher, or both; large, contiguous blocks of forest cover with a high percentage of late-seral stages, abundant snags and large logs, low road densities and overall low human disturbance, and potential connectivity to currently unoccupied source habitats.
7. (In support of strategy no. 4) Minimize new construction of secondary roads and close unneeded roads after timber harvest.
8. (In support of strategy no. 5) Use clearcutting to regenerate aspen. Where aspen regeneration is inhibited by domestic or wild ungulate browsing, use exclosures to protect regenerating stands and modify management to reduce browsing pressure.
9. (In support of strategy no. 5) Survey and map existing old forests of cottonwoods and reference their locations in land management planning documents. Monitor conditions of cottonwood stands to ensure that sufficient seedling or vegetative regeneration, or both, is occurring. Identify factors limiting regeneration so that appropriate corrective measures can be taken. For example, return natural hydrologic regimes to portions of large river systems that support cottonwood riparian woodlands.
10. (In support of strategy no. 6) Manage risks of catastrophic loss by using prescribed fire and thinning to reduce fuel loading and to encourage the development of forest openings, shrub openings, and shade-intolerant and fire-, insect-, and disease-resistant tree species.
11. (In support of strategy no. 7) Identify an area around each active goshawk nest site to be maintained in old-forest condition, and identify possible replacement stands. The Northern Goshawk Scientific Committee for the FS recommends three 12-ha (30-acre) nest stands per breeding pair and three additional 12-ha (30-acre) replacement stands be located within a 2430-ha (6,000-acre) area that functions as a potential home range (Reynolds and others 1992).
12. (In support of strategy nos. 6 and 8) Use silvicultural prescriptions in conjunction with restoration of fire regimes to create a desired mix of cover types and structural stages within the potential home range of each active goshawk nest. The Northern Goshawk Scientific Committee for the FS (Reynolds and others 1992) has identified two larger habitat use areas that extend beyond the nest site: a postfledgling-family area, encompassing about 170 ha (420 acres) around the nest and used by a nesting pair and offspring from the time the young leave the nest until they are independent, and a foraging area of about 2190 ha (5,411 acres) that provides the food resource during and after the breeding period (Reynolds and others 1992). For forests in the southwestern United States, they recommended that four-fifths of each post-fledgling family area and each foraging area be equally divided among four seral stages: young, mid aged, mature, and old forests, and the remaining one-fifth be equally divided between the seedling-sapling stage and grass-forb stage. These recommendations should be reviewed in light of different ecological conditions within the basin.

GROUP 6 - VAUX'S SWIFT, WILLIAMSON'S SAPSUCKER, PILEATED WOODPECKER, HAMMOND'S FLYCATCHER, CHESTNUT-BACKED CHICKADEE, BROWN CREEPER, WINTER WREN, GOLDEN-CROWNED KINGLET, VARIED THRUSH, SILVER-HAIRED BAT, AND HOARY BAT

Results

Species ranges, source habitats, and special habitat features--Group 6 consists of migratory breeding habitat for brown creepers, Hammond's flycatchers, Vaux's swifts, and Williamson's sapsuckers; resident summer habitat for varied thrushes, winter wrens, silver-haired bats, and hoary bats; and year-round habitat for chestnut-backed chickadees, golden-crowned kinglets, and pileated woodpeckers. Ranges within the basin for the 11 species in this group (fig. 18) tend to fit one of four broad patterns. Silver-haired bats and hoary bats occur throughout the basin in forested areas or woodlands. Brown creepers, Hammond's flycatchers, winter wrens, and golden-crowned kinglets generally occur throughout the forested areas of the basin. The range of Williamson's sapsucker differs from these four species as it does not extend all the way to the crest of the Cascade Range or to the southern extremes of the Central Idaho Mountains or Upper Klamath ERUs. Pileated woodpeckers, varied thrushes, chestnut-backed chickadees, and Vaux's swifts are distributed across forested areas in the western half of the basin, but their ranges do not extend to the southeastern portion of the Central Idaho Mountains below the Salmon River, or into the Snake Headwaters or Upper Snake ERUs.

Vol. 2, Figure 18a, b, c--Ranges of species in group 6 within the basin.

Source habitats for the 11 species in group 6 are generally late-seral stages of the subalpine, montane, lower montane, and riparian woodland community groups (vol. 3, appendix 1, table 1). Source habitats shared in common by more than one-half of the species are the old-forest single- and multistrata stages of grand fir-white fir, interior Douglas-fir, western larch, western white pine, western red cedar-western hemlock, Sierra Nevada mixed conifer, and mountain hemlock; and the old-forest multistrata stage of Engelmann spruce-subalpine fir, Pacific silver fir-mountain hemlock, and red fir (vol. 3, appendix 1, table 1). Source habitats used by less than one-half the species include old-forest Pacific and interior ponderosa pine (used by brown creepers, Hammond's flycatchers, Williamson's sapsuckers, hoary bats, and silver-haired bats); old-forest whitebark pine and alpine larch (used by golden-crowned kinglets); old-forest lodgepole pine (used by golden-crowned kinglets, Hammond's flycatchers, hoary bats, and silver-haired bats); old-forest aspen (used by Williamson's sapsuckers, chestnut-backed chickadees, Hammond's flycatchers, hoary bats, and silver-haired bats); and old-forest cottonwood-willow (used by Williamson's sapsuckers, hoary bats, and silver-haired bats) (vol. 3, appendix 1, table 1). Hoary bats also use the stand initiation stage of all montane and lower montane forest types and of aspen and cottonwood-willow (vol. 3, appendix 1, table 1).

Several special habitat features exist for species in this group (vol. 3, appendix 1, table 2). Six of the bird species (brown creepers, chestnut-backed chickadees, pileated woodpeckers, Vaux's swifts, Williamson's sapsuckers, and winter wrens) depend on snags for nesting or roosting, or both (Bull and Hohmann 1993; Bull and others 1986a, 1992; Raphael and White 1984). Brown creepers, pileated woodpeckers, Vaux's swifts, and Williamson's sapsuckers use large (>53 cm [21 in] d.b.h.) snags (Bull and others 1986a, 1992; Bull and Hohmann 1993, Raphael and White 1984). Winter wrens and chestnut-backed chickadees use smaller diameter snags (Thomas and others 1979). Pileated woodpeckers forage on large snags and logs (Bull and Holthausen 1993, Mannan 1984), and winter wrens forage around and

under logs (Van Horne and Bader 1990). Pileated woodpeckers and Vaux's swifts depend on large, hollow live or dead trees for roosting (Bull 1991, Bull and others 1992).

Special habitat features for both bat species include shrub/herbaceous wetland/riparian areas (vol. 3, appendix 1, table 2). Both species use contrasting habitats; forested areas for roosting and open areas for foraging. Snags are a special habitat feature for silver-haired bats. They roost in trees, snags, mines, caves, crevices, and buildings (Christy and West 1993). Day roost trees are usually characterized by being large (>53 cm [21 in] d.b.h.), dead or live with some defect, with loose bark and cracks. In an Oregon study, Betts (1996) found silver-haired bats roosting in live western larch and ponderosa pine, and in grand fir and ponderosa pine snags. The average diameter of these roost trees was 59.6 cm (23.5 in), and they were generally located on relatively densely forested slopes. The hoary bat is an edge-associated species, often roosting in deciduous trees or conifers at the edge of clearings (Perkins and Cross 1988, Shump and Shump 1982). Hoary bats are foliage roosters, with males, nonbreeding females, and breeding females located in different levels in the canopy (Christy and West 1993).

Broad-scale changes in source habitats--Source habitats for species in this group occur in all 13 ERUs (fig. 19), but amounts of habitat are relatively small in the Northern Great Basin, Owyhee Uplands, and Upper Snake ERUs. Basin-wide, source habitats for species in this group have declined moderately or strongly in more than 50 percent of watersheds containing appropriate habitat types (fig. 20). The pattern of habitat change, however, was highly variable across the basin with the northern part of the basin marked by generally strong declines and the southern part by strong increases (fig. 19). Moderate or strong declines in habitat from historical to current were projected in more than 50 percent of the watersheds in six ERUs: the Northern Cascades, Northern Glaciated Mountains, Lower Clark Fork, Upper Clark Fork, Upper Snake, and Snake Headwaters (fig. 20). The declines were particularly strong across the northern basin in the Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork ERUs. Moderate or strong increases were projected in more than 50 percent of watersheds in the Southern Cascades, Upper Klamath, Northern Great Basin, and Columbia Plateau (fig. 20). More balanced mixes of increases and decreases were projected for the remaining three ERUs: Blue Mountains, Owyhee Uplands, and Central Idaho Mountains (fig. 20).

Vol. 2, Figure 19--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 6.

Vol. 2, Figure 20--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 6, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--The projected decline in source habitats reflected basin-wide declines in late-seral forest conditions (USDA Forest Service 1996). Changes in late-seral forests, however, have varied among ERUs (tables 3.141 to 3.165 in Hann and others 1997). Late-seral lower montane multi-layer forests and late-seral subalpine multi-layer forests declined significantly in all six ERUs in which source habitats declined in more than 50

percent of watersheds; late-seral montane multi-layer forests declined in five of them; and late-seral lower montane single-layer forests declined in four of them (Hann and others 1997).

Late-seral montane multi-layer and single-layer forests each increased significantly in three of the four ERUs (Southern Cascades, Upper Klamath, Northern Great Basin and Columbia Plateau) in which source habitats increased in more than 50 percent of watersheds. Much of this change was due to shifts from shade-intolerant, late-seral lower montane forest types to shade-tolerant, late-seral montane forest types. The increase in the fourth ERU, the Columbia Plateau, appears to be somewhat anomalous. It was likely the result of a moderate increase in the open canopy stem exclusion stage of interior ponderosa pine (vol. 3, appendix 1, table 4), which serves as source habitat only for hoary bats.

Condition of special habitat features--Snags are a special habitat feature for seven of the species in this group, and large hollow trees for two species. Densities of large-diameter (>53 cm [21 in] d.b.h.) snags likely have declined basin-wide from historical to current levels (Hann and others 1997; Hessburg and others 1999; Quigley and others 1996). Historical to current trends in smaller diameter snags were variable, with no clear basin-wide trend emerging (Hann and others 1997).

The number of caves available for roosts across the basin likely has stayed the same, and mines may now provide additional roost or hibernacula areas. Cave and mine suitability, however, can be affected by recreational use, such as cave exploration, which increases with higher road densities near caves. Historical road densities were lower than current densities. Road densities are high in intensively managed forest lands of both public and private ownership, and the highest densities typically occur in developed urban-rural areas (USDA Forest Service 1996, p. 85).

Across the basin, there were widespread declines in shrublands in riparian zones (USDA Forest Service 1996, p. 101). Forest conversion and streamside disturbances have degraded and fragmented riparian vegetation. This may have negatively impacted the shrub/herbaceous wetland/riparian foraging areas for the bats.

Other factors affecting the group--Four of the species in this group (brown creepers, Hammond's flycatchers, Vaux's swifts, and Williamson's sapsuckers) are Neotropical migrants and may be affected by habitat conditions on their wintering grounds. The bat species also are thought to winter outside the basin, although exact migration routes and winter ranges are not clear (Christy and West 1993).

Hoary bats eat moths, beetles, and mosquitos (Barclay 1985, 1986; Rolseth and others 1994; Shump and Shump 1982; Whitaker and others 1977). The silver-haired bat is an opportunistic feeder, and eats moths, flies, beetles and various other insects (Whitaker and others 1981). Management activities such as the use of pesticides that cause declines of insect species may negatively affect these bats. Also, direct contact with pesticides can cause illness or death in bats. Although most organochlorine pesticides that cause accumulation of chemicals up the food chain have been banned or highly restricted in the United States, the relatively short-lived organophosphates can provide high risks during application (Clark 1988). For example, a large die-off of bats was observed in Arizona after the application of methyl parathion, and was believed to be linked to direct contact with this chemical (Clark 1988).

Grazing can have an adverse impact on the insect prey of bats (Clark 1988, Nagorsen and Brigham 1993, Perlmeier 1995, Ports and Bradley 1996). The presence of roads also may facilitate harvest of snags for firewood and so may have an indirect effect on habitat for the species that use snags.

Population status and trends--Saab and Rich (1997) reported stable population trends, based on data from BBS routes within the basin, for Williamson's sapsuckers, Vaux's swifts, Hammond's flycatchers, brown creepers, and golden-crowned kinglets. BBS data analyzed within other geographic boundaries (Sauer and others 1996), however, indicates a significant decline from 1966 to 1994 for brown creepers

in eastern Oregon and Washington (-7.4 percent per year, $\bar{n} = 15$, $P < 0.01$). BBS data also indicate a significant increase in pileated woodpeckers in northwestern Montana (6.1 percent per year, $\bar{n} = 41$, $P < 0.01$, 1966 to 1994; Sauer and others 1996) but a significant decrease in eastern Oregon and Washington (-7.8 percent per year, $\bar{n} = 8$, $P < 0.05$, 1966 to 1979; Sauer and others 1996). A significant increase is shown for winter wrens in eastern Oregon and Washington (7.8 percent per year, $\bar{n} = 9$, $P < 0.05$, 1966 to 1979). Population data are not available for the bat species.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 6 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The following issues were identified from our analysis of source habitat trends:

1. Reductions in the extent of late-seral lower montane, montane, and subalpine forest (Hann and others 1997), particularly in the Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork ERUs.
2. Reductions in large snags and logs in landscapes that have been managed under traditional silvicultural practices (Hann and others 1997).
3. Possibly unsustainable conditions in late-seral stage montane forests where there have been large transitions from shade-intolerant to shade-tolerant species.
4. Degradation and loss of riparian habitat.
5. Abandonment of bat roosts because of human disturbance.
6. Reductions in the insect prey base for bats because of both land management activities and to the use of pesticides.
7. Bat population declines because of use of pesticides and insecticides.

Potential strategies--The following strategies would benefit species in group 6:

1. (To address issue nos. 1 and 2) Accelerate development of late-seral conditions in lower montane, montane, and subalpine forest types and retain large snags and logs in all forest seral stages. Habitat restoration efforts would be most beneficial if concentrated in the northern portions of the basin.
2. (To address issues nos. 1, 2, and 3) In the southern portion of the basin, retain sufficient habitat to support species in this group while restoring forest conditions that are more resistant to catastrophic fire, insect, and disease problems. This could require management activities, including prescribed fire, that reduce the dominance of shade-tolerant tree species and increase the presence of shade-intolerant species (i.e., those most resistant to catastrophic fire and insect and disease problems).
3. (To address issue no. 4) Across the basin, maintain or improve riparian shrubland and riparian woodland communities.
4. (To address issues nos. 2 and 5) Protect known and potential bat roosts across the basin. Specifically, maintain caves, mines, snags, and other such features for use as roosting areas and

potential nurseries across the basin. Minimize human disturbance in these areas.

5. (To address issues nos. 6 and 7) Minimize direct physiological effects on bats, as well as indirect effects on their insect prey, stemming from use of insecticides and pesticides.
6. (To address issue no. 6) Modify management practices as appropriate to enhance the insect prey base for bats.

Practices that support strategies---The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Various silvicultural practices including thinning, burning, and uneven-age management could be used to help accelerate the development of old-forest conditions.
2. (In support of strategy nos. 1 and 2) Both the retention and creation of snags are important to retention and development of old-forest characteristics. Techniques for snag management are well studied (Bull and others 1980, Bull and Partridge 1986) and have been extensively applied on National Forests (Bull and others 1986b). Retain existing snags, particularly if > 53 cm (21 in) and provide measures for snag replacement. Review existing snag guidelines or develop guidelines that reflect local ecological conditions and address snag numbers, diameter, height, decay class, species, and distribution. Consider closing roads in areas that are snag deficient and where cutting of snags or remnant trees for firewood contributes to the low snag densities. In addition or as an alternative to road management, actively enforce fuel wood regulations to minimize removal of snags.
3. (In support of strategy no. 2) To continue meeting habitat needs of species in this group, habitat retention efforts should be designed to maintain an appropriate network of old-forest habitats. Bull and Holthausen (1993) suggested managing areas of 1000 ha (2,471 acres) to meet needs of multiple pairs of pileated woodpeckers. Features of these areas were to include a substantial old forest and unlogged component, at least 8 snags per ha (3 snags per acre) with at least 20 percent of these >51 cm (20 in) d.b.h., and at least 100 logs per ha (40 logs per acre) with a preference for logs 38 cm (15 in) in diameter and larger. Such strategies could be coordinated with needs for ecosystem health by focusing old-forest retention areas in geographic locations where fire, insect, and disease risks are lowest.
4. (In support of strategy no. 3) Maintain or restore riparian vegetation around permanent and seasonal water sources.
5. (In support of strategy no. 4) Protect building roost sites. If possible, stabilize old structures that are important roosts.
6. (In support of strategy no. 6) Modify grazing practices to improve condition of degraded riparian areas for bat foraging.

GROUP 7--BOREAL OWL

Results

Species ranges, source habitats, and special habitat features--Group 7 consists of the boreal owl. Within the basin, this species occurs in forested portions of eastern Washington, northern and central Idaho, western Montana, and the Blue Mountains and Cascade Range of Oregon (fig. 21). The boreal owl is a

year-round resident of the basin.

Vol. 2, Figure 21--Ranges of species in group 7 within the basin.

Source habitats for boreal owls include old-forest and unmanaged young forest stages of subalpine and montane forests and riparian woodlands (vol. 3, appendix 1, table 1). Specific cover types and structural stages that provide source habitat are the old-forest multi-story stages of Engelmann spruce-subalpine fir, Pacific silver fir-mountain hemlock, and aspen; and the old forest single- and multi-forest stages of interior Douglas-fir, western larch, and lodgepole pine. Unmanaged young-forest stages of all these cover types and of grand fir-white fir also serve as source habitats if suitable large-diameter snags are present. Source habitats typically support abundant lichens and fungal sporocarps, which provide important foods for southern red-backed voles, the principal prey of boreal owls (Hayward 1994c). These lichens and fungi are associated with coarse woody debris.

Boreal owls require snags or large trees with either natural cavities or cavities excavated by other species (vol. 3, appendix 1, table 2). Cavities excavated by pileated woodpeckers and northern flickers are the most common nest sites (Hayward 1994c). Tree and snag diameters used for nesting are generally large. For example, in Idaho, diameters of nest trees ranged from 26 to 61 cm (10 to 24 in) with an average of 41 cm (16 in). Of 19 nests, 10 were in snags while the remainder were in live trees (Hayward and others 1993).

At the home range scale, boreal owls are adapted to patchy landscapes and use several cover types and structural stages to meet different life history requirements (Hayward and others 1993). Landscapes that contain various old-forest cover types may support the greatest abundance of boreals (Hayward and others 1993). In portions of their range, boreal owls may occur in a patchy geographic pattern resulting in a metapopulation structure, with the long-term persistence of each population determined in part by its relation to other populations (Hayward 1994a).

Broad-scale changes in source habitats--Historically, the most concentrated areas of source habitat for boreal owls were in the Northern Cascades, Northern Glaciated Mountains, and Snake Headwaters ERUs (fig. 22A). Other ERUs that historically supported significant source habitat were the Southern Cascades, Lower Clark Fork, Upper Clark Fork, and Central Idaho Mountains ERUs.

Overall, source habitats were projected to have declined moderately or strongly in nearly 80 percent of the watersheds in the basin (fig. 23). Moderate or strong declines were projected for over 50 percent of watersheds in the Northern Cascades, Northern Glaciated Mountains, Upper Clark Fork, Lower Clark Fork, Snake Headwaters, and Central Idaho Mountains ERUs (fig. 23). Moderate or strong declines in over 50 percent of watersheds also were projected for the Columbia Plateau and Upper Snake, but these ERUs are peripheral to the range of boreal owls. Source habitats were projected to have increased moderately or strongly in over 50 percent of watersheds in the Southern Cascades, and there was a mixed pattern of change in the Blue Mountains ERU (fig. 23).

These trends have resulted in a broad shift in the geographic distribution of source habitats away from the northern ERUs and towards the central portions of the basin. Habitat losses have outweighed the gains, and current habitat distribution is substantially more disjunct than historically in the northern part of the basin (fig. 22).

Vol. 2, Figure 22--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 7.

Vol. 2, Figure 23--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 7, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Across the northern portion of the basin, the trend in forest structure has been an increase in mid-seral stages at the expense of both early- and late-seral stages (Hann and others 1997). Ecologically significant declines (Hann and others 1997) from historical to current times were projected for late-seral montane multi-story and single-story forests for the Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork ERUs. Late-seral subalpine multi-story forests also were projected to have declined significantly in two of these ERUs (Hann and others 1997). Specific habitat types for which there was greatest decline in areal extent within the three northern ERUs were western larch, interior Douglas-fir, and Engelmann spruce-subalpine fir old forests (vol. 3, appendix 1, table 4).

In the Southern Cascades, the source habitats that increased most strongly were single-storied old-forest Douglas-fir and multi-storied old-forest lodgepole pine (vol. 3, appendix 1, table 4). Increases in source habitats in portions of the Blue Mountains were associated largely with increases in multi-storied old-forests of Douglas-fir. In the Central Idaho Mountains ERU, the source habitats that decreased most in areal extent were old-forest single- and multi-storied Douglas-fir (vol. 3, appendix 1, table 4).

Condition of special habitat features--Densities of large-diameter snags and trees (>53 cm [21 in] d.b.h.) declined basin-wide from historical to current levels (Hann and others 1997; Hessburg and others 1999; Quigley and others 1996). Historical trends in smaller diameter snags were extremely variable (Hann and others 1997), so the overall basin-wide trend is unclear.

Other factors affecting species within the group--Cavity availability is dependent on the presence of primary excavators, most notably the pileated woodpecker and northern flicker (Hayward 1994c). Changes in population levels of these and other cavity excavators could affect boreal owl nesting opportunities.

Changes in forest structure could alter habitat suitability for voles and other important prey species and affect population levels of these species. In particular, changes in the abundance of coarse woody debris, snags, lichens, and fungi could significantly alter habitat suitability for many species found in older structural stages. This could affect the food resource for boreal owls and have a direct bearing on reproductive success.

Population status and trends--No reliable estimates of boreal owl population densities or trends in North America are available (Hayward 1994c).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 7 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The following issues have been identified as potentially influencing boreal owl conservation:

1. Declines in late-seral subalpine and montane forests, particularly in the Northern Cascades, Northern Glaciated Mountains, Lower Clark Fork, Upper Clark Fork, and Snake Headwaters ERUs.
2. Declines in large aspen trees and forests primarily because of fire suppression. Hayward and others (1993) found a relatively high use of aspen for nesting compared to available habitats.
3. Increasingly disjunct distribution of source habitats that may affect population structure (Hayward 1994a, 1997) and persistence of boreal owls.
4. Loss of large-diameter snags (>45 cm [18 in] d.b.h. recommended by Hayward [1994a]).
5. Loss of microenvironments for small-mammal prey. Changes in forest structure and composition (such as loss of snags and logs) could alter habitat for primary prey species (Hayward 1994a).

Potential strategies--The following strategies can be used to address the issues listed above:

1. (To address issue no. 1) Maintain existing habitats and accelerate development of subalpine and montane old-forest conditions within stands that are currently in mid-seral structural stages, particularly in the Northern Glaciated Mountains, Upper Clark Fork, and Lower Clark Fork ERUs.
2. (To address issue no. 2) Restore aspen forests throughout the basin where they have been reduced. This is particularly important in areas where aspen provides most of the nesting habitat for boreal owls (Hayward 1997).
3. (To address issue no. 3) Provide adequate links among subpopulations. Evaluate the links among subpopulations and use that information to identify areas that are highest priority for retention and restoration of habitat. This is of particular concern in the Northern Glaciated Mountains, Upper Clark Fork, and Lower Clark Fork ERUs, where reduction in the extent of source habitats has increased the isolation of remaining habitat patches.
4. (To address issue nos. 4 and 5) Retain large-diameter snags in all source habitats and provide for snag replacement over time.
5. (To address issue no. 5) Include boreal owl conservation within a larger, ecosystem context that addresses management of primary cavity nesters, small mammals, and forest structural components (Hayward 1994a).

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Adjust management activities to maintain and restore source habitats, particularly in the northern ERUs. Avoid extensive use of clearcuts, which may reduce habitat quality for 100 to 200 years (Hayward 1997). Small patch cuts implemented on long rotations may be compatible with maintenance of habitat quality for boreal owls (Hayward 1997). Thinning from below

and single tree selection may provide for development of nest structures.

2. (In support of strategy no. 2) Use clearcutting to regenerate aspen, focusing on the maintenance, at a landscape scale, of large aspen that provide nesting habitat for boreal owls (Hayward 1997). Where aspen regeneration is inhibited by domestic or wild ungulate browsing, use exclosures to protect regenerating stands and modify management to reduce browsing pressure.
3. (In support of strategy no. 4) Determine potential snag densities for each cover type used as source habitats by conducting surveys within remote areas, reserves, and natural areas. Use these baseline data to determine whether snags are below potential in other areas. Provide measures for snag protection and recruitment in all timber harvest plans.

GROUP 8--GREAT GRAY OWL

Results

Species ranges, source habitats, and special habitat features--Group 8 consists of breeding habitat for the great gray owl, a year-round resident of the basin. Great gray owls are distributed holarctically across the boreal forests of North America and Eurasia; they also inhabit other forests types at the southern extent of their range within the United States (Duncan and Hayward 1994). Within the basin, the great gray owl is widely distributed, although at low population levels, across most forested areas (fig. 24).

Vol. 2, Figure 24--Ranges of species in group 8 within the basin.

Within the basin, source habitats for great gray owls are old-forest, unmanaged young forest, and stand-initiation stages of montane forests, Engelmann spruce-subalpine fir, and riparian woodlands (vol. 3, appendix 1, table 1). Shrub or herb-tree regeneration also provide source habitats (vol. 3, appendix 1, table 1). Source habitats in the stand-initiation stage and herb-tree regeneration are used primarily for foraging. Old and unmanaged young forests are used for nesting and roosting, and more open stands (11 to 59 percent canopy cover [Bull and Henjum 1990]) are used for foraging. Great gray owls are a contrast species, requiring the juxtaposition of habitats used for foraging and for nesting-roosting (vol. 3, appendix 1, table 2).

Snags are a special habitat feature for great gray owls (vol. 3, appendix 1, table 2). They do not build their own nests but rely on existing platforms such as stick nests originally created by other birds or formed by dwarf mistletoe brooms, depressions in broken-topped dead trees, stumps, or artificial platforms (Bull and Henjum 1990, Duncan 1992, Mikkola 1983, Nero 1980). In one study in northeastern Oregon (Bull and Henjum 1990), 51 percent of the nests were stick platforms, 29 percent were on artificial platforms, and 20 percent were in natural depressions on broken-topped dead trees ($n = 49$). Of the stick nests, 68 percent were made by northern goshawks, 12 percent made by red-tailed hawks, and 20 percent were natural platforms formed by dwarf mistletoe brooms. Large branches are needed to support large stick-nests averaging 74 cm (29 in) long, 65 cm (26 in) wide, and 27 cm (11 in) high (Bull and Henjum 1990), and nests in broken-topped trees must be wide enough to accommodate a family of owls. Such trees range from 46 to 94 cm (18 to 37 in) in d.b.h. (Bull and Henjum 1990).

Broad-scale change in source habitats--Historically source habitats for the great gray owl presumably

were broadly distributed throughout forested portions of the basin (fig. 25A). The greatest concentrations of habitat were in the northern portion of the basin in the Northern Cascades, Northern Glaciated Mountains, Lower Clark Fork, Upper Clark Fork, and Snake Headwaters ERUs (vol. 3, appendix 1, table 3). Source habitat is projected to have declined moderately or strongly in 50 percent of watersheds basin-wide, and to have increased moderately or strongly in nearly 40 percent of watersheds (fig. 26). Although the overall change in source habitat has not been great, there has been a significant shift in its geographic distribution with habitat becoming more extensive in the western and central portions of the basin and less abundant in the northeastern part (fig. 25C). Of the ERUs that support substantial source habitat, moderate or strong increases in more than 50 percent of watersheds were projected for the Southern Cascades, Upper Klamath, Blue Mountains, and Central Idaho Mountains. Decreases in more than 50 percent of watersheds were projected for the Columbia Plateau, Northern Glaciated Mountains, Lower Clark Fork, Upper Clark Fork, and Snake Headwaters (fig. 26). Mixed trends were projected for the Northern Cascades ERU.

Vol. 2, Figure 25--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 8.

Vol. 2, Figure 26--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 8, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--The increase in habitat in the Southern Cascades, Upper Klamath, and Blue Mountains was primarily attributed to an increase in late seral montane forests (Hann and others 1997). In the Blue Mountains, an increase in the stand-initiation structural stage also contributed to the increase in source habitats. In the Northern Cascades, increases in source habitats primarily were due to an increase in early-seral montane forests. Habitat also has increased in the Central Idaho Mountains where the increasing trend is primarily the result of an increase in late-seral multi-layer and early seral montane forests.

In the ERUs where habitat for this species has declined (primarily the northern and eastern parts of the basin), habitat loss can be attributed primarily to the substantial reduction in late-seral montane and subalpine forests and early-seral montane forests (Hann and others 1997). The only exception is the Columbia Plateau, where source habitat declined primarily because of the reduction in abundance of shrub or herb-tree regeneration habitat (vol. 3, appendix 1, table 4). In all of the ERUs where source habitats are projected to have declined, there has been a significant increase in managed mid-seral montane forests since the historical period (Hann and others 1997).

Our evaluation at the broad-scale did not assess the distribution of foraging habitat in relation to that for nesting habitat. Further analysis of the juxtaposition of foraging with nesting habitats is needed at a finer scale of resolution. Average breeding home range size of individual adult great gray owls has been

calculated as 4.5 km² (1.7 mi²) (Bull and Henjum 1990) and 2.6 km² (1.0 mi²) (Craighead and Craighead 1956), and the ranges of adults are overlapping (Bull and Henjum 1990). Within each home range, a mixture of foraging and nesting habitat is needed. Analyses completed for the basin do not reveal landscape patterns at the scale of individual home ranges. Results for source habitats shown here for both the current and historical time periods are likely overestimates as they do not take into account the need for juxtaposition of habitats.

Condition of special habitat features--According to the landscape assessment (Hann and others 1997), the forests of the current period are more homogeneous than historical forests. Old-forest structures, remnant large trees, and the presence of medium to large trees in all forest structural classes have been reduced (Hann and others 1997). Densities of large-diameter snags (>53 cm [21 in] d.b.h.) likely declined basin-wide from historical to current levels (Quigley and others 1996, USDA Forest Service 1996). Presumably, the overall loss in large and medium trees and snag structures has reduced the availability of nest sites for great gray owls.

Other factors affecting the group--An additional factor may be the use of poisons to control pocket gopher populations. Such programs likely reduce the prey base for great gray owls (Hayward 1994b).

Population status and trends--No long-term, rigorous, or standardized surveys have been done of great gray owl populations within the basin (Duncan and Hayward 1994).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 8 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The following issues were derived from the analysis of source habitats and from published literature.

1. Decline of late-seral and early-seral stages of montane and subalpine forests, particularly in the northern and eastern parts of the basin.
2. Decline in availability of large trees and snags in all seral stages of montane and subalpine forests.
3. Encroachment of conifers into natural meadow systems, eliminating potential foraging habitat.
4. Reduced duration of early seral stages because of intensive planting and thinning.
5. Decline in prey resulting from use of poisons to control pocket gophers.

Potential strategies--Habitat for great gray owls would benefit from the following strategies that address the issues listed above:

1. (To address issue no. 1) Conserve existing older forest that is considered source habitat for this species, particularly in the Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork. The older forests that are source habitats for great gray owls have greater likelihood of being used for nesting if such stands are near open or early forests, which are used for foraging.
2. (To address issue no. 1) Accelerate the development of old-forest conditions in existing mid-seral stands.

3. (To address issue no. 2) Maintain and recruit large (>50 cm [20 in] d.b.h.) (Bull and Henjum 1990) live trees and snags for potential nesting strata.
4. (To address issue no. 3) Maintain and restore natural meadow systems that are adjacent to or near areas of old forest that have nesting platforms for great gray owls.
5. (To address issue nos. 1 and 4) Maintain a spatial and temporal mix of nesting (late seral) and foraging (early seral) habitats. Continuity of foraging habitat must be maintained through prudent long-term planning of timber harvest and other forest management activities.
6. (To address issues nos. 1 and 2) In evaluating and managing for long-term habitat quality, consider factors that influence populations of nest-building species (goshawk, red-tailed hawks, and ravens) and tree pathogen-insect interactions that can influence branch development (dwarf mistletoe brooms).
7. (To address issue no. 5) Avoid the use of poisons to control pocket-gopher populations near nesting habitat for great gray owls.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Focus retention efforts for late-seral montane and subalpine forests on sites where risks of catastrophic loss are relatively low.
2. (In support of strategy no. 2) Use prescribed burning and precommercial thinning to accelerate the development of old-forest conditions in mid-seral stands.
3. (In support of strategy no. 3) Maintain and restore natural meadow systems with the use of prescribed burning and removal of encroaching conifers.
4. (In support of strategy no. 3) Close roads to minimize removal of snags where such removals are reducing habitat quality for great gray owls. In addition or as an alternative to road management, actively enforce fuel wood regulations to minimize removal of snags.

GROUP 9--BLACK-BACKED WOODPECKER

Results

Species ranges, source habitats, and special habitat features--The black-backed woodpecker is a year-round resident that occurs in various forest types throughout the basin, except in southern Idaho ERUs (fig. 27). Source habitats of the black-backed woodpecker include old-forest stages of subalpine, montane, and lower montane forests and riparian woodlands (vol. 3, appendix 1, table 1). Both managed and unmanaged young forest stages of lodgepole pine also provide source habitat (vol. 3, appendix 1, table 1).

Vol. 2, Figure 27--Ranges of species in group 9 within the basin.

Burned conifer forests (Caton 1996, Hoffman 1997, Hutto 1995, Marshall 1992, Saab and Dudley 1998) and other insect-infested forests (Goggans and others 1988) provide key conditions necessary for both nesting and foraging. Habitat requirements for nesting include mature and old trees infested with disease, heart rot, or in early stages of decay (Goggans and others 1988). This species forages almost exclusively on the larvae of bark beetles (Scolytidae) and wood-boring beetles (Cerambycidae and Buprestidae) (Marshall 1992), which are obtained from tree trunks by scaling or flaking bark (Bull and others 1986a) and by excavating logs and the base of large-diameter tree trunks (Villard 1994). Thus, black-backed woodpeckers require conditions that produce bark and wood-boring beetle sources, including fire-, wind- or insect-killed mature or old pines, and other trees that have flaky bark (Dixon and Saab, in prep.; Marshall 1992). Both live and dead trees are used for foraging. Once trees have dried out 2 to 3 yr after mortality, bark beetles decline and use by this woodpecker also declines (Bull 1980). Populations are irruptive in response to bark beetle outbreaks in recently fire-killed forest stands or where trees become susceptible to bark beetle attacks through maturity (Baldwin 1968, Blackford 1955, Lester 1980). In the northern Rockies, early postfire conditions (1 to 5 yr after fire) are critical for supporting populations (Hutto 1995). Black-backed woodpecker abundance was not correlated to burn size but best correlated to the number of small snags remaining after fire in the northern Rockies (Hutto 1995). Summer home ranges for single birds differ in size from 72 to 328 ha (178 to 810 acres), depending on the quality of habitat (Goggans and others 1988). Goggans and others (1988) estimated that a single black-backed woodpecker requires an area of 193 ha (477 acres) of which 59 percent should be mature to old-forest conditions. They also suggested that a minimum management area for a nesting pair in lodgepole forests should be 387 ha (956 acres) of mature or old-forest conditions.

Snags are a special habitat feature for black-backed woodpeckers (vol. 3, appendix 1, table 2). Nest cavities are excavated in live trees with heart rot or recently killed trees (dead < 5 yr). This species nests in ponderosa pine, lodgepole pine, and western larch trees in the Blue Mountains (Bull and others 1986a). In central Oregon, they nested in mixed-coniferous and lodgepole forests that were undergoing a mountain pine beetle outbreak (Goggans and others 1988). Selection for mature and old stands was reported in central Oregon based on nest, foraging, and roost sites (Goggans and others 1988). Nesting birds favor unlogged compared to salvage logged stands of burned forests in western Idaho (Saab and Dudley 1998) and western Montana (Caton 1996). Black-backed woodpeckers generally select relatively small-diameter trees for nesting compared with other cavity nesters of similar size. In the Blue Mountains, mean d.b.h. of nest trees was 37 cm (14.6 in) ($n = 15$), and trees were generally tall (>15 m [49 ft]) and recently dead (<5 yr) (Bull and others 1986a). The mean d.b.h. of nest trees in central Oregon was 28 cm (11 in) ($n = 35$) (Goggans and others 1988). In burned ponderosa pine forests of western Idaho, nest tree d.b.h. averaged 32 cm (12.6 in) ($n = 17$), nest trees had relatively light decay, nest sites were located in tree clumps, and tree (>23 cm [9 in] d.b.h.) densities surrounding nests averaged 125 per ha (51 per acre) (104 per ha [42 per acre] in logged and 151 per ha [61 per acre] in unlogged units [Saab and Dudley 1998]).

In an Oregon forest with a bark beetle epidemic, overall nesting success averaged 68.5 percent ($n = 19$ nests) (Goggans and others 1988). In contrast, nest success was 100 percent for nests monitored in burned forests of western Idaho ($n = 27$) (Saab and Dudley 1998) and northwestern Wyoming ($n = 14$) (Hoffman 1997). Nest losses in Oregon were attributed to predation by flying squirrels and Douglas squirrels (Goggans and others 1988). Few mammalian nest predators were observed recolonizing the large-scale burns of western Idaho or the burns in northwestern Wyoming during the first 3 yrs after fire (Dixon and Saab, in prep.). This suggests that large burned forests during early postfire years are potentially important source habitats for black-backed woodpecker.

Broad-scale changes in source habitats--The following analysis does not account for recently burned habitats that are likely important as source habitats for black-backed woodpeckers. Such areas are generally at too fine a scale, and too ephemeral, to have been reliably estimated in the landscape analysis.

Historically, source habitats for black-backed woodpeckers were broadly distributed throughout the range of the species within the basin (fig. 28A). The most concentrated areas of habitat occurred in portions of the Blue Mountains, Columbia Plateau, Upper Klamath, Southern Cascades, Northern Cascades, and Central Idaho Mountains ERUs (fig. 28A).

The current distribution of source habitats is more concentrated in the southern half of the basin and diminished in the northern half. The Upper Klamath, Southern Cascades, Blue Mountains, southern watersheds of the Columbia Plateau, and the Central Idaho Mountains currently support the greatest concentrations of habitat (fig. 28B). In contrast, source habitats in the northern portion of the basin are scarcer and less well distributed than historically (fig. 28B).

Moderate or strong declines in source habitats were projected in nearly 70 percent of watersheds throughout the basin, with moderate or strong increases in 23 percent of watersheds (fig. 29). The most widespread declines were in the northern and far eastern parts of the basin (fig. 28). Moderate or strong declines were projected in over 90 percent of watersheds within the Northern Glaciated Mountains, Lower and Upper Clark Forks, and Snake Headwaters ERUs (fig. 29). Moderate or strongly declining trends also were projected for over 50 percent of watersheds in the Northern Cascades, Columbia Plateau, and Blue Mountains ERUs. Moderately or strongly increasing trends were projected for the Upper Klamath ERU. More mixed trends were projected for remaining ERUs.

Vol. 2, Figure 28--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 9.

Vol. 2, Figure 29--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 9, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Source habitat declined in more than 50 percent of watersheds in seven ERUs--the Northern Cascades, Columbia Plateau, Blue Mountains, Northern Glaciated Mountains, Lower Clark Fork, Upper Clark Fork, and Snake Headwaters. In all but one of these (Snake Headwaters), there were ecologically significant declines in late-seral lower montane forests (Hann and others 1997). In addition, there were also significant declines in late-seral montane forests in the three ERUs in the north end of the basin where source habitats declined most dramatically (Northern Glaciated Mountains, Upper Clark Fork, and Lower Clark Fork) (Hann and others 1997). The declines in the Snake Headwaters resulted from declines in both montane and subalpine late-seral forests (Hann and others 1997). Increases in the Upper Klamath ERU were due to increases in both lower montane and montane late-seral forest (Hann and others 1997).

Condition of special habitat features--Basin-wide declines from historical to current conditions were estimated for late-seral forest stands and for large snags (USDA Forest Service 1996) as well as for medium and large trees in all forest structural classes (Hann and others 1997). Based on these declines in late-seral forests and in residual large trees in other forests, a decline in medium to large snags (23 to

53 cm d.b.h.) is a reasonable assumption (see Quigley and others 1996 and USDA Forest Service 1996).

Other factors affecting the group--The natural pattern of beetle outbreaks has been altered through silvicultural practices and fire management policies. Silvicultural practices directed at maximizing wood production by harvesting trees before they are susceptible to bark beetle attacks, and salvage logging of beetle-infested, fire-killed, and wind-killed trees reduced the occurrence of beetles in some areas. Elsewhere, fire management policies have lengthened natural fire regimes and allowed more frequent occurrences of beetles.

Road densities have increased significantly throughout the basin (Hann and others 1997), thereby allowing greater human access into forested regions and subsequent increases in snag removal for firewood.

Usurpation of nest cavities by hairy woodpeckers (Goggans and others 1988) and by Lewis' woodpeckers (Saab and Dudley 1995) could have negative effects on black-backed woodpeckers. Stress and elevated energetic costs associated with territorial encounters with hairy and Lewis' woodpeckers and potentially reduce reproductive success of black-backed woodpeckers.

Population status and change--Breeding Bird Surveys indicate that population trends from 1966 to 1995 have been stable within western North America ($n = 16$ routes) (Sauer and others 1996). Trend data generated by the BBS, however, may be inadequate for monitoring populations of black-backed woodpeckers because of their relatively uncommon status and because the species is often difficult to detect (Goggans and others 1988, Marshall 1992).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 9 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The following issues were developed from our analysis of source habitat trends and findings from other studies:

1. Decline of old forests, particularly in the northern portion of the basin.
2. Decline in availability of medium to large (23 to 53 cm [9 to 21 in]) trees and snags infected with bark beetles, disease, heart rot, or in the early stages of decay.
3. Decline in availability of large (>387 ha [956 acre]) forest stands with bark beetle outbreaks because of salvage logging, particularly in the northern basin.
4. Altered frequency of stand-replacing fires.

Potential strategies--The issues identified above suggest the following broad-scale strategies would be effective in facilitating the long-term persistence of the black-backed woodpecker.

1. (To address issue no. 1) Maintain existing old forests that include interior ponderosa pine, interior Douglas-fir, western larch, lodgepole pine, grand fir-white fir, Engelmann spruce-subalpine fir, aspen, and red fir cover types over the short term. Accelerate development of old-forest conditions in stands that are currently in mid-seral or early-seral stages. Maintenance and restoration of old forests is especially important within the range of this species where declines in old forests have been most

pronounced. Areas of emphasis include Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork ERUs.

2. (To address issue no. 2) Where suitable nesting and foraging trees and snags are limited, retain mature and old trees and snags susceptible to bark beetle infestations, disease, and heart rot, or in the early stages of decay.
3. (To address issue no. 3) Throughout the ranges of the species, manage watersheds to maintain foraging and nesting habitat, with the location of that habitat shifting through time. Maintain stands that have experienced beetle outbreaks and stand-replacing burns.
4. (To address issue no. 4) Restore fire as an ecological process in montane and lower montane forests.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Use silvicultural treatments of prescribed underburning and thinning only of small-diameter trees (≤ 25 cm [10 in] d.b.h.) to accelerate development of mid-successional stages to old forests, particularly in cover types of ponderosa pine, Douglas-fir, and western larch.
2. (In support of strategy no. 2) Develop guidelines for retention of existing snags (≥ 25 cm [10 in] d.b.h.) in all forests, especially those with recent stand-replacement fire, insects, and disease to lengthen the time that those stands are suitable for nesting by black-backed woodpecker. Close roads, particularly after postfire salvage, to minimize removal of snags for firewood. In addition or as an alternative to road management, actively enforce fuel wood regulations to minimize removal of snags.
3. (In support of strategy no. 2) Develop measures for snag recruitment in unburned forests. Snag recruitment in unburned forests, with high risks of stand-replacing fires, will provide nest trees during the first few years after wildfire.
4. (In support of strategy no. 3) Maintain some large (>387 ha [956 acre]) forest stands with bark beetle outbreaks for 5 yrs, when beetle occupancy diminishes.
5. (In support of strategy no. 3) Avoid postfire salvage logging in portions of large burned forests to maintain contiguous burned stands of at least 387 ha [956 acres].
6. (In support of strategy no. 3) Where postfire salvage logging is planned in burned, lower montane forests, retain snags in clumps rather than evenly spaced distributions and retain at least 104 snags per ha (42 per acre), of d.b.h. size >23 cm (9 in).
7. (In support of strategy nos. 3 and 4) Allow wildfires to burn in some forests with high fire risk to produce stand-replacing conditions, and avoid postfire salvage logging in portions of large burned forests for about 5 yrs postfire.

GROUP 10--OLIVE-SIDED FLYCATCHER

Results

Species range, source habitats, and special habitat features--Group 10 consists of migratory breeding habitat for olive-sided flycatchers. Their range within the basin extends throughout forested areas (fig. 30). Winter range for olive-sided flycatchers includes the Central American highlands, the Andes, and the Amazon (Willis and others 1993a).

Vol. 2, Figure 30--Ranges of species in group 10 within the basin.

Olive-sided flycatchers are a contrast species using coniferous old forests for nesting and either openings or gaps in old forests for foraging (vol. 3, appendix 1, table 2; Sharp 1992). Their source habitats are old-forest single- and multi-storied and stand-initiation stages of subalpine, montane, and lower montane forests. Specific cover types that serve as source habitat are Engelmann spruce-subalpine fir, interior Douglas-fir, red fir, grand fir-white fir, Sierra Nevada mixed conifer, and Pacific ponderosa pine. Olive-sided flycatchers are positively associated with recent burns (Hejl 1994).

Broad-scale changes in source habitats--The extent of source habitat for olive-sided flycatchers is substantial in nine ERUs: the Northern Cascades, Southern Cascades, Upper Klamath, Blue Mountains, Northern Glaciated Mountains, Lower Clark Fork, Upper Clark Fork, Snake Headwaters, and Central Idaho Mountains (fig. 31B). Basin-wide, the trend in source habitat for olive-sided flycatchers is nearly neutral, with source habitats increasing and decreasing in almost equal numbers of watersheds (fig. 32). Trends differed geographically with habitat decreasing moderately or strongly in more than 50 percent of watersheds in three ERUs in the northern basin (Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork) and increasing moderately or strongly in more than 50 percent of watersheds in three ERUs in the southern basin (Southern Cascades, Upper Klamath, and Blue Mountains) (fig. 32). Trends were more mixed in the remaining three ERUs with significant source habitat (fig. 32).

Vol. 2, Figure 31--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 10.

Vol. 2, Figure 32--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 10, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure associated with changes in source habitats--Increases in late-seral montane

forests (Hann and others 1997) were consistent across the three ERUs (Southern Cascades, Upper Klamath, and Blue Mountains), with increasing trends in more than 50 percent of watersheds. The greatest contributors to the increases were old-forest single-storied interior Douglas-fir and grand fir-white fir in the Southern Cascades; old-forest single- and multi-storied interior Douglas-fir in the Upper Klamath; and old-forest multi-storied interior Douglas-fir and grand fir-white fir in the Blue Mountains (vol. 3, appendix 1, table 4). For the three ERUs with decreasing trends in more than 50 percent of watersheds (Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork), there were consistent decreases in early seral lower montane and montane forests; late-seral lower montane and montane multi-layered and single-layered forests; and late-seral subalpine multi-layered forests (Hann and others 1997).

Condition of special habitat features--Changes in fire regimes (Hann and others 1997) have likely resulted in poorer habitat conditions for olive-sided flycatchers, but the magnitude of the change is unknown. Where altered fire regimes result in fewer but larger fires, it seems likely that the juxtaposition of the early-seral and late-seral habitats used by olive-sided flycatchers becomes less favorable. Likewise, decreases in both early-seral and late-seral forests in the Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork likely have resulted in a strong decrease in areas of contrasting habitat condition used by olive-sided flycatchers. Our evaluation at the broad-scale, however, did not assess the distribution of foraging habitat in relation to that for nesting habitat. Further analysis of the juxtaposition of foraging with nesting habitats is needed at a finer scale of resolution.

Other factors affecting the group--It has been suggested that changes in winter habitat have had a negative effect on olive-sided flycatchers (Marshall 1988).

Population status and trends--BBS data indicate a significant decline from 1966 to 1994 for olive-sided flycatchers in eastern Oregon and Washington (-2.5 percent per yr, $n = 25$, $P < 0.01$) (Sauer and others 1996). Saab and Rich (1997) reported significant 10-yr and 26-yr declines (4.2 percent per year and 2.9 percent per year, respectively) for flycatchers on BBS routes within the basin. They included the olive-sided flycatcher as one of 15 Neotropical migrants in the basin that are of high concern under all future management themes.

Management Implications

The following issues, strategies and practices may be useful to managers as a starting point for integration of potential resource objectives for group 10 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The following issues were identified from our analysis of source habitat trends:

1. Reductions in early- and late-seral subalpine, montane, and lower montane forests, particularly in the Northern Glaciated Mountains and Upper and Lower Clark Forks.
2. Changes in fire regimes that result in fewer, larger, and more destructive fires, thereby reducing the areas of juxtaposed early- and late-seral forests.

Potential strategies--The following strategies would benefit species in group 10:

1. (To address issue no. 1) Accelerate development of late-seral conditions in lower montane, montane, and subalpine forests, particularly in the Northern Glaciated Mountains and the Upper and Lower Clark Fork.

2. (To address issues nos. 1 and 2) Increase the amounts of early-seral lower montane and montane forests, focusing on early-seral conditions that result from fire. Such restoration efforts would be most beneficial if concentrated in the northern portions of the basin.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy nos. 1 and 2) Various silvicultural practices including thinning from below, burning, and uneven-age management could be used to help accelerate the development of old-forest conditions and the juxtaposition of early-seral and late-seral habitats used by olive-sided flycatchers.

GROUP 11 - THREE-TOED WOODPECKER AND WHITE-WINGED CROSSBILL

Results

Species ranges, source habitats, and special habitat features--Group 11 consists of the three-toed woodpecker and white-winged crossbill, both of which occur at upper elevations throughout the basin. The range of the three-toed woodpecker is somewhat broader than that of the crossbill, occupying a greater portion of western Montana and central Oregon (fig. 33). The three-toed woodpecker is a year-round resident of the basin, whereas the white-winged crossbill is primarily a winter migrant, although occasional summer flocks have been observed (Harrington-Tweit and Mattocks 1985).

Vol. 2, Figure 33--Ranges of species in group 11 within the basin.

Source habitats for group 11 are late-seral subalpine and montane forests. Source habitats shared in common by the two species are old forests of lodgepole pine, grand fir-white fir, and Engelmann spruce-subalpine fir. The three-toed woodpecker also uses whitebark pine and mountain hemlock, and the white-winged crossbill occurs in western larch and Pacific silver fir-mountain hemlock (vol. 3, appendix 1, table 1).

Specific habitats used by the three-toed woodpecker are mature and overmature stands with bark beetles, disease, and heart rot (Goggans and others 1988) and recent stand-replacing burns with abundant wood-boring insects (Caton 1996, Hutto 1995). Three-toed woodpeckers forage predominantly on wood-boring beetle larvae (Stallcup 1962) and are attracted to areas with high concentrations of beetles, particularly in spruce and lodgepole pine (Bock and Bock 1974, Hogstad 1976, Villard 1994). Snags, a special habitat feature used for nesting (vol. 3, appendix 1, table 2), generally fall within the diameter range of 22 to 50 cm (9 to 20 in) (Bull 1980, Lester 1980). Because snags are used for foraging as well as nesting, large burns and beetle-infested stands are strongly favored for breeding over unburned or non-infested stands (Caton 1996, Goggans and others 1988). The period when burns and beetle-infested stands are useful for foraging is limited to about 5 yr, because beetles no longer use snags after they have dried out (Bull 1980). For nesting, however, the presence of heartrot may be required for cavity excavation (Goggans and others 1988), and fire-killed conifers generally do not develop this stage of decay until more than 5 yr postfire (Caton 1996). Older snags within burns or beetle outbreaks generally satisfy nesting requirements.

Crossbills are highly dependent on conifer cone crops and congregate where seed production is locally

abundant (Benkman 1992). The initiation of reproduction is triggered by abundance of conifer seeds. Nesting has been recorded every month of the year and occurs whenever the seed intake rate is sufficient for egg formation in females (Benkman 1990).

Broad-scale changes in source habitats--Trends in habitat availability for group 11 differ geographically. Historically, source habitats likely were distributed throughout most of the mountainous regions of the basin but generally occupied <25 percent of any given watershed (fig. 34A). The current extent of source habitats seems to have roughly the same geographic distribution, but the amount of habitat in the northern portion of the ranges of the species generally declined, whereas habitat in the south increased (fig. 34B). Basin-wide, source habitats increased moderately or strongly in 38 percent of the watersheds and decreased moderately or strongly in 54 percent (fig. 35). The ERUs that support significant amounts of habitat for the group and had moderately or strongly increasing trends in more than 50 percent of watersheds were the Southern Cascades, Upper Klamath, Blue Mountains, and Central Idaho Mountains (fig. 35). The ERUs for which moderate or strong declines were projected in more than 50 percent of watersheds were the Northern Cascades, the Northern Glaciated Mountains, Lower Clark Fork, Upper Clark Fork, and the Snake Headwaters (fig. 35).

Vol. 2, Figure 34--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 11.

Vol. 2, Figure 35--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 11, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Ecologically significant increases were projected by Hann and others (1997) for late-seral montane forests in all four ERUs in which source habitat increased in more than 50 percent of watersheds. For the five ERUs for which source habitats were projected to decline in more than 50 percent of watersheds, ecologically significant declines were projected in late-seral subalpine forests in the Northern Cascades; for late-seral montane forests in the Lower Clark Fork; and for both late-seral subalpine and late-seral montane forests in the Northern Glaciated Mountains, Upper Clark Fork, and Snake Headwaters.

Condition of special habitat features--Trends in snag availability within group 11 source habitats are unknown at the broad scale. Densities of large-diameter snags (>53 cm [21 in] d.b.h.) likely declined basin-wide from historical to current levels (Quigley and others 1996). The trend in smaller snags (22 to 50 cm [9 to 21 in]) used by three-toed woodpeckers is, however, unknown.

Other factors affecting the group--Three-toed woodpeckers are adapted to shifting their foraging areas to coincide with high concentrations of wood-boring beetles (Koplin 1969). Availability of this shifting food resource could be affected by salvage logging of large burns and beetle-infested stands, and maintenance of conifer stands in vigorous condition through silvicultural thinning.

Population status and trends--There are insufficient sightings in the BBS data records to determine population trends for either white-winged crossbills or three-toed woodpeckers within the basin. Summarized across the West, however, three-toed woodpecker occurrences on 14 BBS routes have declined an average of 0.7 percent annually between 1966 and 1995 ($\bar{n} = 14$, $\bar{P} < 0.05$; Sauer and others 1996).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 11 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The following issues were identified from our analysis of source habitat trends and from the findings of current research on group 11 species:

1. Decline in late-seral subalpine and montane forests. Cover types with basin-wide decline are western larch and whitebark pine. Declines of Engelmann spruce-subalpine fir are most notable in northern portions of the basin.
2. Potential decline in key components of the shifting food and nesting resource that is characterized by large areas of conifer trees infected with bark beetles, disease, heart rot, or in the early stages of decay.

Potential strategies--The following strategies could be used to maintain habitat in the southern and western portions of the basin and to reverse broad-scale declines in the northern and eastern regions:

1. (To address issue no. 1) Basin-wide, maintain remaining old forests of western larch and whitebark pine, and actively manage to promote their long-term sustainability.
2. (To address issue no. 1) In the Northern Glaciated Mountains, Upper Clark Fork, and Snake Headwaters ERUs, accelerate development of old-forest conditions in montane and subalpine forests within areas currently dominated by mid-seral stages.
3. (To address issue no. 2) Throughout the ranges of the species, manage watersheds to maintain foraging and nesting habitat, with the location of that habitat shifting through time. For three-toed woodpeckers, maintain stands that have experienced beetle outbreaks and stand-replacing burns.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy nos. 1 and 2) Use understory thinning and prescribed burns, or both, to enhance development and sustainability of western larch and whitebark pine old forests.
2. (In support of strategy no. 3) Maintain some large (>214 ha [528 acres]) (Goggans and others 1988) forest stands with bark beetle outbreaks for at least 5 yrs, until beetle occupancy diminishes.
3. (In support of strategy no. 3) Where suitable nesting and foraging trees are under represented, retain mature and old trees susceptible to bark beetle infestations, disease, and heart rot, or in the early stages of decay.
4. (In support of strategy no. 3) Allow wildfires to burn in some forests with high fire risk to produce

stand-replacing conditions, and avoid postfire salvage logging in portions of large burned forests to maintain contiguous burned stands of at least 214 ha (528 acres) (Goggans and others 1988) for about 5 yrs postfire.

GROUP 12--WOODLAND CARIBOU

Results

Species ranges and source habitats--Group 12 consists of the woodland caribou, a year-round resident of the basin. Woodland caribou have never been widely distributed in the basin (fig. 36). They are currently restricted to an area within the Northern Glaciated Mountains that includes parts of northeastern Washington, northern Idaho, and northwestern Montana. Evidence of their continued persistence in Montana is scant (USDI Fish and Wildlife Service 1994). The suspected historical range of the woodland caribou (ICBEMP 1996i) included parts of five ERUs: Northern Glaciated Mountains, Lower Clark Fork, Central Idaho Mountains, and small portions of the Columbia Plateau and Upper Clark Fork (fig. 36). Woodland caribou were federally listed as endangered in 1984.

Vol. 2, Figure 36--Ranges of species in group 12 within the basin.

Source habitats for woodland caribou are late-seral subalpine and montane forests (vol. 3, appendix 1, table 1). In total, five cover type-structural stage combinations provide source habitats for the woodland caribou. These are western redcedar/western hemlock old-forest single- and multi-storied stands; grand fir-white fir old-forest single- and multi-storied stands; and Engelmann spruce-subalpine fir old-forest multi-storied stands (vol. 3, appendix 1, table 1).

Broad-scale change in source habitats--This analysis of source habitats was based on the historical caribou range. Source habitats were projected to occur in five ERUs: the Columbia Plateau, the Northern Glaciated Mountains, the Lower Clark Fork, the Upper Clark Fork, and the Central Idaho Mountains (fig. 37). Source habitats in the Upper Clark Fork and Columbia Plateau were scarce (fig. 37).

Basin-wide, the trend in source habitats for caribou (historical to current periods) was mixed with 53 percent of watersheds projected with moderately or strongly negative trends and 41 percent with moderately or strongly positive trends (fig. 38). The three ERUs that supported significant caribou habitat each displayed a different trend. Trend in the Northern Glaciated Mountains was predominantly negative with a moderately or strongly negative trend projected for 65 percent of watersheds (fig. 38). For the Lower Clark Fork, a strongly positive trend was projected for 50 percent of watersheds and a strongly negative trend for 38 percent (fig. 38). Finally, a mixed trend was projected for the Central Idaho Mountains with watersheds split almost evenly among those showing a moderately or strongly negative trend (58 percent) and those showing a moderately or strongly positive trend (52 percent) (fig. 38).

Vol. 2, Figure 37--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 12.

Vol. 2, Figure 38--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 12, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--The predominantly negative trend for source habitat in the Northern Glaciated Mountains resulted largely from a strong decline in the old-forest multi-story stage of Engelmann spruce-subalpine fir (vol. 3, appendix 1, table 4). In the Lower Clark Fork ERU, the decrease in Engelmann spruce-subalpine fir old forest was offset by increases in western redcedar-western hemlock and grand fir-white fir old forests (vol. 3, appendix 1, table 4). In the Central Idaho Mountains, western redcedar-western hemlock, grand fir-white fir, and Engelmann spruce-subalpine fir old forests all increased (vol. 3, appendix 1, table 4), apparently masking geographic variation in trends that resulted in the mixed trend of watersheds increasing and decreasing (fig. 38).

Other factors affecting the group--Analyses by Zager and others (1995) indicated that adult mortality most limits population growth in the Selkirk population, and that at least 30 percent of this mortality is predator related. They suggested that high mortality rates may be associated with an increasing population of mountain lions, that in turn responded to expanding moose and white-tailed deer populations.

Woodland caribou populations are also subject to high rates of neonatal mortality, often approaching 50 percent. Calves typically make up 30 percent of the population at birth, but by recruitment age (1 yr) they typically make up <20 percent of the population (Scott and Servheen 1985).³

Both roads and human disturbance have been documented as causes of direct mortality for woodland caribou. Fatal collisions with automobiles occur on open roads in woodland caribou habitat (Scott and Servheen 1985). A high percentage of the annual mortality in the 1980s was attributed to illegal harvest by hunters and poachers (Scott and Servheen 1985). Caribou mortality due to illegal shootings has decreased since the species was federally listed as endangered in 1984, but illegal shooting has not been eliminated. Road densities and the potential for human disturbance have both increased from historical to current periods. In woodland caribou range, current average road densities are estimated to be moderate to high (Hann and others 1997).

High levels of disturbance by snowmobiles can cause caribou to abandon portions of their range, although low levels of snowmobile use are believed to be compatible with caribou occupancy of an area (Simpson 1987).

Population status and trends--Historically, caribou were distributed throughout the northeastern, north-central, and Northwestern United States. Their range within the basin included northwestern Montana and Idaho south to the Salmon River (USDI Fish and Wildlife Service 1994). By the 1960s, their range in the United States was restricted to the Selkirk Mountains of northeastern Washington and northern Idaho (USDI Fish and Wildlife Service 1994). The reduction in the range of the caribou was probably due to a combination of habitat fragmentation (resulting from both fires and timber harvest) and excessive mortality from overharvest and vehicle collisions.

In the 1950s, the Selkirk population of caribou in northeastern Washington, northern Idaho, and southeastern British Columbia was estimated at about 100 animals (Evans 1960, Flinn 1956). By the early 1980s, this population had declined to 25 to 30 animals whose distribution centered around Stagleap Provincial Park, British Columbia (Scott and Servheen 1985). The population in Idaho was augmented with animals from British Columbia three times between 1987 and 1990. The result was the establishment of a herd in the Idaho portion of the Selkirk Mountains. Populations are declining, however (see footnote 3; Zager and others 1995). Additional augmentation efforts occurred in the Washington portion of the Selkirks in 1996 and 1997.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 12 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The primary issues for woodland caribou are reported in the Selkirk Mountain Woodland Caribou Recovery Plan (USDI Fish and Wildlife Service 1994).

1. Reductions in source habitat in key portions of caribou range.
2. Illegal shooting, including accidental shooting by deer and elk hunters.
3. Predation by mountain lions, bears, wolves, and coyotes.
4. Mortality from vehicle collisions.
5. Displacement resulting from other human disturbance (for example, snowmobiles [Simpson 1987]).

Potential strategies--The U.S. Fish and Wildlife Service has established the following strategies that would provide recovery benefits for woodland caribou:

1. (To address all issues) Maintain the two existing caribou herds in the Selkirk ecosystem, and establish a third herd in the western portion of the Selkirk Mountains in eastern Washington.
2. (To address issue no. 1) Provide for at least 179 415 ha (443,000 acres) of suitable and potential caribou habitat in the Selkirk Mountains to support a self-sustaining population.

Practices that support strategies---The following practices would be effective in implementing the strategies listed above (taken from the Selkirk Mountain Woodland Caribou Recovery Plan [USDI Fish and Wildlife Service 1994]):

1. (In support of strategy no. 1) Reduce the impacts of poaching and hunting through outreach programs, restriction of access, and more effective law enforcement.
2. (In support of strategy no. 1) Reduce impacts of caribou-vehicle collisions by identifying areas where collisions are most likely and taking corrective actions (for example, reducing vehicle speeds, rerouting or closing roads, or increasing driver awareness.).
3. (In support of strategy no. 1) Identify most important additional sources of mortality by following radio-collared animals. Reduce other causes to the extent possible, recognizing that some mortality is unavoidable (for example, predation by other listed species).

4. (In support of strategy no. 1) Reduce impacts because of genetic and demographic influences by continuing augmentation and monitoring the success of augmentation efforts (but see Zager and others [1995] for cautions concerning the prognosis for augmentation efforts).
5. (In support of strategy no. 2) Maintain existing late-seral montane and subalpine forests within the areas designated to support caribou herds. Accelerate the development of old-forest conditions in currently mid-seral stands within these areas.
6. (In support of strategy no. 1) Evaluate the effects of roads, motorized vehicles, and recreational activities on caribou. Where such uses are not compatible with recovery (for example, where intensive snowmobile use is displacing caribou) implement standards (such as access timing or area closures) to address the issues.

GROUP 13 --NORTHERN FLYING SQUIRREL

Results

Species ranges, source habitats, and special habitat features--This group consists of the northern flying squirrel, which is a year-round resident of the basin. Flying squirrels occur throughout forested portions of the basin (fig. 39). Source habitats for this species include old-forest and unmanaged young-forest stages of subalpine, montane, lower montane, and riparian woodland cover types (vol. 3, appendix 1, table 1). The understory reinitiation stage of most of these types also is shown as source habitat (vol. 3, appendix 1, table 1; ICBEMPC). This stage is characterized by varying levels of canopy closure, and may contain large trees and other structures (vol. 1, table 4; Hann and others 1997) characteristic of northern flying squirrel habitat (Carey 1995). Because the understory reinitiation stage is highly variable (Hann and others 1997), however, its suitability as source habitat for flying squirrels is also variable.

Vol. 2, Figure 39--Ranges of species in group 13 within the basin.

Two special habitat features have been identified for northern flying squirrels (vol. 3, appendix 1, table 2). Flying squirrels nest in cavities that result from either damage to trees or excavation by woodpeckers (Carey 1995). Thus, snags are a special habitat feature, although squirrels also use cavities in live trees and external stick nests (Carey 1995, Waters and Zabel 1995). In a study in western Oregon, Carey (1991) found that snags containing nests average 89 cm (35 in) d.b.h. Down woody material is also an important feature of flying squirrel habitat (Carey 1991), presumably because of its role in supporting lichens and fungi that are the principle components of the diet of squirrels.

Broad-scale changes in source habitats--Historically, source habitats likely occurred throughout the forested portions of the basin (fig. 40A). Changes from historical have resulted in a reduction in the concentration of habitat across much of the range of the squirrel, with areas of increased habitat in the northeastern, central, and southwestern portions of the basin (figs. 40B, C). Overall, habitat has declined moderately or strongly in nearly 60 percent of watersheds in the basin and increased moderately or strongly in 27 percent of watersheds (fig. 41).

In eight ERUs, source habitat declined moderately or strongly in more than 50 percent of watersheds. These ERUs are the Northern Cascades, Southern Cascades, Columbia Plateau, Blue Mountains,

Northern Glaciated Mountains, Upper Clark Fork, Lower Clark Fork, and Snake Headwaters. Source habitat increased moderately or strongly in more than 50 percent of watersheds in the Upper Klamath and had mixed trends in the Central Idaho Mountains. Only relatively small amounts of habitat are present in the remaining three ERUs.

Vol. 2, Figure 40--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 13.

Vol. 2, Figure 41--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 13, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Ecologically significant decreases were projected (Hann and others 1997) for late-seral lower montane forests in seven of the eight ERUs for which source habitat declined in more than 50 percent of watersheds. The exception was the Snake Headwaters where significant declines were projected in late-seral montane and subalpine forests but not in late-seral lower montane forests. In addition to the declines in late-seral lower montane forests, there were declines in late-seral montane and late-seral subalpine forests in the Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork (Hann and others 1997). Declines in late-seral subalpine forests also contributed to the decreases in source habitat in the Northern Cascades and Blue Mountains.

Unmanaged young forest and understory reinitiation stages declined throughout the basin, including substantial losses in unmanaged young forest in the Northern Cascades and Upper Snake for cover types used as source habitat by northern flying squirrels (vol. 3, appendix 1, table 4). An exception to this general pattern of decreases were increases in understory reinitiation in the Northern Glaciated Mountains and Lower Clark Fork. These increases likely account for the areas of increasing source habitat concentration that were projected (fig. 40) within these ERUs for which there were otherwise general declines in source habitat. Because these mid-seral stages, and particularly the understory reinitiation stage, are quite variable, these projected increases merit further evaluation at a finer scale.

In the Upper Klamath, the only ERU for which an increase in source habitat was projected in more than 50 percent of watersheds, there were ecologically significant increases in late-seral lower montane, montane, and subalpine forests (Hann and others 1997).

Condition of special habitat features--Densities of large-diameter snags (>53 cm [21 in] d.b.h.) likely declined basin-wide from historical to current levels (Quigley and others 1996, USDA Forest Service 1996).

Other factors affecting the group--Forest management practices may have a significant effect on the hypogeous sporocarps of mycorrhizal fungi, a principal food source for flying squirrels. In a study in the Klamath Mountains, hypogeous sporocarps were nearly absent from clearcuts and were strongly

associated with coarse woody debris in late seral forests (Clarkson and Mills 1994). The negative association with clearcuts was thought to be due to microclimatic conditions and the effects of postharvest slash burns (Clarkson and Mills 1994). In a study in northeastern California, flying squirrel abundance was associated with the frequency of hypogeous sporocarps (Waters and Zabel 1995), but no correlation was found between sporocarp abundance and either thinning or broadcast burning (Waters and others 1994, cited in Waters and Zabel 1995). This study, however, did not examine sporocarp abundance in relation to clearcuts versus mature forests.

Population status and trends--No population trend information is available for northern flying squirrels within the basin.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 13 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The following issues were identified from the results of our analysis and other empirical research:

1. Widespread loss of old forests and associated structures (snags, logs, and cavities).
2. Reduced availability of remnant large trees and snags in all seral stages (Hann and others 1997).
3. Negative effect of forest management activities on fungus and lichen diversity and abundance (Carey 1991).

Potential strategies--The following strategies could be used to reverse broad-scale declines in source habitats and populations:

1. (To address issues nos. 1, 2, and 3) Maintain existing late-seral forests and encourage the development of appropriate habitat structures (snags, decayed down wood, and abundance of fungi and lichens) in mid-seral forests in all cover types used as source habitats, particularly in the northern half of the basin (Northern Cascades, Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork ERUs).

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) In the northern basin, give high priority to retention of old forests that have relatively low risk of loss through catastrophic fire. Priority should be given to large blocks having high interior-to-edge ratios and few large openings.
2. (In support of strategy no. 1) Actively recruit snags and logs from green trees to increase the representation of old-forest structures (snags and logs) in mid-seral stands and in old forests where snags and logs are in low density or absent.
3. (In support of strategy no. 1) Manage early- and mid-seral stands for increased vegetative diversity in order to encourage fungus and lichen diversity and abundance (Carey 1991).

GROUP 14--HERMIT WARBLER

Results

Species ranges and source habitats--Group 14 consists of the hermit warbler, a migrant that breeds in the basin and winters in high-elevation forests in Mexico and Central America. Most of the range of the hermit warbler occurs outside the basin along the west coast of British Columbia, Washington, Oregon, and California, overlapping the basin only along the crest of the Cascade Range (fig. 42) in three ERUs: the Northern Cascades, Southern Cascades, and Upper Klamath.

Vol. 2, Figure 42--Ranges of species in group 14 within the basin.

Habitat for hermit warblers is characterized by medium to large conifers (>31 cm [12.2 in] d.b.h.) (Morrison 1982). Source habitats within the basin include the old-forest and young-forest structural stages of interior Douglas-fir, red fir, grand fir-white fir, and Sierra Nevada mixed conifer (vol. 3, appendix 1, table 1). Both managed and unmanaged young forest support source habitat.

Broad-scale changes in source habitats--Source habitats for hermit warblers occur along the crest of the Cascade Range (fig. 43). Within this area, source habitat was projected to have increased moderately or strongly in over 75 percent of watersheds (fig. 44). Habitat decreased moderately or strongly in only 17 percent of watersheds. Source habitat increased moderately or strongly in 62 percent of watersheds in the Northern Cascades, in 90 percent of watersheds in the Southern Cascades, and in 100 percent of watersheds in the Upper Klamath (fig. 44).

Vol. 2, Figure 43--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 14.

Vol. 2, Figure 44--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 14, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--In the Northern Cascades, the increase in source habitat was due to increases in managed young-forest stages of interior Douglas-fir and grand fir-white fir (vol. 3, appendix 1, table 4). In the Southern Cascades, increasing source habitat was associated with increases in interior Douglas-fir and grand fir-white fir old forests and interior Douglas-fir managed young forest (vol. 3, appendix 1, table 4). In the Upper Klamath, increases were driven by increasing old-forest stages of interior Douglas-fir and grand fir-white fir (vol. 3,

appendix 1, table 4).

Other factors affecting the group--Hermit warblers forage along conifer branches, and sometimes deciduous trees and shrubs, for beetles, caterpillars, small flying insects, and spiders (Terres 1991). Thus, measures taken to control insects may impact hermit warblers.

The hermit warbler winters in high-elevation forests in Mexico and Nicaragua and sparingly into Costa Rica (Sharp 1992). Impacts to wintering habitats may negatively affect the species.

Population status and trends--There are insufficient data in the BBS information to be able to predict a population trend for the hermit warbler across the basin (Saab and Rich 1997). The BBS data analyzed within other geographic boundaries (Sauer and others 1996), however, showed an increasing trend in hermit warbler populations in eastern Oregon and Washington (7.6 percent per year, $n = 7$, $P < 0.01$, 1966 to 1979).

Management Implications

No significant issues were identified for hermit warblers or their habitat.

GROUP 15--PYGMY SHREW AND WOLVERINE

Results

Species ranges, source habitats, and special habitat features--This group consists of the pygmy shrew and wolverine, which are year-round residents of the basin. Wolverines occur in parts of all ERUs in the basin, although they are absent from the middle portion of the Columbia Plateau, and the south-central portion of the basin (fig. 45). The range of the pygmy shrew is restricted to the northeastern portion of the basin, primarily within the Northern Glaciated Mountains and Lower Clark Fork ERUs (fig. 45).

Vol. 2, Figure 45--Ranges of species in group 15 within the basin.

Both species should be considered generalists. Source habitats for pygmy shrews include virtually all structural stages of all subalpine and montane forests with the exception of Sierra Nevada mixed conifer (vol. 3, appendix 1, table 1). All stages of the shrub-herb-tree regeneration type also serve as source habitat for pygmy shrews. Source habitats for wolverines include alpine tundra and all subalpine and montane forests (vol. 3, appendix 1, table 1). Within the forest types, all structural stages except the closed stem exclusion stage provide source habitat.

Wolverines are predominantly scavengers, especially in winter when their diets consist primarily of ungulate carcasses (Banci 1994). In summer, they use a wider variety of foods including small mammals, birds, carrion, and berries (Weaver and others 1996). Copeland (1996) found that carrion-related food supplied 46 percent of wolverine diets in Idaho during both summer and winter. Banci (1994) suggested that diversity of habitats and foods is important to wolverines.

Several special habitat features have been identified for wolverines (vol. 3, appendix 1, table 2). Natal

dens in Idaho were primarily located in subalpine cirque basins in isolated talus surrounded by trees (Copeland 1996). There is also evidence that wolverines use down logs and hollow trees for denning (Copeland 1996; Pulliainen 1968, as cited in Banci 1994), and cavities in live trees also may be used (Ognev 1935, cited in Banci 1994; Pulliainen 1968). Both talus and areas associated with large, fallen trees were used as maternal den sites in Idaho (Copeland 1996).

No special habitat features were identified for the pygmy shrew.

Broad-scale changes in source habitats--Historically, source habitats likely occurred throughout the forested portions of the basin, with some of the greatest concentrations in the northeast (fig. 46A). From historical to current times, source habitat has increased in the central and western portions of the basin and undergone minor decreases in the north (fig. 46B).

Basin-wide, source habitat was projected to have increased moderately or strongly in 56 percent of watersheds and to have decreased moderately or strongly in 22 percent (fig. 47). Within the nine ERUs that support significant amounts of source habitat (fig. 47), five (Northern Cascades, Southern Cascades, Columbia Plateau, Blue Mountains, and Central Idaho Mountains) have undergone moderate or strong increases in more than 50 percent of watersheds, one (Upper Clark Fork) has undergone decreases in 50 percent or more of watersheds, and three (Northern Glaciated Mountains, Lower Clark Fork, and Snake Headwaters) have had mixed trends.

Vol. 2, Figure 46--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 15.

Vol. 2, Figure 47--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 15, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Causes for source habitat increases and decreases differed across ERUs (Hann and others 1997). Community types that most influenced habitat increases were early seral montane in the Northern Cascades, late-seral subalpine in the Southern Cascades, mid-seral montane in the Columbia Plateau, mid- and late-seral montane in the Blue Mountains, and early-seral subalpine and late-seral montane in the Central Idaho Mountains. In the Upper Clark Fork, community types that contributed most to the decline in habitat were early- and late-seral montane.

Condition of special habitat features--Densities of large-diameter snags (>53 cm [21 in] d.b.h.) and of large, remnant trees likely declined basin-wide from historical to current levels (Hann and others 1997; Hessburg and others 1999; Quigley and others 1996). Trends in snag abundance ultimately affect the availability of large down logs and cavities, whereas the decrease in large, remnant trees would likely translate to a decrease in large, hollow trees. Talus likely exists currently where it existed historically.

Other factors affecting the group--The clearcut method of timber harvest can negatively affect wolverines. Snow-tracking and radio telemetry in Montana indicated that wolverines avoided recent clearcuts and burns (Hornocker and Hash 1981). Copeland (1996), however, found that wolverines in Idaho commonly crossed natural openings, burned areas, meadows, or open mountain tops.

Populations of wolverines can be impacted by fur harvesting if trapping is not carefully regulated (Banci 1994). Within the basin, trapping is only allowed in Montana, and most of the harvest is believed to be incidental in traps set for other fur bearers (Banci 1994).

Copeland (1996) found that human disturbance near natal denning habitat resulted in immediate den abandonment but not kit abandonment. Disturbances that could affect wolverine are heli-skiing, snowmobiles, backcountry skiing, logging, hunting, and summer recreation (Copeland 1996, Hornocker and Hash 1981, ICBEMP 1996f). Wolverine densities in Montana, however, did not differ between the wilderness and nonwilderness portions of one study area, nor was their behavior or habitat use different, based on snow tracking and radio telemetry (Hornocker and Hash 1981). In addition, Hornocker and Hash (1981) concluded that movements of wolverines in Montana were not affected by highways.

Weaver and others (1996) argued that wolverines are less resilient than other large carnivores due to their low lifetime reproductive capability, susceptibility to natural fluctuations in scavenging opportunities, and vulnerability to trapping. They suggested that wolverines, along with grizzly bears, have a greater requirement for large, contiguous reserves than do other large carnivores such as gray wolves and mountain lions.

No information is available on other factors that might affect the pygmy shrew.

Population status and trends--Hash (1987) described a contraction in the North American range of the wolverine beginning around 1840 with the onset of extensive exploration, fur trade, and settlement. State records suggest very low wolverine numbers in Montana, Idaho, Oregon, and Washington from the 1920s through 1950s, with increases in wolverine sighting since the 1960s (Banci 1994). The increases in Montana (Newby and McDonald 1964, Newby and Wright 1955) and in Washington (Johnson 1977) may have resulted from dispersal from Canada.

Throughout its range, the pygmy shrew is considered rare (Feldhamer and others 1993), and basin-wide trends in pygmy shrew populations are unknown.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 15 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The following issues were identified from the results of our analysis and other empirical research:

1. Loss of montane and subalpine old-forests and associated structures (snags, logs, and cavities), particularly in the northern portion of the basin.
2. Low population numbers.
3. Increased negative effects from humans, resulting from higher road densities, increased technological advances in vehicular capabilities, and interest in winter recreation.

Potential strategies--The following strategies could be used to reverse broad-scale declines in source habitats and populations:

1. (To address issue no. 1) Increase the representation of late-seral stage forests in all cover types used as source habitats, particularly in the northern half of the basin (Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork ERUs).
2. (To address issue nos. 2 and 3) Identify refugia for long-term management of wolverine (Banci 1994).
3. (To address issue nos. 2 and 3) Provide adequate links among existing populations. These dispersal corridors likely do not require the same habitat attributes needed to support self-sustaining populations (Banci 1994).
4. (To address issue no. 3) Reduce human disturbances, particularly in areas with known or high potential for wolverine natal den sites (subalpine talus cirques).

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) In the northern basin, retain existing old forests and identify mid successional forests where attainment of old-forest conditions can be accelerated.
2. (In support of strategy no. 1) Actively recruit snags and logs from green trees to increase the representation of old-forest structures (snags and logs) in mid-seral stands and in old forests where snags and logs are uncommon or absent.
3. (In support of strategy no. 1) Retain slash piles and decks of cull logs to substitute for down logs over the short term.
4. (In support of strategy no. 2) Maintain current wilderness areas and other congressionally designated reserves as refugia for wolverine, and reduce human disturbances near den sites in these areas.
5. (In support of strategy no. 2) Identify existing areas with the following desired conditions, or manage selected areas to create the following desired conditions for species strongholds: large, contiguous blocks of forest cover with abundant snags and large logs and low road densities with connectivity to subalpine cirque habitats required for denning, security, and summer foraging habitat.
6. (In support of strategy no. 3) Identify isolated populations and unoccupied habitats and use interagency planning to develop broad-scale links over the long term.
7. (In support of strategy no. 4) Minimize new construction of secondary roads and close unneeded roads after timber harvests.

No explicit recommendations are available in the literature nor are any available from our results for the pygmy shrew.

GROUP 16--LYNX

Results

Species ranges, source habitats, and special habitat features--The lynx, a year-round resident of the basin, is the only member of group 16. The range of the lynx includes the northern, eastern, and central portions of the basin (fig. 48). There are limited records of lynx occurring in the Southern Cascades ERU (McKelvey and others 1999), but these records were not included in the range map delineated by Marcot and others (in prep.). In 1997, the U.S. Fish and Wildlife Service (USFWS) determined that the lynx warranted protection under the Endangered Species Act (ESA), but postponed listing because other species were in greater need of protection. In 1998, the USFWS settled litigation surrounding the lynx by proposing that lynx be listed as endangered in the lower 48 States. A final decision is expected in 2000.

Vol. 2, Figure 48--Ranges of species in group 16 within the basin.

Primary habitat for lynx is found in subalpine and montane forests that are cold or moist forest types (vol. 3, appendix 1, table 1; McKelvey and others 1999). Within the montane forest community, source habitats are provided by all vegetation types except Pacific silver fir-mountain hemlock, red fir, and Sierra Nevada mixed conifer. Within the subalpine forest community, only Engelmann spruce-subalpine fir provides source habitat. Lynx habitat includes various structural stages (Koehler and Aubry 1994, Ruggiero and others 1999).

Lynx forage primarily in early-seral forests and in some mid-seral forests that support high numbers of prey; lynx also use late-seral forests for denning and rearing young as well as for hunting alternative sources of prey (Ruggiero and others 1999). Consequently, source habitats for lynx are provided by most of the coniferous forest structural stages with the exception of old-forest single-storied stands (vol. 3, appendix 1, table 1). Riparian woodlands and shrublands are also source habitats.

Hollow down logs are a special habitat feature for lynx (vol. 3, appendix 1, table 2); logs are used both as den sites and resting places (ICBEMP 1996e, Koehler 1990).

Broad-scale changes in source habitats--Basin-wide, amounts of source habitats for lynx increased moderately or strongly in 47 percent of watersheds and decreased in 23 percent from historical to current periods (figs. 49 and 50). Habitat increased in more than 50 percent of the watersheds in two ERUs, the Blue Mountains and the Northern Glaciated Mountains (fig. 50). Trends were mixed in the remaining ERUs that contain significant habitat: Northern Cascades, Upper Clark Fork, Lower Clark Fork, Snake Headwaters, and Central Idaho Mountains (fig. 50).

Vol. 2, Figure 49--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 16.

Vol. 2, Figure 50--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 16, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure associated with changes in source habitats--A strong increase in mid-seral montane forests, along with increases in early- and mid-seral subalpine forests (Hann and others 1997), accounted for the increasing source habitat trend in the Northern Glaciated Mountains. Increases in mid- and late-seral montane forests and early- and mid-seral subalpine forests (Hann and others 1997) contributed to the overall increase in source habitats in the Blue Mountains. Mid-seral montane and subalpine forests also increased in the Lower Clark Fork, Upper Clark Fork, and Snake Headwaters ERUs; however, these increases were offset by decreases in early-seral montane forests and late-seral montane and subalpine forests (Hann and others 1997). In the Northern Cascades, increases in early-seral montane and subalpine forests were offset by decreases in mid- and late-seral subalpine forests (Hann and others 1997). There were increases in early- and late-seral montane and subalpine forests in the Central Idaho Mountains (Hann and others 1997), but these increases were not widespread enough to result in an overall moderate or strong ERU trend.

Condition of special habitat features--Hann and others (1997) reported a decrease in abundance and occurrence of large down logs in areas of traditional forest management. Large down logs are used by lynx for denning and rearing young (Ruggiero and others 1999).

Other factors affecting the group--Trapping can be a significant source of mortality for lynx (Bailey and others 1986, Carbyn and Patriquin 1983, Mech 1980, Nellis and others 1972, Parker and others 1983, Ward and Krebs 1985). Trappers are capable of removing from 60 to 80 percent of the individuals in a given lynx population (Bailey and others 1986, Parker and others 1983). In the basin, lynx can be legally trapped in Montana. Increased roading of Federal land has provided trappers greater access to lynx populations. Also, incidental takes of lynx during bobcat and coyote trapping seasons may be cause for concern, especially with low-density lynx populations.

Other forms of human disturbance also affect lynx. According to Koehler and Brittell (1990), minimal human disturbance is important to denning site selection. Winter recreation may have a significant effect on lynx populations. The packing effect of snowmobile trails may open areas of deep snow to foraging competition from other predators such as bobcats and lynx (Kohler and Aubry 1994, Ruggiero and others 1999). In the north Cascades, snowmobiling and other winter recreation have increased in the past decade, with suspected negative effects on lynx.⁴ The increase in interactions between human and lynx, primarily because of increased use of off-highway vehicles (including snowmobiles), may result in increased lynx mortality from intentional and unintentional shooting and collisions with vehicles (Koehler and Brittell 1990). Highways could also pose barriers to lynx movement or increase mortality from vehicle collisions (Ruediger 1996, Terra-Berns and others 1997).

Lynx populations are closely tied to snowshoe hare population trends, especially north of the basin (Butts 1992, Murray and Boutin 1991, Parker and others 1983, Weaver 1993). Lynx populations in the basin, however, may not be as cyclic as those at more northern latitudes (Brittell and others 1989, Koehler 1990). Within the basin, several other predators (bobcat, red fox, and some hawks and owl species) compete with lynx for snowshoe hare as prey, unlike areas to the north; many of these competing predators possibly respond more positively to human-induced habitat alterations (Roloff 1995). This

increased competition for prey may increase the vulnerability of lynx (Witmer and others 1998) as well as limit the size of lynx populations (Boutin and others 1986, Keith and others 1984).

Forest management practices have varying effects on both lynx and lynx prey habitat (Ruggiero and others 1999). Lynx do not hunt in large, open areas with little or no cover (Koehler 1990, Koehler and Brittell 1990), making large clearcut blocks potential barriers to movement (Koehler and Aubry 1994). Early-seral habitats created by fire or logging, however, are essential to maintain foraging areas for lynx prey, principally snowshoe hare (Koehler and Aubry 1994, Koehler and Brittell 1990). Koehler and Aubrey (1994) proposed that frequent, small patches of habitat alteration that mimic natural disturbance patterns would be beneficial. Postclearcut areas may not become suitable for snowshoe hare habitat for more than 10 years, and may not become optimal hare habitat for another 20 years (Koehler and Aubrey 1994). Relatively small patches of old forest (1 ha [2.5 acres]) are needed for denning, though these areas must be near and connected to good foraging habitat (Koehler and Brittell 1990). Travel corridors generally have a closed-canopy cover >2 m high (Brittell and others 1989).

Population status and trends--Empirical data for distribution of lynx within the basin are scarce, and data on abundance of lynx populations are not available. McKelvey and others (1999) recently summarized all known lynx locations in the United States, which provides a framework for designing and conducting future surveys and demographic studies of lynx populations.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 16 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The following issues for lynx were taken from the literature.

1. The lack of empirical information on population ecology, foraging ecology, den site characteristics, habitat relations at the landscape scale, and distribution and status in the basin (Ruggiero and others 1999).
2. Altered mosaic of source habitats because of fire suppression and logging (Hann et al. 1997).
3. Negative effects of human activities on lynx (Koehler and Aubry 1994).
4. The peninsular and disjunct distribution of suitable lynx habitat in the western mountains (Koehler and Aubry 1994), and the associated potential for population isolation or limited metapopulation structure to cause local or regional extirpations (Ruggiero and others 1999).

Potential strategies--

1. (To address issue no. 1) Develop an interagency research, inventory, and monitoring effort aimed at gathering information on population ecology, foraging ecology, den site characteristics, habitat relations at the landscape scale, and distribution and status in the basin.
2. (To address issue no. 2) Restore fire as an ecological process or use other forest management practices in montane and upper montane community types to provide for a suitable mosaic of early seral habitat rich in shrubs and well connected to late-seral habitat with abundant large down logs.

3. (To address issue no. 3) Design silvicultural treatments at a landscape scale with the needs of snowshoe hare and other lynx prey as one consideration.
4. (To address issue no. 3) Provide areas of high-quality lynx habitat that are protected from human disturbance (Koehler and Aubry 1994).
5. (To address issue no. 4) Develop a strategy to allow for interactions among lynx populations, including the provision of travel corridors (Koehler 1990) and broader landscape connectivity.
6. (To address issue no. 4) Develop a strategy to allow for population reintroductions as appropriate.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy nos. 2 and 3) Management of stand dynamics for lynx and snowshoe hares focuses on the creation of early and late old-forest structural stages consistent with historical variability. In designing forest landscapes, give management consideration to habitats for alternate prey species such as red squirrel, voles, and mice in addition to denning habitat for lynx. Down wood is an important denning habitat component. When thinning stands to meet timber management objectives, stands should either be thinned early before they are recolonized by snowshoe hares, or thinned when they are older (for example, 30 to 40 yr) and are little used by hares.
2. (In support of strategy no. 4) In areas of known or suspected lynx populations, close roads and areas to all vehicles as needed to minimize human disturbance, limit potential increase in competing predators, and provide for landscape connectivity among and within populations. Improve highway passage by using fencing and overpasses and underpasses.
3. (In support of strategy nos. 5 and 6) Identify areas that currently support high-quality lynx habitat, have low road densities, and are sites of recent lynx observation. Identify such sites as species strongholds, and use them as the backbone of a metapopulation strategy.

GROUP 17--BLUE GROUSE (SUMMER) AND MOUNTAIN QUAIL (SUMMER)

Results

Species ranges, source habitats, and special habitat features--Group 17 consists of summer habitats for both blue grouse and mountain quail. The range of the blue grouse includes the western, northern, central, and eastern portions of the basin (fig. 51). The range of the mountain quail includes southern Washington, Oregon, and western Idaho (fig. 51; Ehrlich and others 1988). Blue grouse are ground nesters that forage primarily on seeds, berries, and insects; the young feed heavily on insects (Ehrlich and others 1988). Mountain quail are also ground nesters and feed primarily on bulbs, greens, and insects (Ehrlich and others 1988).

Vol. 2, Figure 51--Ranges of species in group 17 within the basin.

Source habitats for group 17 include all structural stages except stem exclusion of interior Douglas-fir,

Sierra Nevada mixed conifer, and Pacific and interior ponderosa pine (vol. 3, appendix 1, table 1). In addition, blue grouse source habitats also include western larch, aspen, mixed-conifer woodlands, antelope bitterbrush-bluebunch wheatgrass, and wheatgrass bunchgrass. Chokecherry-serviceberry-rose is also source habitat for both species.

A special habitat feature for the mountain quail is riparian shrub (vol. 3, appendix 1, table 2). Mountain quail within the basin primarily are found within 100 to 200 m (328 to 656 ft) of a water source (Brennan 1989). The blue grouse (summer) is considered a contrast species as it is typically found at the interface of forest and open areas (Zwickel 1992; vol. 3, appendix 1, table 2).

Broad-scale changes in source habitats--Source habitats for blue grouse (summer) and mountain quail (summer) occur primarily in the forested ERUs across the basin (fig. 52A, B). The overall trend in source habitats since historical times has been neutral (fig. 53), with increasing trends occurring primarily in the western and southeastern part of the basin, and more decreasing trends occurring in the northeast part of the basin. The ERUs with increasing trends are the Southern Cascades, Upper Klamath, Northern Great Basin, Upper Snake, and Snake Headwaters. The ERUs with decreasing trends are the Lower Clark Fork, Upper Clark Fork, and Central Idaho Mountains. The remaining ERUs are overall neutral (Northern Cascades, Columbia Plateau, Blue Mountains, Northern Glaciated Mountains, and Owyhee Uplands).

Vol. 2, Figure 52--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 17.

Vol. 2, Figure 53--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 17, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Increases in source habitats in the Northern Cascades are primarily because of increases in managed young forests of interior Douglas-fir and interior ponderosa pine, whereas a similar decline occurred in old-forest ponderosa pine (vol. 3, appendix 1, table 4). Increases in source habitats in the Southern Cascades, Upper Klamath, and Blue Mountains, and southern portions of the Columbia Plateau are due primarily to increases in old forest. Decreases in source habitats in much of the northeastern part of the basin are due to declines in both late-seral and early-seral community types.

The primary changes in source habitats in the Upper Snake were an increase in wheatgrass bunchgrass (vol. 3, appendix 1, table 4). Hann and others (1997), however, suspect that in some areas that show increases in upland herblands (including wheatgrass bunchgrass), these areas may in fact be areas of early seral forests attributable to relatively recent timber harvest or large-scale wildfires, and were misclassified as upland herbland. In such a case, recent timber harvest or wildfire may have increased the quantity and quality of source habitat because of potential increases in shrubs. Increases in wheatgrass bunchgrass, however, also may be attributable to increases in exotic wheatgrasses such as crested wheatgrass, which does not provide source habitat for blue grouse. The increase in source

habitat in the Snake Headwaters is primarily due to an increase in both early- and mid-seral interior Douglas-fir (vol. 3, appendix 1, table 4).

Condition of special habitat features--Basin-wide analysis of riparian vegetation found significant changes, including widespread declines in riparian shrublands (Quigley and others 1996). Because of the scale of our analysis and the fine-scale nature of riparian shrubland habitats, likely the results of our analysis do not reveal the true loss in this important habitat component for mountain quail. Water impoundments, grazing, residential developments, and agricultural activities can alter the extent, composition, and structure of mountain quail habitat (Brennan 1990, Murray 1938, Vogel and Reese 1995). Remaining habitat in the basin is fragmented, and populations exist often in islands of habitat connected by narrow corridors of vegetation (Vogel and Reese 1995).

Because the blue grouse (summer) is a forest-open areas contrast species, the scale of this analysis does not allow determination of change in the juxtaposition of these contrasting habitats. Thus, this special habitat feature is not accounted for in the results presented above, and a finer scale analysis is needed to fully evaluate the status of their source habitats. A loss of interspersed early- and late-seral stages of forest partly because of altered fire regimes was identified by Lehmkuhl and others (1997) as a reason for a declining trend since the historical period of both habitat and populations of the blue grouse.

Other factors affecting the species--Some mountain quail populations migrate to lower elevations to winter (Brennan 1990, Ehrlich and others 1988, Leopold and others 1981). Winter habitat availability may be more limited than summer habitat because of severe winter weather in some mountainous areas (Edminster 1954). Low-elevation riparian shrub habitat is especially important during severe winters. Hydroelectric impoundments along the Columbia River and its tributaries have flooded thousands of acres of low-elevation winter habitat for mountain quail (Brennan 1990). One of the last remaining Idaho populations can be found along the Salmon River drainage in an area that experiences mild winters, thought to be one of the important variables for the continued presence of quail in this area (Brennan 1989).

Both blue grouse and mountain quail most often are found in areas with a high abundance of shrubs, which most likely are used for cover as well as forage (Brennan and others 1987, Zwickel 1992). Traditional forest managers commonly replanted harvested areas, thus hastening the rate of succession and shortening the time that a stand remains in the early seral stage (Hann and others 1997). This practice coupled with ground-disturbing site preparation before planting, often eliminates the herb, forb, and shrub structures from stands. Management activities such as salvage logging and planting in postfire habitats also may shorten the duration of these early-seral, shrub-dominated sites.

Grazing of domestic livestock may negatively impact blue grouse (Mussehl 1963, Zwickel 1972), as well as mountain quail (Brennan 1990).

The frequency and areal extent of wildfires declined since the early to mid 1900s because of suppression activities (Hann and others 1997). With the increased fuel loads in fire-suppressed areas, however, the trend since 1960 has changed, and the current extent of wildfires is approaching the early 1900s. This increase in postfire areas should benefit both blue grouse (summer) and mountain quail if these fires result in an increase in shrub vegetation.

Both species are negatively affected by human disturbance, primarily during the nesting/brood-rearing season (ICBEMP 1996h). The human population in the basin is estimated at 3 million, which is a substantial increase from the pre-European settlement period (McCool and others 1997). This change in population increases human encounters, thus having a potentially negative effect on both blue grouse and mountain quail. In particular, the introduction of human residents to an area also introduces domestic cats, which can be an effective predator of mountain quail (Edminster 1954, Jewett and others

1953, McLean 1930.)

There are open hunting seasons for blue grouse throughout the basin, whereas hunting for mountain quail is only allowed in some parts of Oregon.

Population status and trends--Blue grouse still occupy most of their original range, although historical populations may have been stronger in some areas (Zwickel 1992). Although mountain quail populations to the west of the basin seem to be stable, populations in the basin have experienced dramatic declines (Brennan 1990, Robertson 1989, Washington Department of Wildlife 1993a).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 17 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--Issues identified for group 17 were based on our analysis of source habitats as well as knowledge of finer scale habitat features for these species:

1. Decline in late- and early-seral source habitats, particularly in the northeastern part of the basin.
2. Changes in vegetation composition and structure of understory shrub habitat.
3. Loss of riparian shrubs.
4. Increased interaction with humans.
5. Isolated and disjunct populations of mountain quail vulnerable to extinction by stochastic events (that is demographic, environmental, or genetic stochasticity).

Proposed strategies--

1. (To address issue no. 1) Maintain and restore late-seral montane and lower montane forests.
2. (To address issue nos. 1 and 2) Increase the representation of shrub-dominated early seral forests.
3. (To address issue nos. 1 and 2) Restore fire as an ecological process in the montane and lower montane community groups.
4. (To address issue no. 3) Maintain and restore riparian shrubland habitats, including protecting existing areas from the encroachment of exotics.
5. (To address issue no. 3) Reduce habitat degradation by livestock grazing in areas currently occupied by mountain quail.
6. (To address issue no. 4) Restrict human access in areas of known nesting use by blue grouse and mountain quail.
7. (To address issue no. 5) Expand the current range of mountain quail within their historical habitats.

Practices that support strategies--The following practices would be effective in implementing the

strategies listed above:

1. (In support of strategy no. 1) Maintain existing old forest until mid-seral forests have developed into old forest at a level that is within the range of historical variability.
2. (In support of strategy no. 2) Leave some postfire areas unaltered to regenerate naturally.
3. (In support of strategy no. 3) Use prescribed fire to enhance growth and regeneration of understory or mountain shrub development. Use caution especially during spring breeding season, as fires can cause direct mortality to mountain quail (Clark 1935, McLean 1930, Spaulding 1949).
4. (In support of strategy no. 4) Reduce exotic weed invasions by plantings of native shrub and herbaceous vegetation in riparian shrubland habitats.
5. (In support of strategy no. 5) Remove or explicitly control the timing and intensity of grazing to discourage weed invasions and to minimize losses and allow for restoration of native riparian and mountain shrubs.
6. (In support of strategy no. 6) Reduce road densities and timing of management activities to reduce human interactions with these species, especially during the nesting and brooding season. In addition or as an alternative to reductions in road density, implement seasonal road closures during nesting and brooding periods.
7. (In support of strategy no. 7) Reintroduce and augment mountain quail after habitat enhancement.

GROUP 18--LAZULI BUNTING

Results

Species ranges and source habitats--Group 18 consists of the Lazuli bunting, a migratory breeder that occurs throughout the basin (fig. 54). Source habitats for Lazuli buntings are grass-forb-shrub edges, burns, early seral stages of conifer forest, and dense low vegetation along streams (Sharp 1992). Hutto (1995) found that Lazuli buntings demonstrated a strong positive response to early successional burned forests, resulting from stand-replacing fires that occurred in a broad spectrum of coniferous forest types across western Montana and northern Wyoming. This bunting was also a common nesting species in recently burned ponderosa pine/Douglas-fir forests of western Idaho (Saab and Dudley 1998). The Lazuli bunting is a shrub-nesting insectivore, foraging primarily by gleaning off foliage (Ehrlich and others 1988).

Vol. 2, Figure 54--Ranges of species in group 18 within the basin.

Source habitats analyzed in this report are the stand-initiation stage of the montane, lower montane, riparian woodland terrestrial communities and also chokecherry-serviceberry-rose (vol. 3, appendix 1, table 1). Among landscape and microhabitat features of cottonwood forests in eastern Idaho, the most important predictor of Lazuli bunting occurrence was shrub density and cover (Saab 1999). Other significant predictors of their occurrence included herbaceous ground cover and willow subcanopies, providing foraging and nesting habitat, respectively. Additionally, their relative abundance was

significantly reduced in forest patches managed for grazing compared with unmanaged patches (Saab 1996, 1998). In cottonwood forests of western Montana, the abundance of Lazuli buntings also was reduced in heavily grazed areas, as compared to lightly grazed areas (Mosconi and Hutto 1981).

Broad-scale changes in source habitats--Historically source habitats for group 18 were broadly distributed throughout the mountainous regions of the basin, though most watersheds with source habitats contained less than 25 percent area in source habitats (fig. 55A). Currently source habitats are more patchily distributed and absent from many watersheds that historically contained these habitats (fig. 55B).

The trend in source habitats from historical to current periods was negative to strongly negative for nearly 60 percent of the watersheds in the basin (figs. 55C and 56). About 33 percent of the watersheds basin-wide had positive trends in source habitats (fig. 56). Eight ERUs had negative to strongly negative trends, including the Upper Klamath, Northern Great Basin, Columbia Plateau, Blue Mountains, Northern Glaciated Mountains, Lower Clark Fork, Upper Clark Fork, and Upper Snake. Trends were neutral in the Southern Cascades and Owyhee Uplands. Three ERUs, the Northern Cascades, Snake Headwaters, and Central Idaho Mountains, had positive trends.

Vol. 2, Figure 55--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 18.

Vol. 2, Figure 56--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 18, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--For the ERUs with positive trends, increased area of various cover types, especially Douglas-fir, Englemann spruce, lodgepole pine, and aspen, were responsible for the trend (vol. 3, appendix 1, table 4). For the eight ERUs with negative trends, the loss of early seral Douglas-fir, lodgepole pine, interior ponderosa pine, and western larch contributed most to the trend. Nearly 100 percent of the western larch stand-initiation stage was eliminated in these ERUs.

In addition, there have been basin-wide declines in riparian woodlands at the broad scale (Hann and others 1997). Smaller patches of riparian vegetation, especially riparian shrublands have declined in extent basin-wide, because of disruption of hydrologic regimes from dams, water diversions, and road construction. Additionally, grazing and trampling of riparian vegetation by livestock, and increased recreational use along stream courses have reduced riparian habitats (USDA Forest Service 1996). Low-elevation wetlands in Idaho are considered "endangered" based on 85- to 98-percent decline since European settlement (Noss and others 1995)

Other factors affecting the group--Traditional forestry practices commonly tried to accelerate the regeneration process in harvested areas by planting, thus hastening the rate of succession and shortening the time that a stand remained in the early seral stage (Hann and others 1997). This practice

coupled with ground-disturbing site preparation activities before planting often eliminated the herb, forb, and shrub structure from stands. Planting in postfire habitats also shortens the duration of the stand-initiation stage. Salvage logging in postfire habitats may reduce the availability of tall structures used for singing perches. Hutto (1995) found that the relative abundance of many bird species including the Lazuli bunting differed between recently burned and recently harvested forests. Composition of trees, snags, and shrubs subsequent to a burn can differ depending on fire intensity and postfire timber harvest.

According to Hann and others (1997), the frequency and areal extent of wildfires declined since the early-to mid-1900s because of suppression activities. With the increased fuel loads in fire-suppressed areas, however, the trend since 1960 has changed, and the current extent of wildfires is approaching the early 1900s. This increase in postfire areas should benefit Lazuli buntings if these fires result in an increase in shrub vegetation.

Lazuli buntings are Neotropical migratory birds. The availability of suitable habitats used during migration, as well as their winter habitat, are critical components. Status of habitats, however, effects of nonhabitat factors on populations, and management practices in migratory and wintering areas are unknown.

Population status and trends--Recent BBS data indicate that the population was stable from 1968 to 1994 ($\bar{n} \geq 14$; $P < 0.10$) across the basin (Saab and Rich 1997). Sauer and others (1996) identified increasing trends for Lazuli buntings in the Western United States from 1980-1995 (+2.9 percent per yr, $\bar{n} = 147$; $P < 0.01$).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 18 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The results of our habitat trend analysis and the literature suggest the following issues are of high priority for group 18:

1. Altered frequency of stand-replacing fires.
2. Loss of shrub-dominated early seral vegetation types.
3. Loss and degradation of riparian vegetation.

Potential strategies--The issues suggest the following broad-scale strategies would be effective in supporting the long-term persistence of the Lazuli bunting. strategies would apply basin-wide.

1. (To address issue no. 1) Restore fire as an ecological process in the montane and lower montane community groups. Natural fire frequencies and intensities should be considered where appropriate.
2. (To address issue no. 2) Increase the representation of shrubs in the early seral stages of forest communities.
3. (To address issue no. 3) Reduce impacts to shrubs from grazing, recreation, and other activities.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Leave some postfire and postharvest areas unaltered to regenerate naturally.
2. (In support of strategy no. 2) Use prescribed fire to increase the representation of shrubs in the early seral stages of forest communities.
3. (In support of strategy no. 3) Remove or explicitly control the timing and intensity of grazing to develop and promote the long-term persistence of shrub communities.
4. (In support of strategy no. 3) Restrict activities in riparian areas that negatively affect riparian vegetation. Areas that currently support healthy shrub communities should be a priority for conservation.

GROUP 19--GRAY WOLF AND GRIZZLY BEAR

Results

Species ranges and source habitats--Group 19 consists of the grizzly bear and gray wolf. Historically these two species ranged across most of the basin (fig. 57), although use of lower elevations within the Northern Great Basin and Owyhee Uplands ERUs was probably incidental. This distribution has been greatly reduced, and both species currently persist only in small, disjunct populations. Gray wolf populations occur in western Montana, central Idaho, and western Wyoming; grizzly bear populations remain in the North Cascades, northern Idaho, western Montana, and western Wyoming (fig. 57).

Vol. 2, Figure 57--Ranges of species in group 19 within the basin.

The grizzly bear was listed as federally threatened under the ESA on July 28, 1975. The original recovery plan was approved in 1982 and amended in 1993. The northern Rocky Mountain gray wolf was listed as endangered on June 4, 1973, and a recovery plan was released in 1987 (USDI Fish and Wildlife Service 1987). Wolves have been state protected in Montana since 1975 and in Idaho since 1977 (USDI Fish and Wildlife Service 1987).

Source habitats for group 19 span a broad elevational range and include all terrestrial community groups except exotic herbland and agriculture. About 80 percent of all possible cover type-structural stage combinations are source habitats (vol. 3, appendix 1, table 1).

Source habitats for wolves must include suitable denning and rendezvous sites and a sufficient, year-round prey base of ungulates and alternate prey (USDI Fish and Wildlife Service 1987). Den sites are used for rearing pups and are typically near forested cover and removed from human activity. Wolves are sensitive to human disturbance near dens from mid-April to July (Weaver and others 1996). Rendezvous sites are resting and gathering areas used by wolf packs after the pups are mobile and typically include meadow vegetation and adjacent forest with resting sites under trees (USDI Fish and Wildlife Service 1987). Home ranges can be exceedingly large, based on estimates from radio telemetry--in Minnesota for example, home range estimates were from 49 to 135 km² (19 to 52 mi²) (van Ballenberg and others 1975), and in Alberta, winter home ranges fell between 357 and 1779 km² (138 and 687 mi²) (Fuller and Keith 1980). The principal foods of wolves in the Rocky Mountains are deer, elk,

and moose (USDI Fish and Wildlife Service 1987; Weaver 1994, cited in Weaver and others 1996).

Grizzly bear habitat selection is affected by (1) abundance and quality of foods; (2) gender-specific orientation to different nutrients; (3) reproductive status of females and concerns about security of dependent young; (4) presence and identity of other bears, especially adult males; and (5) presence of humans and prior contact with humans.⁵ Grizzly bears are omnivorous, but their use of certain high-quality foods with limited spatial or temporal distribution often results in seasonal shifts in habitat selection (Hamer and Herrero 1987; Mace and others 1996; Mattson and others 1991a, 1991b; McLellan and Hovey 1995; Servheen 1983). Also, food availability fluctuates among years, and habitat selection may therefore differ from one year to the next (Green and others 1997; Mattson and others 1991a, 1991b; McLellan and Hovey 1995).

A selection process also seems to be used for the location of dens for hibernation and the birth and rearing of young. Typical dens are either dug by bears or occur in natural cavities in subalpine, montane, and rock community groups. Den sites tend to be clustered, thereby suggesting that certain areas possess more favorable combinations of environmental factors for denning (USDI Fish and Wildlife Service 1993). Grizzly home ranges encompass large areas. For example, based on several studies, annual home ranges of males in the Northern Continental Divide Ecosystem are between 165 and 1406 km² (64 and 543 mi²), with an average of 489 km² (189 mi²) (USDI Fish and Wildlife Service 1993).

Broad-scale changes in source habitats--Source habitats for the grizzly bear and gray wolf likely occurred throughout the basin historically (fig. 58A). The current extent of habitat, albeit largely unoccupied, is similar to the historical distribution except for the Columbia Plateau, Lower Clark Fork, and Upper Clark Fork ERUs, where habitat is more patchily distributed than it was historically (fig. 58B).

Basin-wide, the overall trend in source habitats for group 19 was neutral (fig. 58C). Source habitats remained relatively stable in 9 of 13 ERUs (figs. 58C and 59). Fifty percent of all watersheds, located primarily in the southern half of the basin and along the western and northern borders, showed no trend in habitat (fig. 59). Source habitats were projected to have decreased in four ERUs: the Columbia Plateau, Lower Clark Fork, Upper Clark Fork, and Upper Snake (fig. 59).

Vol. 2, Figure 58--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 19.

Vol. 2, Figure 59--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 19, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Despite the overall neutral trend for source habitats for group 19, many of the terrestrial communities were projected to have changed dramatically from historical conditions. In general, mid-seral forests increased in areal extent basin-wide, whereas both early- and late-seral forests declined (Hann and others 1997). Some forest

cover types, including western white pine, whitebark pine, western larch, and limber pine no longer occur in stands large enough to map at the broad scale, whereas Pacific silver fir-mountain hemlock and western redcedar-western hemlock increased, respectively, 1,700 and 853 percent basin-wide (Hann and others 1997).

Within nonforest terrestrial communities, upland herbland and upland shrubland both strongly declined, whereas three new terrestrial communities, urban, agricultural, and exotic herbland, have emerged since the historical period (Hann and others 1997). Examples of declining nonforest cover types are native forb and mountain big sagebrush, which declined, respectively, by 91 and 34 percent basin-wide (Hann and others 1997). Within the four ERUs having overall declining trends in source habitats for group 19, declines were mostly in western white pine, whitebark pine, western larch, limber pine, big sagebrush, and native forb (vol. 3, appendix 1, table 4).

Other factors affecting the group--Human-caused mortality is the major factor limiting the recovery of wolves and grizzly bears (Fritts and Mech 1981; Knight and others 1988; Mattson and others 1996a, 1996b; Pletscher and others 1997; USDI Fish and Wildlife Service 1987, 1993). About 84 percent of all known mortalities of wolves on the Montana-British Columbia-Alberta border were human-caused, primarily legal shootings in Canada (Pletscher and others 1997). In the northern Rockies, between 1974 and 1996, 85 to 94 percent of all deaths of marked grizzly bears >1 year old were due to humans (Mattson and others 1996a).

For wolves, human-caused losses are due to shooting, trapping, and vehicle accidents (Fritts and others 1985). Six of the nine mortalities that occurred in the first 20 months after the reintroduction into Yellowstone National Park were human caused: three wolves were illegally shot, one was killed by Animal Damage Control personnel after repeated sheep depredations, and two were killed by vehicles (Bangs and Fritts 1996). In many cases, wolf mortalities are related to real and perceived depredations of livestock.

For grizzly bears, human-caused mortalities stem from (1) direct human-bear conflicts in wilderness areas and parks (for example, hikers, photographers, or hunters); (2) attraction of grizzly bears to improperly stored food or garbage; (3) attraction of grizzly bears to improperly disposed dead livestock; (4) chance interactions between livestock and grizzly bears; (5) increased human occupancy of grizzly bear habitat, causing increased interactions and stress; and (6) hunting (USDI Fish and Wildlife Service 1993). Legal hunting of grizzly bears no longer occurs in the basin, but grizzly bears are taken by poachers and occasionally are mistakenly shot during the black bear hunting season.

Wolves, particularly juveniles, are susceptible to canine parvovirus and distemper, and these diseases could affect recovery in the northern Rocky Mountains if not monitored (USDI Fish and Wildlife Service 1987). Parasites and diseases do not appear to be significant causes of natural mortality of grizzly bears (Jonkel and Cowan 1971, Rogers and Rogers 1976, both cited in USDI Fish and Wildlife Service 1993).

Both species are negatively affected by roads. Roads per se are not a physical barrier; wolves use gated roads as travel corridors (Thurber and others 1994), and grizzly bears in Montana exhibit neutral or positive selection for areas with roads having <10 vehicles per day (Mace and others 1996). However, roads usually increase human presence and the likelihood of negative contacts. A disproportionate number of human-caused mortalities occur near roads, both for wolves (Mech 1970, as cited in Frederick 1991) and grizzly bears (Mattson and others 1996a). These mortalities are mostly legal and illegal shootings resulting from human access provided by roads (Mace and others 1996, McLellan and Shackleton 1988); vehicle collisions also play a role (Bangs and Fritts 1996, Knight and others 1988). Thurber and others (1994) cited three studies (Jensen and others 1986, Mech and others 1988, Thiel 1985) indicating wolf packs would not persist where road densities exceeded about 1.0 mi per mi² (0.6 km per km²).

An additional, indirect effect of roads is that road avoidance leads to underutilization of habitats that are otherwise high quality. Mace and others (1996) found that grizzly bears in Montana avoided roads having >10 vehicles per day. In southeastern British Columbia, grizzly bears underutilized about 9 percent of available habitats by avoiding areas 100 m (328 ft) from roads, regardless of traffic volume (McLellan and Shackleton 1988). Several other studies have documented road avoidance by grizzly bears in or near the basin (Green and others 1997, Kasworm and Manly 1990, Mattson and Reinhart 1997, Mattson and others 1987). Similar effects have been observed with wolves: packs in the Great Lakes region avoided habitats with high road and human densities even though densities of deer, a principal prey, were also high in these areas (Mladenoff and others 1995). In northern Montana, wolf travelways were at least 4 to 22 km (2.5 to 13.6 mi) from the nearest driveable road, which precluded their use of high-quality habitats and food resources (Singer 1979).

Road access also increases the likelihood of habituation to humans. Individual wolves and grizzly bears can become accustomed to human presence, leading to nuisance situations that can result in the death of the habituated animal (Mattson and others 1992, Meagher and Fowler 1989).

The neutral trends in source habitats projected for the basin do not reflect loss of habitat effectiveness because of roads and human activities. Road densities in the basin have substantially increased from historical levels and are estimated to be moderate to high in most ERUs (Hann and others 1997). Moreover, the human population in the basin has increased and is estimated currently at 3 million (McCool and others 1997). The increase in road densities and human population are believed responsible for the unoccupied state of many source habitats of grizzly bears and wolves in the basin. For example, Merrill and others (1999) included roads, level of human activity, and distance and size of nearby human populations in their model of environmental suitability for grizzly bears in Idaho, and found that suitable environments did not exceed 37 100 km² (14,342 mi²) under any model assumptions.

The demographic impact of human-caused mortality is intensified for grizzly bears by their low reproductive rate. Litters range from one to four cubs with an average of two, and females generally do not begin to reproduce until 5.5 yrs old (USDI Fish and Wildlife Service 1993). Each female has the limited potential of adding three to four females to a population during her lifetime (USDI Fish and Wildlife Service 1993). Using this demographic information in conjunction with behavioral plasticity in food acquisition and dispersal capabilities, Weaver and others (1996) concluded that grizzly bears have fairly low resiliency to human disturbances, whereas gray wolves, based on these same factors, are moderately resilient.

Lack of connectivity among habitat reserves is a major factor affecting the long-term persistence of grizzly bears, and perhaps also wolves (Noss and others 1996). Source habitats are currently fragmented by human disturbances to a level where interchange within the entire regional population occurs rarely if at all (Noss and others 1996). Small, isolated populations are susceptible to extirpation from inbreeding, chance breeding events (for example, no female births in a given year), and environmental uncertainty (for example, drought or disease) (Shaffer 1981). This appears to be a concern for small, isolated grizzly bear populations (Allendorf and others 1991, cited in Mattson and others 1996b). Insufficient connectivity among local populations reduces the likelihood of recolonization once a population has been extirpated. The Bitterroot ecosystem is an example of a recent extirpation with extremely low probability of recolonization because of lack of connectivity with other grizzly bear populations (Merrill and others 1999).

Ultimately, human attitudes towards wolves and grizzly bears is what will ensure their survival or extirpation (Bangs and Fritts 1996, Mattson and others 1996a). Many of the negative effects of roads and human activities could be diminished through changes in human attitudes and behavior (Mattson and others 1996a, 1996b).

Population status and trends--Wolf populations were reduced to near extinction within the basin during the 1800s to early 1900s (USDI Fish and Wildlife Service 1987). Wolf numbers have increased, however, within the last 10 years. In addition to natural recolonizations of historical habitats in Washington, Idaho, and northwestern Montana (Marcot and others 1997), wolves have been reintroduced to central Idaho and the Yellowstone area as nonessential experimental populations (Federal Register 1994) beginning in 1995. Natural and experimental populations are currently doing well in all three areas identified for recovery: northwestern Montana, north-central Idaho, and the Greater Yellowstone Ecosystem. As of 1997 (USDI Fish and Wildlife Service 1997), northwestern Montana had about 75 wolves and 6 confirmed breeding pairs; central Idaho contained 75 to 80 wolves with 8 to 11 breeding pairs; and the Yellowstone ecosystem contained 85 to 90 wolves and 8 breeding pairs.

Between 1800 and 1975, grizzly bear populations in the lower 48 States receded from estimates of over 100,000 to <1,000 bears (USDI Fish and Wildlife Service 1993). Extirpations within the basin include Utah (1923) and Oregon (1931) (USDI Fish and Wildlife Service 1993). The Interagency Grizzly Bear Committee⁶ identified five recovery zones south of Canada where grizzly bears and grizzly habitat are managed for recovery, and within which the population parameters will be monitored (Interagency Grizzly Bear Committee 1998). The recovery zones are referred to as ecosystems to emphasize the ecological rather than jurisdictional nature of their boundaries (USDI Fish and Wildlife Service 1993). Four of the recovery zones are within the basin--the Northern Cascades, Selkirk, Cabinet-Yaak, and Northern Continental Divide ecosystems--and the fifth, the Yellowstone ecosystem, occurs on the eastern border of the basin. The Selway-Bitterroot ecosystem is under consideration as a recovery zone, as outlined in the Draft Environmental Impact Statement for Grizzly Bear Recovery in the Bitterroot Ecosystem (USDI Fish and Wildlife Service, 1997).

Grizzly bear population estimates currently are available only for the Northern Continental Divide Grizzly Bear ecosystem (440 to 680 bears) (USDI Fish and Wildlife Service 1993) and the Yellowstone ecosystem (280 to 610 bears) (Eberhardt and Knight 1996). The Selkirk Mountains and Cabinet-Yaak ecosystems are believed to have breeding populations based on sightings of females with young, but populations within each ecosystem may be less than 20 grizzly bears (Knick and Kasworm 1989, Wielgus and Bunnell 1995). Population status within the Northern Cascades is unknown (Interagency Grizzly Bear Committee 1998, USDI Fish and Wildlife Service 1993). No grizzly bears currently live in the Bitterroot Mountains of Idaho (Interagency Grizzly Bear Committee 1998).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 19 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The following issues have been identified as major challenges to the conservation of the grizzly bear and gray wolf:

1. Excessive mortality from conflicts with humans.
2. Excessive mortality related to the presence of roads (accidents, poaching, and increased conflicts).
3. Displacement from suitable habitats because of human activities.
4. Isolation of populations within each recovery area.

The goal of the revised Grizzly Bear Recovery Plan is to identify actions necessary for the conservation

and recovery of the grizzly bear and to remove the grizzly bear from threatened status in each recovery zone (USDI Fish and Wildlife Service 1993). The goal of the recovery plan for gray wolves is to remove the Northern Rocky Mountain wolf from the endangered and threatened species list by securing and maintaining a minimum of 10 breeding pairs of wolves in each of the three recovery areas for a minimum of 3 successive years (USDI Fish and Wildlife Service 1987).

Potential strategies--The following strategies could be used in the Northern Cascades, Northern Glaciated Mountains, Upper Clark Fork, Lower Clark Fork, Central Idaho Mountains, and Snake Headwaters ERUs to support recovery of the gray wolf and grizzly bear:

1. (To address issue no. 1) Reduce the prevalence of conflict situations and the number of human-caused mortalities of bears and wolves. Provide secluded habitats that reduce the potential for conflicts with humans.
2. (To address issue no. 2) Develop a policy for road construction, maintenance, and obliteration on public lands within gray wolf and grizzly bear recovery areas and in source habitats that surround and could potentially connect these habitats.
3. (To address issue no. 3) Reduce human activities in important grizzly bear foraging areas and around known wolf dens.
4. (To address issue no. 4) Provide interregional habitat connectivity across all ERUs with wolf and bear populations (Northern Cascades, Northern Glaciated Mountains, Upper Clark Fork, Lower Clark Fork, Central Idaho Mountains, and Snake Headwaters).

Practices that support strategies--Action items and practices for the recovery of the gray wolf and grizzly bear are in the Northern Rocky Mountain Wolf Recovery Plan (USDI Fish and Wildlife Service 1987), the Grizzly Bear Recovery Plan (USDI Fish and Wildlife Service 1993), the Interagency Grizzly Bear Guidelines (Interagency Grizzly Bear Committee 1986) and the Grizzly Bear Compendium (LeFranc and others 1987). The following practices have been drawn from these documents as examples, and would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Alter the timing and location of livestock grazing to reduce the need for wolf and grizzly bear depredation control.
2. (In support of strategy no. 1) Implement sanitation practices, including law enforcement to support these practices, to minimize the likelihood of grizzly bear attraction to human food, garbage, and dead livestock.
3. (In support of strategy no. 1) Increase extent and scope of public education programs regarding the role of human-bear and human-wolf conflicts in the conservation of these species.
4. (In support of strategy nos. 1 and 2) Minimize or avoid road construction within unroaded areas in grizzly bear ecosystems and wolf recovery areas. Obliterate or restrict use of roads in important seasonal habitats, such as low-elevation riparian areas (spring habitat for grizzly bears).
5. (In support of strategy nos. 1 and 3) Reduce or temporarily discontinue activities such as livestock grazing, timber harvests, backcountry use, mining, and oil and gas development in important grizzly bear foraging areas during peak foraging periods. Restrict human access near wolf dens from April 15 to July 1.
6. (In support of strategy no. 4) Use concepts described in Noss and others (1996) to design habitat

connectivity among recovery areas. Identify existing and potential dispersal corridors for wolves and bears, and seek opportunities with all landowners and affected parties to modify the timing, intensity, and location of human activities within these corridors.

GROUP 20--MOUNTAIN GOAT

Results

Species ranges, source habitats, and special habitat features--Group 20 consists of the mountain goat, a year-round resident of the basin. Within the basin, the mountain goat occurs in the mountains of central and northeast Washington, northeast Oregon, central and northern Idaho, and western Montana. These areas correspond to five ERUs: the Northern Cascades, Northern Glaciated Mountains, Lower Clark Fork, Upper Clark Fork, and Central Idaho Mountains (fig. 60). The range also includes small, bordering areas of the Southern Cascades and Columbia Plateau ERUs (fig. 60). Most populations are native, but mountain goats have been introduced into portions of Montana, and reintroduced into the Elkhorn and Blue Mountains of Oregon. Although the Hells Canyon population stems from a transplant, recent archeological evidence suggests historical occupancy of the Hells Canyon area and the Wallowa Mountains (Matthews and Coggins 1994).

Vol. 2, Figure 60--Ranges of species in group 20 within the basin.

Source habitats for mountain goats include 15 cover types within six community groups: alpine, subalpine forest, montane forest, lower montane forest, upland shrubland, and rock-barren (vol. 3, appendix 1, table 1). Mountain goats show no apparent preference for any cover type, as long as they occur on steep terrain or near cliffs and talus. Mountain goats seem to use all structural stages within forested cover types except for the stem-exclusion stage of montane and lower montane forests (vol. 3, appendix 1, table 1). Upland shrublands provide important foraging habitat, and forests provide both foraging habitat and protection from inclement weather (Johnson 1983).

Special habitat features identified for mountain goats are cliffs, talus, and seasonal wetlands (vol. 3, appendix 1, table 2). Cliffs and talus are central to mountain goat distribution and habitat use (Hjeljord 1973). Cliffs provide escape terrain from predators (Johnson 1983, Rideout 1978), and both cliffs and talus provide foraging areas with little competition from other herbivores (Rideout 1978).

Mountain goats forage on various plant species depending on local and seasonal availability. Grasses and sedges comprise a major portion of the diet in most locales (Adams and Bailey 1983, Hjeljord 1973, Saunders 1955), along with mosses, lichens, ferns, and shrubs (Rideout 1978). Mountain goats exhibit localized shifts in habitat use in response to changes in food availability because of snow accumulation, moisture, wind, and solar exposure (Rideout 1978). Mountain goats are subject to predation from mountain lions, golden eagles (Rideout 1978), wolves, and grizzly bears (Smith 1986, Smith and others 1992).

Broad-scale changes in source habitats--The following trends in source habitats for mountain goats were derived without reference to the proximity of cliffs and talus and therefore include habitat patches that are not available to mountain goats. Trends derived from a restricted subset of habitats near cliffs could differ substantially in magnitude from those reported here, but the general direction of the trends likely

would remain the same.

The historical distribution of source habitats was essentially the same as it is now, occurring in the mountains of central and northeast Washington, northeast Oregon, central and northern Idaho, and western Montana (figs. 61A, B). Because mountain goats use various cover types, trends in the extent of source habitats differed basin-wide. Trends were projected to be neutral in 32 percent of the watersheds and positive in 42 percent of the watersheds basin-wide (fig. 62). Positive trends were projected in more than 50 percent of watersheds in the Blue Mountains and Central Idaho Mountains ERUs, and declining trends were most prevalent in the Lower Clark Fork and Upper Clark Fork ERUs (figs. 61C and 62). All other ERUs with source habitats exhibited mixed trends.

Source habitats for mountain goats were most prevalent in the Northern Cascades ERU historically, and this has not changed. The area occupied by source habitats in this ERU comprised 51 percent of the weighted area of watersheds included in mountain goat range during both time periods (vol. 3, appendix 1, table 3).

Vol. 2, Figure 61--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 20.

Vol. 2, Figure 62--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 20, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Neutral trends in source habitats were partly because alpine and rock-barren community groups did not change in areal extent from historical to current periods (Hann and others 1997; vol. 3, appendix 1, table 4). Within other community groups, neutral trends resulted from declines in some cover types that were offset by increases in other cover types used as source habitats. For example, in the Northern Cascades ERU, a major transition occurred from interior ponderosa pine to both interior Douglas-fir and grand fir-white fir (Hann and others 1997), but this resulted in static trends in habitat extent because all three cover types are source habitats (vol. 3, appendix 1, table 1).

Declining trends in the Lower Clark Fork ERU were due to total losses at the broad scale of old forests of interior ponderosa pine, as well as declines in the stand-initiation stage of lodgepole pine and Engelmann spruce-subalpine fir (vol. 3, appendix 1, table 4). Declines in the Upper Clark Fork were chiefly because of nearly total losses of interior Douglas-fir and interior ponderosa pine old forests (vol. 3, appendix 1, table 4). Although less extensive in area, strong declines in whitebark pine old forests also occurred in both the Lower and Upper Clark Fork ERUs (vol. 3, appendix 1, table 4). In the Central Idaho Mountains, increases in source habitat were primarily due to areal increases in Engelmann spruce-subalpine fir, grand fir-white fir, interior Douglas-fir, lodgepole pine, mountain mahogany, and shrub or herb-tree regeneration (vol. 3, appendix 1, table 4). Increases in the Blue Mountains were associated mostly with

increases in grand fir-white fir (Hann and others 1997).

Condition of special habitat features--The areal extent of cliffs and talus has not changed between historical and current periods (Hann and others 1997). Seasonal wetlands are highly dependent on annual hydrologic cycles and therefore have fluctuated widely in occurrence and productivity over time.

Other factors affecting the group--Young of the year and yearlings incur the highest mortality rates, primarily because of harsh weather in conjunction with predation, internal parasites, and diseases (Johnson 1983). Adults are highly susceptible to hunting mortality, both legal and illegal (Kuck 1977, Matthews and Coggins 1994, Smith 1986, Swenson 1985).

Human activities disrupt mountain goats and can cause displacement from source habitats. Low-flying aircraft cause mountain goats to run, take alert defense postures, or take refuge under trees (Chadwick 1973). Road blasting and sonic booms also cause defensive reactions in mountain goats (Chadwick 1973). Mountain goats can become habituated to human disturbance, especially where they are not hunted, as in Glacier National Park (Pedicivillano and others 1987, Singer and Doherty 1985), but more typically, mountain goats exhibit signs of stress when exposed to human disturbances. In Montana's Rocky Mountain Front, mountain goat reproduction and kid survival was lower in a herd exposed to much human activity (such as energy exploration, a downhill ski resort, and developed recreation) compared to a herd in a more remote area (Joslin 1986).

Timber harvests can have both positive and negative effects on mountain goats. Overstory removal can increase forage productivity in areas where fire suppression has reduced the extent of open habitats (Johnson 1983). Sufficiently large stands of mature forests, however, must be retained for winter cover (Johnson 1983). Timber harvests also increase human access to mountain goat habitat through road construction (Chadwick 1973), and this has led to increased hunting mortality in some herds that were formerly less accessible (Johnson 1983).

Roads, particularly highways, also increase mortality rates through vehicle collisions (Singer 1978). In Glacier National Park, however, highway mortality was reduced by placing two highway underpasses on Highway 2 to allow goats to reach two mineral licks (Pedicivillano and others 1987).

Many goat populations are small because of habitat fragmentation, hunting pressure, and the establishment of new herds with few individuals. A potential consequence of low numbers is a high probability of deleterious effects from inbreeding. For example, even after hunting of the Wallowa Mountain goat population was discontinued, the population remained static for many years until new genetic stock was introduced in the 1980s (Matthews and Coggins 1994).

Population status and trends--Mountain goat population trends differ across the basin. Populations in Washington state⁷ and Montana⁸ have declined, whereas populations in the Wallowa and Elkhorn Mountains in northeastern Oregon have increased (Matthews and Coggins 1994). Native populations in Idaho have decreased, whereas introduced populations are stable or increasing.⁹

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 20 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--Important issues affecting mountain goats were taken both from the literature and our habitat analysis.

1. Increased human disturbance in formerly isolated habitats.
2. Reduction in forage quantity and quality because of successional changes in source habitats from fire suppression.
3. Habitat fragmentation because of human land uses and successional changes in source habitats from fire suppression.

Potential strategies--

1. (To address issue no. 1) Reduce human activities, particularly where mountain goat herds are static or declining.
2. (To address issue no. 2) Restore quality and quantity of forage where forage has declined because of successional changes and changes caused by fire suppression.
3. (To address issue no. 3) Seek opportunities to reduce fragmentation in historical range caused by human land uses and fire suppression.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Incorporate mitigation measures for human activities within or adjacent to known mountain goat herds into all relevant planning documents.
2. (In support of strategy no. 1) Carefully regulate frequency and height of low-flying aircraft over known mountain goat herds, including military exercises, helicopter logging, recreational flights, and wildlife surveys.
3. (In support of strategy nos. 2 and 3) Use understory thinning and prescribed burns to improve the quantity and quality of forage, and increase links with isolated herds.
4. (In support of strategy no. 3) Use land acquisitions, exchanges, and easements to consolidate blocks of suitable mountain goat habitat, including blocks of currently unoccupied habitat.

GROUP 21--LONG-EARED OWL

Results

Species ranges, source habitats, and special habitat features--Group 21 is comprised of the long-eared owl. Long-eared owls are year-round residents of the basin, but some individuals move long distances suggestive of migratory behavior during fall and spring (Marks and others 1994). The current range of the long-eared owl includes all 13 ERUs (fig. 63).

Vol. 2, Figure 63--Ranges of species in group 21 within the basin.

Source habitats for the long-eared owl include a broad range of vegetation types from mid-elevational forests to low-elevational shrublands. The six vegetation community groups in which source habitats occur are montane forests, upland woodlands, upland shrublands, upland herblands, riparian woodlands, and riparian shrublands (vol. 3, appendix 1, table 1). Source habitat cover types within the montane forest community include interior Douglas-fir, western larch, grand fir-white fir, Sierra Nevada mixed conifer, and red fir. Nearly all structural stages within these cover types except for managed young forests are considered source habitats.

Long-eared owls tend to nest and roost in dense vegetation, but they hunt almost exclusively in open habitats (Getz 1961, ICBEMP 1996h, Marks and others 1994, Thurow and White 1984). As such, they are considered a contrast species (vol. 3, appendix 1, table 2), requiring a juxtaposition of contrasting vegetative structure to meet all aspects of their ecology. Where forests are adjacent to open areas, trees are typically used for nest sites. Where forests are not present, nests are placed in tall shrubs (Holt 1997). This owl typically lays its eggs in abandoned stick nests of other species, especially common raven, American crow, and black-billed magpie nests (Marks and others 1994).

Broad-scale changes in source habitats--The historical distribution of source habitats was most concentrated in the Columbia Plateau, Northern Great Basin, and Owyhee Uplands (fig. 64A). The current distribution is about the same (fig. 64B), although significant declines have occurred in the northern half of the Columbia Plateau and in the eastern basin, and significant increases have occurred in the north, the central basin, and in the southwest (fig. 64C).

Trends in extent of source habitats are mixed across the basin: 29 percent of watersheds with source habitats showed no change in areal extent between the historical and current periods; 40 percent of watersheds had declining trends, and 31 percent had increasing trends (fig. 65). Four ERUs had declining and strongly declining trends in source habitats in >50 percent of watersheds. These were the Columbia Plateau (53 percent of watersheds), the Upper Clark Fork (75 percent of watersheds), the Upper Snake (76 percent of watersheds), and the Snake Headwaters (67 percent of watersheds). Increasing and strongly increasing trends occurred in > 50 percent of watersheds in the Upper Klamath (63 percent of watersheds) and Blue Mountains (52 percent of watersheds) ERUs, and the Southern Cascades had increasing trends in 9 percent of watersheds (figs. 64C and 65).

Vol. 2, Figure 64--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 21.

Vol. 2, Figure 65--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 21, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Most vegetation types that provide source habitats for the long-eared owl have changed in extent from the historical period, but these changes have resulted in no net increase or decrease in source habitats.

Within the Upper Klamath, Blue Mountains, and Southern Cascades ERUs, increases in source habitats were largely due to increases in interior Douglas-fir, grand fir-white fir, juniper/sagebrush woodland, and big sagebrush (Hann and others 1997; vol. 3, appendix 1, table 4). Declines in the northern portion of the Columbia Plateau and the Upper Snake are primarily due to transitions from big sagebrush to agriculture and the conversion of many cover types in the upland shrubland and riparian shrubland community groups to exotic forbs-annual grass (Hann and others 1997; vol. 3, appendix 1, table 4). Declines in the Upper Clark Fork are due to areal increases in cropland and Engelmann spruce-subalpine fir cover types (Hann and others 1997), neither of which are source habitats for the long-eared owl, and areal declines in all structural stages of interior Douglas-fir (Hann and others 1997). Declines in the Snake Headwaters are due to transitions in both the upland herbland and upland shrubland communities to agriculture (Hann and others 1997).

Condition of special habitat features--No special habitat features were identified for the long-eared owl. The amount of edge habitat, however, may be a landscape-level variable of some importance to long-eared owls. The mid-scale analysis of vegetation changes in the basin (Hessburg and others 1999) indicated that the amount of edge increased significantly in 6 of 13 ERUs. Assuming that this scale of analysis is appropriate for long-eared owls, and assuming that interspersed habitats is beneficial to this species, the increase in edge is considered a positive change in habitat condition.

Other factors affecting the group--The long-eared owl generally nests in trees, using stick nests created by other bird species, especially common raven, American crow, and black-billed magpie. Programs designed to reduce these species could therefore have a negative effect on the long-eared owl.

Little is known about effects of pesticides on this species. Henny and others (1984) discovered organochlorine residues in one-third of all long-eared owl eggs they examined.

Roads apparently do not impact long-eared owls. Mean distance to nearest road was not different for successful and unsuccessful nests (Marks 1986).

Population status and trends--Long-eared owls are common in most Western states, although they are considered rare in Montana (Craig and Trost 1979). Long-eared owl numbers appear to be stable in most states (Marti and Marks 1989). Within the basin, populations seem to attain peak densities in southern Idaho (Craig and Trost 1979).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 21 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The primary issue related to long-eared owl conservation is degradation and loss of native upland shrublands, riparian shrublands, and riparian woodlands.

Potential strategies--

1. Maintain and restore native upland shrublands, riparian shrublands, and riparian woodlands across the basin, particularly in the northern half of the Columbia Plateau and in the Upper Snake and Snake Headwaters ERUs.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. Limit livestock grazing and recreational activities in riparian shrublands and woodlands to allow growth of dense vegetation for nest sites.
2. Explore options under the Conservation Reserve Program (CRP) (Johnson and Igl 1995), or develop other incentive programs, to encourage restoration of agricultural areas to native cover types.
3. Restore native vegetation by appropriate treatments and seedings of native shrub, grass, and forb species.

GROUP 22--CALIFORNIA BIGHORN SHEEP AND ROCKY MOUNTAIN BIGHORN SHEEP

Results

Species ranges, source habitats, and special habitat features--Group 22 consists of two subspecies of bighorn sheep, the California and Rocky Mountain bighorn sheep; both are year-round residents of the basin. Although they use similar habitats, the two subspecies are separated by disparate ranges of remnant populations and by different geographic areas that have been designated for their reintroduction. In general, California bighorn occur in the western and southern portions of the basin, and Rocky Mountain bighorn occupy the eastern and northern portions of the basin (fig. 66).

Vol. 2, Figure 66--Ranges of species in group 22 within the basin.

Historically, California bighorns occurred in central and southeastern Oregon, the eastern slope of the Cascade Range in Washington, northwestern Nevada, and the mountains of southwestern Idaho (fig. 66). Populations declined in the late 1800s and bighorns were extirpated from all four states between 1900 and 1930 (Thorne and others 1985). Because of a series of reintroductions, California bighorns currently are found in many disjunct populations within their former range (fig. 66).

Rocky Mountain bighorns historically occurred in northeastern Oregon, central Idaho, Montana and Wyoming, and northeastern Nevada (Thorne and others 1985) (fig. 66). After a severe population decline in the early 1900s, bighorns remained in only a few isolated areas of their former habitat. The current range represents an increase in occupied habitat since that time, because of a combination of reintroductions and protection of remnant populations (Thorne and others 1985). However, much of the historical range is still unoccupied (fig. 66).

Source habitats for both subspecies are primarily in the alpine, subalpine, upland shrubland, and upland herbland community groups. Old-forest and stand-initiation stages of whitebark pine are source habitat, but only the stand-initiation stage of other forest cover types are used (vol. 3, appendix 1, table 1). Bighorn sheep prefer open habitats with short vegetation, both for high-quality forage (McWhirter and others 1992) and to maintain high visibility for predator avoidance (Risenhoover and Bailey 1985, Wishart 1978), and a negative correlation between forest cover and bighorn occurrence has been observed (Bentz and Woodard 1988). Postfire habitats can benefit bighorn sheep by improving forage quality (McWhirter and others 1992) and increasing visibility (Bentz and Woodard 1988).

In the basin, Rocky Mountain bighorn sheep exhibit more seasonal movements than California bighorn

sheep. Alpine and subalpine community groups are primarily summer range for the Rocky Mountain subspecies, whereas upland herbland and shrubland are used in both seasons, depending on elevation (vol. 3, appendix 1, table 1).

Special habitat features identified for these two subspecies include cliffs, talus, and seasonal wetlands (vol. 3, appendix 1, table 2). The location of cliffs and talus ultimately define the distribution of bighorn sheep because they are essential for escape cover and the secure rearing of young (Wakelyn 1987). Cover types listed as source habitats (vol. 3, appendix 1, table 1) generally are not available to bighorns unless they are near cliffs.

Broad-scale changes in source habitats--The following trends in source habitats for bighorn sheep were derived without reference to the proximity of cliffs and talus and may not accurately represent changes in the more restricted subset of stands available to bighorns. Trends derived from a restricted subset of habitats could differ substantially in magnitude from those reported here, but the general direction of the trends likely would remain the same.

Source habitats (regardless of proximity to cliffs) currently occupy the same general geographic extent as the historical distribution of habitats but are less prevalent within each watershed (figs. 67A, B), thereby resulting in overall negative trends in habitat extent. Many areas that formerly had bighorn sheep habitat in 25 to 50 percent of each watershed now meet source habitat conditions in less than 25 percent of each watershed, particularly in the central and northern regions of the basin (fig. 67B). Habitats declined in 57 percent of the watersheds throughout the basin and in most watersheds in five ERUs: the Blue Mountains, Northern Glaciated Mountains, Lower and Upper Clark Fork, and Upper Snake (fig. 68). Declining trends also were noted in the Northern and Southern Cascades, but these ERUs are on the western edge of the geographic range and contain little habitat (vol. 3, appendix 1, table 3). Most watersheds of the Northern Great Basin and Owyhee Uplands ERUs exhibited no change in the amount of source habitats, whereas watersheds in the Snake Headwaters exhibited mixed trends in habitat extent (fig. 68).

Vol. 2, Figure 67--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 22.

Vol. 2, Figure 68--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 22, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Declines in source habitats were due primarily to declines in big sagebrush, mountain big sagebrush, fescue-bunchgrass, interior ponderosa pine, native forb, western larch, wheatgrass-bunch grass, whitebark pine-subalpine larch, and whitebark pine (vol. 3, appendix 1, table 4). A notable change that has affected bighorn sheep is the widespread conversion of native shrublands and grasslands to agricultural cover types (Hann and others 1997), particularly in historical winter range. Also, source habitats with high visibility for predator

avoidance have been replaced by stands with reduced visibility, primarily through the transition of whitebark pine old forests to Engelmann spruce-subalpine fir and the transition of stand-initiation stage forest cover types to mid-seral stages (Hann and others 1997).

Condition of special habitat features--Cliffs and talus (represented by the community group rock-barren) have not changed between historical and current periods (Hann and others 1997). Cliffs and talus can be significantly altered through direct human disturbance such as blasting and road construction, but this type of activity generally has not occurred in remote areas used by bighorn. Seasonal wetlands are highly dependent on annual hydrologic cycles and therefore have fluctuated widely in occurrence and productivity over time.

Other factors affecting the group--Bighorn sheep are highly susceptible to pneumonia after exposure to bacteria (Pasteurella spp.), viruses (Parainfluenza type-3), lungworm, and stress agents (Foreyt 1994, Wishart 1978). Major reductions or total extirpation of bighorn herds because of pneumonia outbreaks are well documented (Cassirer and others 1996, Coggins 1988, Onderka and Wishart 1984, Spraker and others 1984). A recent episode of Pasteurella-associated pneumonia in the Hells Canyon area resulted in a known loss of 327 bighorn sheep between November 1995 and March 1996, which represented 50 to 75 percent of four herds in Oregon and Washington (Cassirer and others 1996).

Abundant circumstantial evidence indicates that domestic and exotic sheep are the source of nonendemic bacteria and viruses predisposing bighorn sheep to pneumonia (Coggins 1988, Foreyt and Jessup 1982, Martin and others 1996); moreover, direct evidence recently has been acquired through experimental contact between sheep and bighorns in enclosures (Foreyt 1994), and through bacterial swab cultures and DNA analysis of Pasteurella spp. collected from free-ranging bighorn sheep with pneumonia in Nevada and Oregon (Rudolph and others, in prep.). Domestic goats also may be reservoirs, although the evidence is less compelling. A feral goat was associated with diseased bighorn at the start of the outbreak in Hells Canyon and had genetically identical Pasteurella to one of the bighorn ewes; however, these bacteria were not common among bighorns sampled during the episode (Cassirer and others 1996; Rudolph and others, in prep.).

Bighorn sheep also are affected by grazing competition from livestock (USDI Bureau of Land Management 1995). Intensive grazing pressure that occurred between the late 1800s and early 1900s is believed a factor in the reduction in bighorn sheep populations of that era (Johnson 1983). Grazing competition with domestic sheep has been reduced in recent times because of efforts to maintain buffers between sheep and bighorns to reduce the potential for disease transmission. The leading source of grazing competition is from cattle (Blood 1961, Demarchi 1965, and Lauer and Peek 1976, as cited in Van Dyke and others 1983). Late winter grazing by cattle, however, has proven beneficial to the Lower Imnaha bighorn herd in Oregon.¹⁰

The condition of bighorn sheep habitats has been altered over the last century because of changes in historical fire regimes. Fire suppression has resulted in an increase in the density of trees of formerly open stands, reducing forage quality and causing bighorns to avoid these areas because of reduced visibility. Some cliff areas are currently inaccessible to bighorns because the stands of open timber through which bighorns formerly traveled have developed into dense stands that bighorns avoid (Wakelyn 1987). For the Rocky Mountain bighorn, fire-suppressed stands have created barriers between historical winter and summer range, preventing occupancy of the total range even though each isolated range is currently suitable (Wakelyn 1987).

Some historical ranges have become fragmented by urban, mining, agricultural, and recreational developments (USDI Bureau of Land Management 1995). In some cases, this has created a barrier between seasonal ranges, as described above for fire-suppressed habitat. Additionally, fragmentation has resulted in habitat islands that can support only small, isolated herds (USDI Bureau of Land

Management 1995).

Direct disturbance by humans can affect bighorn sheep by shifting their distribution (Hamilton and others 1982, Hicks and Elder 1979) and by increasing physiologic stress (MacArthur and others 1979). Hunted populations generally react more strongly than nonhunted populations (Hamilton and others 1982, Hicks and Elder 1979). Among the human activities that elicit the strongest negative response are low-flying aircraft (helicopters and military air exercises). Hiking in lambing areas is also disruptive to bighorns (USDI Bureau of Land Management 1995). The human population in the basin has increased from a relatively small number of native people to 3 million (McCool and others 1997); therefore, the number of human disturbances in bighorn sheep habitat likely has increased.

Population status and trends--Bighorn sheep populations declined substantially throughout their geographic range in the late 1800s and early 1900s. Because of the establishment of hunting regulations, however, a better understanding of disease transmission, and concentrated reintroduction efforts throughout the West, bighorn numbers have steadily increased over the last 50 years (Thorne and others 1985). By 1995, many reintroductions of California bighorn resulted in the establishment of 6 herds in Idaho, 29 herds in Oregon, and 8 herds in Washington (USDI Bureau of Land Management 1995).

Populations of Rocky Mountain bighorn also have been widely reintroduced into their historical habitats within the basin. As of 1995, the reintroduced and native populations comprised 10 herds in Idaho, 9 herds in Oregon (1 which extends into Washington), 3 additional herds in Washington, and 9 herds in Montana (USDI Bureau of Land Management 1995).

Population trends differ by herd. Some reintroduced herds are still increasing; for example, the Pueblo Mountains herd in southeast Oregon currently numbers 130 and is still growing.¹¹ This herd was started with three reintroductions in 1976, 1980, and 1983 that totaled 40 animals (Coggins and others 1996). Some herds have static trends; for example, the Steens Mountain bighorn herd was started with 11 animals in 1960 (Coggins and others 1996) and increased to 275 (USDI Bureau of Land Management 1995), but currently numbers 250 and seems to be static for unknown reasons (see footnote 11). Several herds in the Hells Canyon area of Washington and Oregon have recently declined because of an outbreak of Pasteurella-associated pneumonia (Cassirer and others 1996).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 22 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--Issues were taken from the literature and the results of our habitat analysis for these two subspecies.

1. Incompatibility with domestic sheep and possibly domestic goats because of the potential for disease transmission and competition for forage.
2. Reduction in forage quantity and quality because of successional changes in source habitats.
3. Habitat fragmentation (poor juxtaposition of seasonal ranges as well as isolation of small herds) because of successional changes in source habitats.
4. Habitat fragmentation because of agricultural, industrial, and recreational development.

5. Disturbance and habitat displacement because of human activities such as low aircraft fly-overs and hiking in lambing areas.

Potential strategies--

1. (To address issue no. 1) Actively control the potential for disease transmission and forage competition between bighorns and domestic livestock.
2. (To address issue no. 2) Restore quality and quantity of forage where forage has declined because of successional changes in vegetation.
3. (To address issue no. 3) Restore habitat links between summer and winter range and access to escape cover that have been lost because of changes in historical fire regimes.
4. (To address issue no. 4) Seek opportunities to reduce fragmentation in historical range caused by human land uses.
5. (To address issue no. 5) Reduce human activities in key foraging and lambing areas.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Avoid direct contact between bighorn sheep and domestic sheep and goats. Guidelines established by the BLM for domestic sheep management in bighorn sheep habitats (USDI Bureau of Land Management 1995) recommend that buffers (having no domestic sheep or goats) are placed around bighorn sheep habitat and that bighorn sheep reintroductions do not occur in areas that have been grazed by domestic sheep or goats within the last 2 years.
2. (In support of strategy no. 1) Reduce forage competition with livestock by factoring bighorn sheep forage consumption into total forage utilization. Light to moderate cattle grazing during spring or early summer can be used to improve forage quality on bighorn sheep winter ranges (Bodie and Hickey 1980).
3. (In support of strategy nos. 2 and 3) Use understory thinning and prescribed burns to improve the quantity and quality of forage and to restore open habitat links between winter and summer ranges and to provide access to cliffs that currently are inaccessible to bighorns.
4. (In support of strategy no. 4) Use land acquisitions, exchanges, and easements to consolidate blocks of suitable bighorn sheep habitat (USDI Bureau of Land Management 1995).
5. (In support of strategy no. 5) Incorporate mitigation measures into all planning documents for mines, highways, canals, and recreational developments within or adjacent to occupied bighorn sheep range to minimize human disturbance.
6. (In support of strategy no. 5) Regulate activities that cause unacceptable disturbance to bighorns, such as flights of low-flying aircraft and back country recreation.

GROUP 23--RUFIOUS HUMMINGBIRD AND BROAD-TAILED HUMMINGBIRD

Results

Species ranges, source habitats, and special habitat features--Group 23 consists of the rufous hummingbird and the broad-tailed hummingbird, both of which are migratory breeders in the basin. The rufous hummingbird is distributed throughout forested portions of the basin (fig. 69), whereas the range of the broad-tailed hummingbird is restricted to small areas of Idaho and Montana (fig. 69). Both of these species are mostly associated with coniferous forests. The rufous hummingbird is found in 12 coniferous forest types and occurs in 53 combinations of forest types and structural stages. The broad-tailed hummingbird has source habitats in four coniferous types: Engelmann spruce-subalpine fir, interior Douglas-fir, grand fir-white fir, and interior ponderosa pine (vol. 3, appendix 1, table 1). Within the forest types, both species use old forests, understory reinitiation, and stand initiation. Source habitats for both species also include shrub-wetlands and aspen, and each species uses some woodland types. These species generally are found in more open forests, forests with openings, or in areas where open areas and forest habitats are adjacent because it is within these areas that the potential for herbaceous shrubs and herbs is higher. Herbaceous shrubs and herbs provide important foraging substrates (flowers) for these birds.

Vol. 2, Figure 69--Ranges of species in group 23 within the basin.

Both species typically nest in conifers in areas that support an abundance of nectar-producing flowers, which serve as a foraging substrate. Nectar-producing flowers are a special habitat feature for hummingbirds (vol. 3, appendix 1, table 2).

Broad-scale changes in source habitats--Historically, source habitats for group 23 were broadly distributed throughout the mountainous regions of the basin (fig. 70A). Currently, source habitats are still widely distributed but more concentrated in fewer watersheds in most of the ERUs (fig. 70B).

Overall, the projected trend in source habitats for group 23 declined from historical to present. Basin-wide, about 36 percent of the watersheds had strong declines in source habitats, and 19 percent had moderate declines (fig. 71). Eight ERUs were projected to have moderate or strong declines in source habitats in more than 50 percent of watersheds (fig. 71). More than 50 percent of the watersheds in the Upper Klamath and Northern Great Basin were projected to have moderate or strong increases from historical to present time (fig. 71). The Northern Cascades, Snake Headwaters, and Central Idaho Mountains generally had no change in amount of source habitats (fig. 71).

Vol. 2, Figure 70--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 23.

Vol. 2, Figure 71--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 23, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--The increase in source habitats in the Upper Klamath and Northern Great Basin is directly related to an increase in late-seral montane forests (vol. 3, appendix 1, table 4). Decreases in source habitats in six ERUs are due primarily to reductions in late-seral ponderosa pine, western larch, and western white pine. Six ERUs (Southern Cascades, Upper Klamath, Columbia Plateau, Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork) also showed substantial declines in early-seral forests, particularly ponderosa pine, western larch, and western white pine. Decreases in the Upper Snake resulted from declines in aspen (understory reinitiation) and chokecherry-serviceberry-rose. The decline in available source habitats in the Owyhee Uplands primarily was because of a decrease of about 2 percent in shrub-wetlands, but this figure may underrepresent the actual loss of habitat due to the small size of shrub-wetland patches relative to mapping unit size at the broad scale.

Condition of special habitat features--An analysis of the abundance of nectar-producing flowers, the primary food source for these hummingbirds, is not possible at the scale of this analysis, and no information on condition or trend is available. The increasing trend in shade-tolerant, multi-storied stands likely decreased the abundance of forest-associated flowers by reducing the amount of sunlight needed for flower development.

Other factors affecting the group--Grazing has an overall negative impact on nectarivores because of these species dependence on understory plants as a food source. Negative effects of grazing on broad-tailed hummingbirds have been documented in two studies (Page and others 1978, Schulz and Leininger 1991, cited in Saab and others 1995). Negative responses to grazing also were reported for the rufous hummingbird (Page and others 1978, cited in Saab and others 1995).

Because both species are Neotropical migratory birds, habitat used during migration and winter also may influence population trends. Russell and others (1994) observed that the quality of "stopover" habitats for migrant rufous hummingbirds differed widely because of the natural variation in flowering, and found a positive correlation between variation in flowering and hummingbird survival. Little is known on the abundance or trend of wintering habitat of these species.

Population status and trends--Based on BBS data from 1968 to 1994, rufous hummingbirds in the basin have shown stable population trends (Saab and Rich 1997). There are insufficient BBS data for the broad-tailed hummingbird to analyze population trends within the basin (Saab and Rich 1997). Specialized monitoring techniques are needed to track population trends for both species of hummingbirds.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 23 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The following issues are drawn from our analysis of source habitat trends in combination with issues identified from other literature:

1. Decline in abundance of natural forest openings specifically within ponderosa pine, interior Douglas-fir, grand fir, and western larch. There also has been a nearly complete loss of open forests of western white pine (all structural stages).
2. Decline in abundance of forest-associated flowering plants because of exclusion of fire, establishment of shade-tolerant trees, and subsequent decrease in shrub and herbaceous understories.
3. Decline in abundance of understory flowering shrubs, particularly in riparian areas, because of cattle grazing.

Potential strategies--Habitat for rufous and broad-tailed hummingbirds would benefit from the following strategies that address the issues listed above:

1. (To address issue no. 1) Promote the development of forest openings and single-layered old forest structures of ponderosa pine, interior Douglas-fir, grand fir, and western larch, particularly in the ERUs where source habitats have declined (Southern Cascades, Columbia Plateau, Blue Mountains, Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork).
2. (To address issue no. 1) Increase the amount of early seral forests in the ERUs where it has declined (Southern Cascades, Upper Klamath, Columbia Plateau, Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork).
3. (To address issue no. 2) Restore fire as an ecological process to encourage development of forest openings and growth of shrubs and forbs.
4. (To address issue no. 3) Reduce impacts to flowering herbs and shrubs from grazing.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Remove shade-tolerant understory trees to promote stand health and longevity in old-forest stands. Hand removal, or in some cases prescribed burning, may be effective.
2. (In support of strategy nos. 2 and 3) Accelerate development of flowering shrubs and forbs with the use of prescribed underburning and thinning, or allow for natural wildfires to occur particularly in the following ERUs: Southern Cascades, Columbia Plateau, Blue Mountains, Northern Glaciated Mountains, Lower Clark Fork, Upper Clark Fork, Owyhee Uplands, and the Upper Snake.
3. (In support of strategy nos. 2 and 3) Select areas that have been burned by wildfire or harvested for timber and try to extend the duration of the seral stage, which is rich in forbs and shrubs, by not planting conifers. Areas of primary importance are the Southern Cascades, Upper Klamath, Columbia Plateau, Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork.
4. (In support of strategy no. 4) Remove or explicitly control the timing and intensity of grazing to develop and promote the long-term persistence of shrub communities.

GROUP 24--SHARPTAIL SNAKE, CALIFORNIA MOUNTAIN KINGSLAKE, AND BLACK-CHINNED HUMMINGBIRD

Results

Species ranges, source habitats, and special habitat features--Group 24 consists of three species that primarily depend on open forest and woodland habitats: the black-chinned hummingbird, the sharptail snake, and the California mountain kingsnake. The range of the black-chinned hummingbird covers the entire basin except the high elevations of the Cascade Mountains in both the Northern and Southern Cascades ERUs and the high elevations of the northern Rocky Mountains (fig. 72). Both species of snakes occur in scattered, isolated populations along the eastern slope of the Cascade Range (fig. 72). The two species of snakes are only known to occur in the same location near the Columbia River Gorge.

Vol. 2, Figure 72--Ranges of species in group 24 within the basin.

These three species primarily group together based on their consistent use of interior ponderosa pine, and interior Douglas-fir vegetation types in all structural stages except stem-exclusion, closed-canopy forests. They also use mixed-conifer woodlands and Oregon white oak (vol. 3, appendix 1, table 1).

The black-chinned hummingbird is the only member of the group whose source habitats include juniper, juniper/sagebrush, chokecherry-serviceberry-rose, mountain mahogany, shrub wetlands, and old-forest aspen (vol. 3, appendix 1, table 1). The sharptail snake uses more source habitats than the kingsnake, including nearly all seral stages of cottonwood-willow (also used by the black-chinned hummingbird), nearly all structural stages of western red cedar-western hemlock, and the stem-exclusion, closed-canopy, and stand-initiation structural stages of western larch (vol. 3, appendix 1, table 1).

Logs and talus are special habitat features for both species of snakes because of their dependency on moist environments (vol. 3, appendix 1, table 2). In the absence of nearby streams, microhabitats with higher moisture are found under logs and within talus (Brown and others 1995). These features also provide protection from predators and habitat for potential prey. Additionally, deciduous tree riparian is also a special habitat feature for the sharp-tail snake (vol. 3, appendix 1, table 2).

Nectar-producing flowers are considered a special habitat feature for the black-chinned hummingbird because of the dependence on nectar as a primary food source.

Broad-scale changes in source habitats--Because the distribution of the two species of snakes is restricted to a few disjunct locations, the results of our analysis for this group are primarily based on source habitats for the black-chinned hummingbird, which is widely distributed throughout the basin both historically (fig. 73A) and currently (fig. 73B). Source habitats are most abundant in northeastern Washington, the Upper Klamath, and central Oregon (figs. 73A, B).

Overall, source habitats generally appeared to increase since the historical period, primarily in Oregon, Washington, and southeastern Idaho, whereas much of northern and central Idaho and Montana experienced declines (fig. 73C). About 53 percent of the watersheds basin-wide were projected to have increasing trends (fig. 74). The three ERUs with declining trends were Lower and Upper Clark Fork and Central Idaho Mountains (fig. 74), whereas mostly neutral trends were projected for the Blue Mountains and Northern Glaciated Mountains ERUs (fig. 74).

Vol. 2, Figure 73--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 24.

Vol. 2, Figure 74--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 24, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Changes in broad-scale habitat trends differed across the basin because of the wide array of cover types and structural stages used by group 24. Declining trends were fairly consistent for interior ponderosa pine old forest (both multi- and single-storied), and for stand-initiation stages of both ponderosa pine and Douglas-fir. Increases in habitat occurred in nearly all ERUs in both ponderosa pine and Douglas-fir young forests and in all woodland types (vol. 3, appendix 1, table 4). The increase in woodlands contributed substantially to the overall increase in source habitats, especially in rangeland-dominated ERUs (Upper Klamath, Northern Great Basin, Columbia Plateau, Snake Headwaters, and parts of the Blue Mountains). The increase in source habitats for group 24 closely reflects the increase in upland woodland reported for the basin (see map 3.58 in Hann and others 1997).

Condition of special habitat features--Trends in the condition of logs, talus, and flowers are not available at the broad scale. Activities that may negatively affect these variables include timber harvesting, road building, grazing, mining, and fire suppression. Timber harvesting and road building can lead to the direct removal of logs and flowers; mining can lead to disturbance of talus. Fire suppression can impact flower abundance by increasing forest canopy closure and reducing the amount of sunlight needed for flower development on herbaceous plants in the understory.

Other factors affecting the group--Humans have had a direct effect on snakes through collection, harassment, and accidental mortalities. Because of its striking coloration, the California mountain kingsnake is known to be in demand by collectors (ICBEMP 1996a). Humans also intentionally kill various snake species because of fear and hate, and are responsible for unintentional mortality caused by motorized vehicles (Brown and others 1995).

Population isolation was raised as a concern by the viability panel that evaluated sharptail snakes (ICBEMP 1996b). Although the viability panel did not evaluate the California mountain kingsnake, the same concerns and considerations are presumably important for this species because of its patchy and restricted range in the basin.

Because the black-chinned hummingbird is a Neotropical migrant, habitat used during migration and wintering habitat may have an impact on its populations. In a study on migrating rufous hummingbirds, researchers found a correlation between abundance of nectar-producing flowers and hummingbird survival in habitat used during migration (Russell and others 1994). A similar correlation likely exists with black-chinned hummingbirds. Little is known on the abundance or trends of the wintering habitat of the black-chinned hummingbirds.

Heavy grazing has had an overall negative impact on nectarivores by reducing the density of understory plants used as a food source (Saab and others 1995). Direct effects on the black-chinned hummingbird are unknown.

Population status and trends--There are no estimates of population change for either the sharptail snake or the California mountain snake within the basin. According to Brown and others (1995), however, loss of snake habitat and population declines in snakes worldwide have increased because of the increased paving of roads, fast cars, intensive agriculture, urban sprawl, desertification of arid lands, deforestation of the tropics, pesticides, hobby collecting, rattlesnake "roundups," and a general aversion to snakes. Sharp-tail snakes have declined in the Willamette Valley of Oregon, just west of the basin (Marshall and others 1996, Oregon Department of Fish and Wildlife 1987).

Population trend estimates for the black-chinned hummingbird in the basin are not available because of insufficient data from established BBS routes (Saab and Rich 1997). Specialized monitoring techniques would be needed to adequately measure population trends because they are difficult to detect (Saab and Rich 1997).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 24 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--Although the results of our analysis show an increase in source habitats across the basin, other sources of information have indicated that habitat and populations have decreased since the historical period. The trend of special habitat features for these species may affect populations more strongly than the broad-scale changes in source habitats. The following are issues that relate to special habitat features:

1. Loss of down logs and surface litter used by snakes as a result of timber harvest.
2. Loss of habitat connectivity for snakes as a result habitat loss and road construction primarily.
3. Decline in availability of understory flowering shrubs, particularly in riparian areas, because of cattle grazing.
4. Decreases in natural forest openings and shrub understories because of exclusion of fire and invasions by shade-tolerant trees.
5. Collection of California mountain kingsnakes.

Potential strategies--The issues identified above suggest the following broad-scale strategies to maintain the long-term persistence of sharptail snakes, California mountain kingsnakes, and black-chinned hummingbirds:

1. (To address issue no. 1) Survey and manage for downed logs and litter for the two species of snakes.
2. (To address issue no. 2) Seek opportunities to improve connectivity between isolated populations of both the sharptail and California mountain kingsnake.
3. (To address issue no. 3) Maintain and restore flowering herbs and shrubs in areas that have been

negatively affected by cattle grazing.

4. (To address issue no. 4) Restore fire as an ecological process, particularly in interior ponderosa pine and interior Douglas-fir plant communities, to encourage forest openings that are occupied by flowering shrubs and forbs.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Maintain and protect down logs at a level that is ecologically sustainable and meets the habitat requirements for snakes.
2. (In support of strategy no. 2) Close roads to minimize human disturbance and maximize dispersal capabilities, particularly in areas known to be occupied by either sharptail snakes or California mountain kingsnakes.
3. (In support of strategy no. 3) Remove or explicitly control the timing and intensity of grazing to develop and promote the long-term persistence of shrub communities.
4. (In support of strategy nos. 3 and 4) Accelerate development of flowering shrubs and forbs by the use of prescribed underburning and thinning, or allow for natural wildfires to occur, particularly in the Douglas-fir and ponderosa pine plant communities. Highest priorities for following these practices are in the Lower Clark Fork, Upper Clark Fork, and Central Idaho Mountains ERUs.

GROUP 25--NORTHERN GOSHAWK (WINTER)

Results

Species ranges and source habitats--Group 25 consists of winter habitat for the northern goshawk. Summer habitat for the northern goshawk is described in group 5. During winter, the range of the goshawk is basin-wide (fig. 75). Throughout North America, little is known about goshawks in winter, but indications are that northern goshawks are partial migrants. A portion of the population regularly winters outside the breeding area, whereas some do not migrate at all (Squires and Reynolds 1997). The degree to which goshawks migrate during winter may relate to prey availability. In the Yukon Territory in winter, goshawk numbers fluctuate with snowshoe hare numbers (Doyle and Smith 1994). Some goshawks may travel short distances in winter to lower elevations or more open habitats (Squires and Reynolds 1997), and migrations may consist of predominately immature birds (Sibley 1993).

Vol. 2, Figure 75--Ranges of species in group 25 within the basin.

Source habitats are found in old forest and unmanaged young forests in montane, lower montane, and riparian woodland community groups and chokecherry-serviceberry-rose (vol. 3, appendix 1, table 1). Also, contrary to summer source habitats, winter source habitats include all of the upland woodland types.

Important attributes of goshawk prey habitat include snags, downed logs, woody debris, large trees,

openings, herbaceous and shrubby understories, and an intermixture of various forest structural stages (Reynolds and others 1992).

Broad-scale changes in source habitats--Goshawk winter source habitats were projected to be broadly distributed, primarily throughout the forested areas of the basin in historical times (fig. 76A). Source habitats are still widely available, although more disjunct in many areas, and there has been an increase in habitats in some areas that provided little or no source habitats historically (fig. 76B).

Trends in source habitat availability differed geographically (fig. 76C). Most areas with strong negative trends were in the northeast portion of the basin, within the Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork ERUs, where habitat loss was generally greater than 90 percent (figs. 76C and 77; vol. 3, appendix 1, table 3). A preponderance of watersheds in the Northern Cascades, Blue Mountains, Snake Headwaters, and Central Idaho Mountains ERUs had moderate and strong negative trends (fig. 77). The most significant gains in source habitats occurred in the Upper Klamath and Northern Great Basin ERUs (fig. 77). About 50 percent of the watersheds in the Columbia Plateau, Owyhee Uplands, and Upper Snake ERUs also experienced strongly increasing trends (fig. 77). Trends in source habitats in the Southern Cascades showed a slight decrease (fig. 77).

Vol. 2, Figure 76--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 25.

Vol. 2, Figure 77--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 25, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--In areas with negative trends, projected declines occurred in nearly all source habitats, though predominately in the old-forest types (vol. 3, appendix 1, table 4). Some old-forest types increased in the Southern Cascades, Upper Klamath, and Blue Mountains ERUs. Further elaboration of the changes in old forest for the goshawk is found in the results for group 5, which includes goshawk (summer).

Large increases in juniper/sagebrush in the Upper Klamath, Northern Great Basin, Columbia Plateau, Blue Mountains, Upper Snake, and Snake Headwaters ERUs contributed to much of the increases in these ERUs or parts of these ERUs (fig. 77; vol. 3, appendix 1, table 4). Areas with increasing trends in source habitats correspond closely with the increases in upland woodlands as shown in map 3.58 in Hann and others (1997).

Other factors affecting the group--Little is known about population dynamics of goshawks, though it is thought that food availability may play an important role (Squires and Reynolds 1997). Goshawks prey primarily on relatively large-bodied mammals and birds, including tree squirrels, ground squirrels, lagomorphs, galliformes, corvids, piciforms, and passerines. Several studies have documented a positive relation of prey abundance with nest success (Doyle and Smith 1994, Linden and Wikman 1983, Ward

and Kennedy 1996). Important components of habitat for many of the prey species listed above are snags, downed logs, woody debris, openings, large trees, herbaceous and shrubby understories, and interspersions of different vegetation structural stages (Reynolds and others 1992). In many areas in the basin, fire suppression, timber harvesting, and livestock grazing have resulted in a decrease in many of the attributes listed above as important characteristics of prey habitat for goshawks (Hann and others 1997).

Some evidence indicates that diet composition may change drastically during the nonbreeding season in Sweden, but winter food habits are unknown in North American populations (Squires and Reynolds 1987, Widen 1987).

Effects of falconry, shooting, and trapping of goshawks in North America are thought to be minimal (Squires and Reynolds 1987).

Human disturbance at nest sites can cause failure, but there is no information on the effects of human activities during the nonbreeding or winter season (Anon. 1989, Boal and Mannan 1994, Speiser 1992, Squires and Reynolds 1987).

Population status and trend--The BBS data for the goshawk were insufficient to determine population trends for the basin (Saab and Rich 1997) or for any state or physiographic region within the basin (Sauer and others 1996), because of low detection of goshawks by using the BBS survey method. Sufficient data, however, were available for western North America to indicate a stable trend in numbers between the years 1966 and 1995 (Sauer and others 1996).

A separate trend estimate was derived from fall migration counts conducted by Hawkwatch International at four locations in Utah and New Mexico. These data indicated an average rate of decline in migrating goshawks of about 4 percent annually between 1977 and 1991 (Hoffman and others 1992). The extent to which the migration data represented local declines near the survey stations was not determined.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 25 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--Conservation issues for goshawk winter habitat, based on results of our analysis of source habitats in combination with empirical literature, include the following:

1. Reduction in the amount of old forests in the montane, lower montane, and riparian woodland community groups.
2. Possibly unsustainable conditions of old forests where there have been large transitions from shade-intolerant to shade-tolerant tree species. This issue stems from the exclusion of fire from many forested communities, which has resulted in increased susceptibility to stand-replacing fires (USDA Forest Service 1996).
3. Loss of important attributes of prey habitat, including large trees, snags, downed logs, forest openings, and herbaceous and shrubby understories because of fire suppression, timber harvesting, and livestock grazing.

Potential strategies--Potential strategies that would be effective for maintaining source habitats for

wintering goshawks within the basin are as follows:

1. (To address issue nos. 1 and 2) Especially in the northern areas of the basin, promote greater diversity in forest structure at the landscape scale. Mid-seral stages currently predominate and do not provide source habitats. Maintain stands with active goshawk nests in old-forest condition, and identify opportunities to increase the representation of old forests in individual watersheds.
2. (To address issue no. 2) Reduce the risk of loss of habitat by focusing old-forest retention and restoration efforts on areas with low probability of stand-replacing fires. In ERUS where old-forest habitat has remained stable or increased from historical conditions, efforts could be focused on retaining existing habitat in areas with lower fire and insect risk while managing other areas to reduce risks of catastrophic loss of habitat.
3. (To address issue nos. 1 and 3) Throughout the basin, provide for an abundant and sustainable prey base for goshawks by increasing the abundance of large trees, snags, downed logs, forest openings, and herbaceous and shrubby understories across the landscape.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) In the northern basin, identify representative stands of old forests for retention, and mid-successional stages for development into old-forest conditions. Priority should be given to large blocks having high interior-to-edge ratios and few large openings.
2. (In support of strategy nos. 1 and 3) Actively recruit snags and logs from green trees to increase the representation of old-forest structures (snags and logs) in mid-seral stands and in old forests where snags and logs are in low density or absent.
3. (In support of strategy no. 2) Thin small-diameter trees, either through hand equipment or prescribed burns, to reduce fuel loading and increase herbaceous and shrubby understories for prey habitat and improve growth of overstory trees.

GROUP 26--YUMA MYOTIS, LONG-EARED MYOTIS, FRINGED MYOTIS, AND LONG-LEGGED MYOTIS

Results

Species ranges, source habitats, and special habitat features--Group 26 is comprised of four species of bats: the Yuma myotis, long-eared myotis, fringed myotis, and long-legged myotis. All four species are year-round residents of the basin, active from spring through fall and hibernating during winter. The species in group 26 are similar in their use of a broad range of forest and woodland habitats for foraging.

The ranges of the long-legged myotis and long-eared myotis encompass the entire basin (fig. 78). The Yuma myotis occurs across most of the basin except for an area in the southeast portion (fig. 78). The fringed myotis occurs in the western half of the basin and in the Upper Clark Fork ERU (fig. 78).

Vol. 2, Figure 78--Ranges of species in group 26 within the basin.

Source habitats shared by all members of group 26 are all cover types in the montane, lower montane, riparian woodland, and upland woodland community groups, and the mountain hemlock cover type in the subalpine community group (vol. 3, appendix 1, table 1). The long-eared myotis ranges somewhat higher than the other species and uses whitebark pine, whitebark pine-alpine larch, and Engelmann spruce-subalpine fir as source habitats. Source habitats for the Yuma myotis and long-eared myotis extend into big sagebrush, mountain big sagebrush, and low sage cover types (vol. 3, appendix 1, table 1).

The long-eared and fringed myotis forage primarily by hover-gleaning insects off of foliage (Barclay 1991, Nagorsen and Brigham 1993, Perkins 1996). The long-eared myotis consumes moths, beetles, and other insects (Whitaker and others 1977, 1981), and the fringed myotis consumes mostly beetles (Black 1974, cited in O'Farrell and Studier 1980). Surveys based on bat vocalizations indicate that in forested habitats, foraging is highest in clearcuts and mature stands, and low in precommercially thinned and young, unthinned stands (Erickson and West 1996). The Yuma myotis is primarily found in association with rivers, lakes, ponds, and streams, where it forages over water and eats midges and emergent aquatic insects (Whitaker and others 1977).

Several special habitat features were identified for group 26 (vol. 3, appendix 1, table 2). Large-diameter (>53 cm [21 in]) snags with exfoliating bark provide maternity roosts for the long-legged myotis (Nagorsen and Brigham 1993, Ormsbee and McComb 1998, Rabe and others 1998), the fringed myotis (Chung-MacCoubrey 1996, Rabe and others 1998), and the long-eared myotis (Chung-MacCoubrey 1996, Rabe and others 1998). Caves, mines, and buildings provide maternity roosts for the fringed myotis, Yuma myotis, and long-eared myotis (Christy and West 1993, Nagorsen and Brigham 1993). Caves and mines also are used as hibernacula by all four species (Nagorsen and Brigham 1993). Various structures are used for day and night roosts, including exfoliating bark, rock crevices, mines, caves, and buildings (Manning and Knox-Jones 1989, Nagorsen and Brigham 1993, O'Farrell and Studier 1980). Ormsbee and McComb (1998) found that snags extending above the canopy were most frequently used by long-legged myotis for day roosts.

Rabe and others (1998) suggested that snag-roosting bats may require higher densities of snags than cavity-nesting birds, because the stage at which snags are suitable for bat roosts (exfoliating bark) is extremely short lived, requiring the use of several snags over the course of a lifetime of a bat. Bats frequently shift maternity roosts, possibly to find snags with better thermal conditions when the bark on the previous roost is no longer suitable (Rabe and others 1998).

The presence of water is considered a special habitat feature for the Yuma myotis because it forages mostly by flying low over water (permanent or seasonal) and feeding on emerging aquatic insects (Whitaker and others 1977). Although less dependent on water, long-legged myotis (Ormsbee and McComb 1998) and long-eared myotis (Ports and Bradley 1996) forage over or near water, and the fringed myotis frequently forages over thickets along streams (Nagorsen and Brigham 1993). In shrubland habitats, nearby riparian woodlands may provide the only available roost sites. Thus, all species in group 26 have a strong association with water and riparian vegetation.

Broad-scale changes in source habitats--When the need for suitable roost sites is ignored, there have been few changes in the extent of source habitats between historical and current periods (figs. 79A, B). Declining trends were most pronounced in the northern half of the Columbia Plateau and in the Upper Snake ERU, and increasing trends occurred mostly in the southern half of the Columbia Plateau, and in a few watersheds of the Northern Glaciated Mountains, Upper Klamath, Central Idaho Mountains, and Snake Headwaters ERUs (fig. 79C). Neutral trends in habitat extent were found in 59 percent of watersheds within the basin, and neutral trends predominated in all 13 ERUs (fig. 80). In most ERUs, the number of watersheds with increasing trends exceeded those with declining trends (fig. 80).

Vol. 2, Figure 79--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 26.

Vol. 2, Figure 80--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 26, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure associated with changes in source habitats---Neutral trends in habitat extent reflect the ability of species in group 26 to use a wide variety of cover types and nearly all structural stages of forests as source habitats. The basin has experienced dramatic declines in old-forest structural stages of all forest cover types (Hann and others 1997; vol. 3, appendix 1, table 4). However, for group 26, these losses have been offset by increases in mid-seral stages that also serve as source habitats, as long as suitable roost sites are available.

Declines in the northern portion of the Columbia Plateau, the southern portion of the Central Idaho Mountains, and portions of the Owyhee Uplands and Upper Snake ERUs are due to losses of big sagebrush and mountain big sagebrush to agriculture (Hann and others 1997). Increases in the Northern Glaciated Mountains are due primarily to areal increases in managed young forests of interior Douglas-fir and interior ponderosa pine (vol. 3, appendix 1, table 4). Increases in the Central Idaho Mountains are due primarily to areal increases in managed young forests and understory reinitiation stages of several forest cover types, including Engelmann spruce-subalpine fir, interior Douglas-fir, grand fir-white fir, lodgepole pine, and western larch (vol. 3, appendix 1, table 4).

Within the riparian woodlands community group, old forests had strongly declining trends throughout the basin (vol. 3, appendix 1, table 4) and generally remain only in stands smaller than the 1-km² (0.4-mi²) mapping unit used in this analysis. These losses occurred from changes in historical hydrologic regimes: reservoirs have eliminated many aspen and cottonwood-willow stands, a lowered water table has reduced others, and loss of periodic flooding has prevented establishment of seedlings (Merigliano 1996, Rood and Heinze-Milne 1989).

Condition of special habitat features---The number of caves has not changed significantly from historical to current times, but human disturbance from recreation has increased, causing some caves to be less available to hibernating bats. Mines proliferated in the early part of the historical period and provided additional habitat, but during the 1980s, thousands of abandoned mines throughout the West were closed with no input from biologists, thereby resulting in unknown loss of established roosts (Idaho State Conservation Effort 1995). The extent of cliffs and rocky areas has not changed since the historical period, but habitat quality of some cliffs has declined because of human disturbances (Lehmkuhl and others 1997).

Large-diameter snags >53 cm (21 in) have been reduced basin-wide in roaded areas with a history of

timber sales (Hann and others 1997; Hessburg and others 1999; Quigley and others 1996). Consequently, the neutral trends in source habitats for the long-legged myotis may give a more positive assessment of habitat availability than is actually the case.

In addition to riparian woodlands large enough to map at the broad scale, smaller patches of riparian vegetation have declined in extent basin-wide, because of disruption of hydrologic regimes from dams, water diversions, and road construction, along with grazing and trampling of riparian vegetation by livestock and increased recreational use along stream courses (USDA Forest Service 1996). These fine-scale changes have caused additional declines in bat foraging habitat and potential roost sites.

Other factors affecting the group--Roost availability has greatly influenced the distribution of all Nearctic bat species (Humphrey 1975), and the conservation of group 26 bats is largely dependent on maintaining suitable roost sites. The most straightforward source of impact is destruction of the structure, that is, loss of snags through timber harvests, and removal of old buildings and bridges or closure of mines and caves for safety reasons (Perlmeter 1995, Pierson and others 1991). Perkins and Peterson (1997) attributed the low detection of bats in the Owyhee Mountains to the lack of suitable roosts, particularly in the form of cottonwood and juniper snags.

The second source of impact is disturbance of roosting bats, primarily by recreational activities in or near caves but also from mining, road construction, road access and any other activities near roosts (Pierson and others 1991). During winter, rising out of torpor requires a large caloric output, and repeated disturbances can drain the energy reserves of a bat and lead to starvation (Nagorsen and Brigham 1993). Recreational use of caves during the hibernation and nursery periods seriously affects persistence of individual colonies if disturbances are frequent (Nagorsen and Brigham 1993).

The third source of impacts at roost sites is purposeful killing of bats. Because of their high visibility at colonial roosts, bats have suffered high mortality rates; total loss of colonies have occurred from shooting by individuals who often are guided by negative folklore regarding bats (Nagorsen and Brigham 1993). Destruction of a single colony may represent a significant impact across large areas because of the patchy distribution of bats related to roost availability.

Roads may indirectly affect bat species by increasing human access to roost sites. Caves have become more accessible, increasing the amount of human visitation and potential harassment of bats. The presence of roads increases the likelihood that snags will be cut for fuel wood (see Hann and others 1997). The additional loss of snags in areas where snag densities are currently low could limit populations of group 26 species.

Direct contact with pesticides can cause illness or death in bats. Although most organochlorine pesticides that cause accumulation of chemicals up the food chain have been banned or highly restricted in the United States, the relatively short-lived organophosphates can provide high risks during application (Clark 1988). For example, a large die-off of bats was observed in Arizona after the application of methyl parathion, and was believed to be linked to direct contact with this chemical (Clark 1988).

Population status and trends--There are insufficient population data on any species in group 26 to determine population trends. In general, however, bats in the basin are believed to be declining because of increased human disturbance of roosts, declining snag densities, decrease of late-seral lower montane and montane forests, decreased acreage and quality of riparian areas, pesticide use, changes in habitats on non-Federal lands, direct killing, and decreases in water quality (Lehmkuhl and others 1997).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 26 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--Our results, combined with literature and other empirical information, suggest that the following issues are important for group 26:

1. Basin-wide loss of large-diameter snags (>53 cm [21 in]) for the long-legged myotis maternity roosts and day roosts.
2. Destruction of roosts, disturbance of roosting bats, or both.
3. Degradation and loss of native riparian vegetation.
4. Impacts of pesticides on bats and their prey.
5. Lack of information on hibernacula, including locations, special features, and numbers of bats associated with them.
6. Lack of population trend data.

Potential strategies--The following strategies could be used to maintain and improve habitat for these bat species:

1. (To address issue no. 1) Actively manage for the retention and recruitment of large-diameter snags in all forest cover types and structural stages.
2. (To address issue no. 2) Protect all roosts and reduce human disturbances near roosts.
3. (To address issue no. 3) Maintain and improve the condition of riparian and wetland vegetation for bat foraging areas.
4. (To address issue no. 4) Alleviate impacts of pesticides on bat populations.
5. (To address issue nos. 5 and 6) In cooperation with other state, Federal, and tribal agencies, establish a coordinated approach to search for hibernacula, and to protect these sites.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Retain existing snags, particularly if >53 cm (21 in) and provide measures for snag replacement. Review existing snag guidelines or develop guidelines that reflect local ecological conditions and address snag numbers, diameter, height, decay class, species, and distribution. Retain snags in clusters to provide adjacent roosts for maternity colonies. Maintain snags at higher than historical levels, to restore loss in previously harvested areas (ICBEMP 1996d).
2. (In support of strategy no. 1) Emphasize retention of snags that provide best solar exposure to bark or cavity roost sites (Betts 1996).
3. (In support of strategy no. 1) Reduce road densities in managed forests where snags are currently in low abundance. Close roads after timber harvests and other management activities, and minimize the period when such roads are open to minimize removal of snags along roads. In addition or as an

- alternative to road management, actively enforce fuel wood regulations to minimize removal of snags.
4. (In support of strategy no. 1) Restrict fuel wood permits to disallow snag cutting where snags are in low abundance, and particularly where existing roads cannot be closed. Blair and others (1995) recommend that public fuel wood harvest should be limited to trees <38 cm (15 in) d.b.h.
 5. (In support of strategy no. 2) Monitor known roosts for potential human disturbances, and initiate closures of recreational or construction activity near roost sites.
 6. (In support of strategy no. 2) If possible, stabilize old structures that are important for maternity roosts and hibernacula.
 7. (In support of strategy no. 2) Survey caves, mines, and abandoned buildings before removal or closure, and protect roosting bats from human presence and disturbance. During closures, use specialized gates designed to allow continued use of mines and caves by bats (Pierson and others 1991).
 8. (In support of strategy no. 2) Assure that construction of roads and rights-of-way are not going to cause siltation, slumping, or water run-off to enter cave habitats or alter other roosting structures (Perkins 1992-1994).
 9. (In support of strategy no. 3) Identify areas of existing riparian and wetland habitats that are important bat foraging areas, and design conservation measures to protect and enhance foraging opportunities for bats.
 10. (In support of strategy no. 3) Modify grazing practices to improve condition of degraded riparian areas for bat foraging and roosting.
 11. (In support of strategy no. 3) Restore degraded areas by appropriate mechanical treatments and with seedings of appropriate native species.
 12. (In support of strategy no. 4) Avoid pesticide use in areas of high bat foraging activity or near nursery colonies.
 13. (In support of strategy no. 5) Use existing interagency cooperative agreements, or develop agreements where needed to conduct surveys for hibernacula.
 14. (In support of strategy no. 5) Use individual project planning (such as timber sales, road construction, mineral extraction, or recreational development) as opportunities for conducting surveys for new roost sites and to assess population status of known roosts.

GROUP 27--PINE SISKIN AND TOWNSEND'S BIG-EARED BAT

Results

Species ranges, source habitats, and special habitat features--Group 27 includes the pine siskin and the Townsend's big-eared bat, both of which are year-round residents of the basin. The pine siskin occurs throughout the basin except for low elevation, nonforested areas, and the Townsend's big-eared bat is found basin-wide (fig. 81).

Vol. 2, Figure 81--Ranges of species in group 27 within the basin.

Both species are forest generalists within the subalpine, montane, upland woodland, and riparian woodland community groups. Most cover types within these community groups are source habitats for both species, but Engelmann spruce-subalpine fir is considered source habitat for only the pine siskin, whereas aspen is used only by the big-eared bat. Source habitat for both species was considered to be in all structural stages except the stem exclusion and stand initiation stages (vol. 3, appendix 1, table 1). Source habitats for the big-eared bat also include several cover types within the upland shrubland, upland herbland, and riparian shrubland community groups (vol. 3, appendix 1, table 1).

No special habitat features were identified for the pine siskin. Breeding takes place in various conifer species, including ornamentals, and foraging occurs in trees, shrubs, and grassy areas (Dawson 1997). Diet consists primarily of small seeds from annual plants, conifers, and deciduous trees (Dawson 1997). Pine siskin populations are highly irruptive on a continental scale, causing local abundance or scarcity of siskins from one year to the next, apparently in response to food availability (Bock and Lepthien 1976, Dawson 1997).

The Townsend's big-eared bat is colonial in its use of caves and cavelike structures for nursery colonies, day roosts, and hibernacula (Idaho State Conservation Effort 1995, Nagorsen and Brigham 1993; vol. 3, appendix 1, table 2). Big-eared bats do not roost in crevices like many other bat species but rather restrict their roosting sites to the ceilings of cavelike structures (caves, mines, and buildings), where they aggregate in large colonies. A stable, cold temperature and moderate airflow may be important criteria for hibernation (Genter 1986, Humphrey and Kunz 1976). The distribution of big-eared bats is patchy across the basin because of their restrictive roosting requirements.

The big-eared bat is a moth specialist (Idaho State Conservation Effort 1995; Nagorsen and Brigham 1993; Whitaker and others 1977, 1981). In central Oregon, they forage in sagebrush, bitterbrush, and open ponderosa pine forests (Dobkin and others 1995).

Broad-scale changes in source habitats--Source habitats were widespread across the basin historically, with greatest concentrations in the mountains of the Northern Cascades, Southern Cascades, Upper Klamath, Blue Mountains, Northern Glaciated Mountains, Upper Snake, and Snake Headwaters ERUs (fig. 82A). Extensive shrubland and grassland habitats suitable only for the big-eared bat occurred in the Columbia Plateau, Northern Great Basin, and Owyhee Uplands. The current extent of habitat is similar to the historical distribution (fig. 82B), although the abundance of habitat has changed in some areas. Watersheds with declining trends were primarily in the northern half of the Columbia Plateau, the Upper Snake, and Snake Headwaters ERUs (figs. 82C and 83). Watersheds with increasing trends were mostly in the Upper Klamath, Blue Mountains, Northern Glaciated Mountains, and Central Idaho Mountains (figs. 82C and 83). Basin-wide, the number of watersheds with declining, increasing, or static trends was nearly equal, representing 34, 34, and 31 percent of watersheds, respectively (fig. 83).

Vol. 2, Figure 82--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 27.

Vol. 2, Figure 83--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 27, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Mixed trends in habitat extent reflect the association of both species in group 27 with several cover types and nearly all structural stages of forests as source habitats. The basin has experienced dramatic declines in old-forest structural stages of all forest cover types (Hann and others 1997; vol. 3, appendix 1, table 4), but for group 27, these losses have been offset by increases in mid-seral stages that also serve as source habitats. Increases in the areal extent of habitats in the Upper Klamath were due to transitions from the fescue-bunchgrass cover type to mixed-conifer woodlands and an areal increase in the extent of interior Douglas-fir, historically less than 2 percent, but currently 15 percent of the ERU (Hann and others 1997). In the Blue Mountains, Northern Glaciated Mountains, and Central Idaho Mountains, increasing trends were largely due to increases in the areal extent of grand fir-white fir. Engelmann spruce-subalpine fir increased in the Central Idaho Mountains as well (Hann and others 1997; vol. 3, appendix 1, table 4).

Static trends in nonforested habitats are partially due to transitions from big sagebrush to juniper/sagebrush and juniper woodlands (Hann and others 1997), which have resulted in no net change in source habitats for the big-eared bat. Declines have occurred in the northern portion of the Columbia Plateau because of transitions from big sagebrush to agriculture (Hann and others 1997).

Condition of special habitat features--The number of caves likely has stayed the same from historical to present periods, but human disturbance from recreation has increased, causing some caves to be abandoned by big-eared bats (Idaho State Conservation Effort 1995). Mines proliferated in the early part of the historical period and provided additional habitat, but during the 1980s, thousands of abandoned mines throughout the West were closed with no input from biologists, thereby resulting in unknown loss of established roosts (Idaho State Conservation Effort 1995).

Other factors affecting the group--Pine siskin foraging behavior, geographic location, and population levels are highly influenced by the combination of current population level and food availability--an abundance of seeds will cause the population to expand, and if the next year's crop is unable to support the expanded population, the birds will move elsewhere (Bock and Lepthien 1976).

Because the distribution of Townsend's big-eared bats is dependent on specialized roosting requirements, alterations and disturbances of any structures used for day roosts, nursery colonies, or hibernacula (caves, mines, old buildings) could affect the persistence of individual colonies. The most straightforward source of impact is destruction of the structure, that is, removal of old buildings or closure of mines and caves for safety reasons (Pierson and others 1991).

The second source of impact is disturbance of roosting bats, primarily by recreational activities in or near caves but also from mining, road construction, and any other activities near roosts (Idaho State Conservation Effort 1995). Females at nursery colonies are alert and readily take flight if disturbed (Perkins and Schommer 1992), and frequent interruptions are known to result in abandonment of the roost site (Idaho State Conservation Effort 1995, Nagorsen and Brigham 1993). During winter, rising out

of torpor requires a large caloric output, and repeated disturbances can drain the energy reserves of a bat and lead to starvation (Nagorsen and Brigham 1993). Recreational use of caves during the hibernation and nursery periods seriously affects persistence of individual colonies if disturbances are frequent (Idaho State Conservation Effort 1995, Nagorsen and Brigham 1993).

The third source of impacts at roost sites is purposeful killing of roosting bats (Idaho State Conservation Effort 1995). Because of their high visibility at colonial roosts, big-eared bats have suffered high mortality rates and sometimes total loss of a colony from shooting by individuals who often are guided by negative folklore (Nagorsen and Brigham 1993). Destruction of a single colony may represent a significant impact on big-eared bats across large areas because of the patchy distribution of bats related to roost availability.

The big-eared bat is negatively affected by the presence of roads. Increased road networks have made caves more accessible and have increased the amount of human visitation and potential harassment.

Because the big-eared bat is insectivorous, use of insecticides in foraging areas has the potential to impact bat species, primarily by reducing the prey base. For example, forest spraying for tussock and spruce budworm moths, although targeted at the larval stage of these insects, ultimately affects the number of flying adults and can cause a sufficient reduction in the prey base to suppress a year or two of Townsend's bat reproduction (Perkins and Schommer 1992). Also, exposure to insecticides can directly affect the health of bats. Although most organochlorine pesticides that cause accumulation of chemicals up the food chain have been banned in the United States or their use highly restricted, the relatively short-lived organophosphates can cause illness or death to bats during application (Clark 1988).

Population status and trends--Population trends for the pine siskin are difficult to obtain because the irruptive tendencies of this species result in highly variable annual numbers at any given locale (Dawson 1997). The BBS data show no significant population trends in most states, Canadian provinces, or BBS physiographic regions because of wide fluctuations in numbers or insufficient routes to determine a trend (Sauer and others 1996). Two areas with significant annual declines from 1966 to 1995, however, have been reported, which reflect possible population trends in the basin: an annual decline of 4.5 percent ($\bar{n} = 52$, $\bar{P} < 0.01$) has occurred on BBS routes in Washington, and an annual decline of 4.1 percent ($\bar{n} = 196$, $\bar{P} < 0.01$) has occurred in USDI Fish and Wildlife Service Region 1 (five Western states) (Sauer and others 1996).

Wintering populations of the big-eared bat seem to have declined, based on a comparison of counts made at hibernacula in central Oregon in the 1960s compared to the 1980s (Perkins 1987). In general, several species of bats in the basin have declined because of increased human disturbance of roosts, declining snag densities, decrease of late-seral lower montane and montane forests, decreased acreage and quality of riparian areas, pesticide use, changes in habitats on non-Federal lands, direct killing, and decreases in water quality (Lehmkuhl and others 1997).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 27 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--Our results, combined with literature and other empirical information, suggest that the following issues are important for group 27:

1. Unknown causes for population declines of pine siskins.

2. Direct loss of big-eared bat roosts because of cave and mine closures and destruction of abandoned buildings.
3. Excessive disturbance of roosting bats because of human activities.
4. High mortality of roosting bats or total loss of colonies because of vandalism and shooting.
5. Reduction in bat prey base (moths) through excessive use of insecticides.

Potential strategies--strategies for reversing the declining trends in pine siskin populations are difficult to formulate because of the irruptive nature of siskin populations at the continental scale. The following strategies have been identified to reverse broad-scale declines in populations of the big-eared bat:

1. (To address issue no. 2) Protect all known roost sites (nursery, day roosts, and hibernacula) of big-eared bats and restore historical roosts where feasible.
2. (To address issue no. 3) Reduce levels of human activities around known bat roosts.
3. (To address issue no. 4) Reduce vandal-related mortalities of roosting bats
4. (To address issue no. 5) Reduce impacts of insecticide use on principal prey of big-eared bats.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Survey all mines and caves scheduled for public closure for big-eared bats before closure. If roosting colonies are found, or if the structure has potential as a roosting colony, carry out the closure with gates that allow bats to enter and exit the structure. Unless superseded by other designs, use the bat gate designs in Tuttle and Taylor (1994), presented in appendix B of Idaho's Conservation strategy for Townsend's big-eared bat (Idaho State Conservation Effort 1995). If possible, stabilize old structures that are important for maternity and hibernacula sites (Perkins 1992-1994).
2. (In support of strategy no. 2) Initiate seasonal public closures of caves used as big-eared bat roosts during critical time periods, by using signs, road closures, and bat gates.
3. (In support of strategy no. 2) Reduce surveys to the minimum needed for assessing colony health and population status. Coordinate research efforts to minimize entry of roosts for data collection.
4. (In support of strategy no. 3) Increase public education and awareness of bat ecology and the current conservation status of big-eared bats.
5. (In support of strategy nos. 2 and 3) Reduce human access to bat roosting structures by closing roads that facilitate access to such habitat.
6. (In support of strategy no. 4) Avoid or minimize application of pesticides near bat roosts (Perkins 1992-1994). Utilize a 3.2-km (2-mi) "no-spray" buffer zone around roost sites (Idaho State Conservation Effort 1995). Within a 16-km (10-mi) radius of known roosts, use a strip-spraying technique to reduce the amount of area sprayed.

GROUP 28--SPOTTED BAT, PALLID BAT, AND WESTERN SMALL-FOOTED MYOTIS

Results

Species ranges, source habitats, and special habitat features--Group 28 consists of three bat species that generally are associated with low-elevation woodlands and shrublands: the spotted bat, pallid bat, and western small-footed myotis. The spotted bat and pallid bat occur in low numbers throughout eastern Washington and Oregon, and the spotted bat also occurs in eastern and southern Idaho (fig. 84). The small-footed myotis is somewhat more abundant and occurs throughout the basin except for high-elevation sites in the Cascade Range (fig. 84).

Vol. 2, Figure 84--Ranges of species in group 28 within the basin.

This analysis addresses year-round source habitat for all three species. The small-footed myotis is known to hibernate in the basin, but it is not known whether the spotted bat and pallid bat hibernate or leave the basin during winter (Nagorsen and Brigham 1993). With no migratory information, we have assumed that source habitats for all three species include winter hibernacula.

Cover types used as source habitats by all species in group 28 include interior ponderosa pine, juniper woodland, juniper/sagebrush, big sagebrush, mountain big sagebrush, and low sage (vol. 3, appendix 1, table 1). Additional cover types used as source habitats by one or two group members include cottonwood-willow (small-footed myotis), interior Douglas-fir and shrub wetlands (spotted bat), and salt desert shrub (spotted and pallid bats). Within interior ponderosa pine, the pallid bat is limited to old-forest structural stages, whereas the spotted bat and small-footed myotis also use young forest and understory reinitiation stages (vol. 3, appendix 1, table 1). All three species use both open- and closed-canopy structures of the shrub cover types.

A special habitat feature associated with all source habitats is the presence of cliffs or other rocky areas for roost sites (vol. 3, appendix 1, table 2). For the spotted and pallid bats, it is not necessary for roost structures to be adjacent to foraging areas because the spotted bat is known to travel up to 10 km (6.2 mi) between day roosts and feeding areas (Wai-Ping and Fenton 1989), and the pallid bat commutes up to 4 km (2.5 mi) (Nagorsen and Brigham 1993). Distances farther than these, however, would render shrub habitats unsuitable as source foraging areas. Commuting distances have not been reported for the small-footed myotis, but it seems to be versatile in its selection of roost sites, using boulders, vertical banks, and talus slopes in addition to cliffs (Nagorsen and Brigham 1993). Within this group, the spotted bat appears most limited in roost site selection, with all roosts reported in crevices of high cliffs (Nagorsen and Brigham 1993, Sarell and McGuinness 1993, Wai-Ping and Fenton 1989). The pallid bat primarily roosts in rock crevices but also uses tree cavities, buildings, and mines (Nagorsen and Brigham 1993).

The small-footed myotis and spotted bat are both aerial feeders, with diets that differ according to local prey availability (Nagorsen and Brigham 1993). In eastern Oregon, the small-footed myotis was reported to consume primarily moths, true bugs, and flies (Whitaker and others 1981). In eastern British Columbia, the spotted bat consumed mostly moths (Wai-Ping and Fenton 1989). The pallid bat can aerial feed, but mostly gleans prey from vegetation and the ground. In eastern Oregon, the diet was grasshoppers and moths (Whitaker and others 1981).

Broad-scale changes in source habitats--Historically, source habitats for group 28 were concentrated in the Columbia Plateau, Northern Great Basin, Owyhee Uplands, and Upper Snake ERUs, and patchily distributed elsewhere in the basin (fig. 85A). The current distribution of habitats resembles the historical extent, but there have been significant losses of habitat in the Columbia Plateau and total loss of the former patchy habitats in the Upper Clark Fork ERU (fig. 85B). Trends in habitat extent were variable across the basin, but in general, habitats declined in the northern portion of the basin and were static to increasing in the south, except for the Snake Headwaters, a southern ERU with declining trends (fig. 85C).

About one-third of the watersheds within the basin had static trends in the areal extent of source habitats, but nearly half had declining or strongly declining trends (fig. 86). Eighty percent of watersheds in the Lower Clark Fork and 54 percent of watersheds in the Columbia Plateau had declining and strongly declining trends (fig. 86). Increasing and strongly increasing trends were projected in 43 percent of the watersheds in the Southern Cascades and 50 percent of the watersheds in the Upper Klamath (fig. 86). These represent the two ERUs with the highest percentages of increasing habitat extent for group 28.

Vol. 2, Figure 85--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 28.

Vol. 2, Figure 86--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 28, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Throughout the basin, declines in source habitats of shrubland bats were associated with declines in big sagebrush, mountain big sagebrush, and old-forest structural stages of interior ponderosa pine and interior Douglas-fir (vol. 3, appendix 1, table 4). Source habitats declined in the Columbia Plateau and Snake Headwaters because of the conversion of 46 and 41 percent of the big sagebrush cover type to agriculture within each ERU, respectively (Hann and others 1997). In the Lower Clark Fork ERU, 66 percent of the interior ponderosa pine cover type was replaced by grand fir-white fir (Hann and others 1997), a cover type that does not serve as source habitat for group 28.

Increases in habitat extent generally were due to increases in juniper woodlands and juniper/sagebrush cover types (vol. 3, appendix 1, table 4). These increases often occurred in ERUs that experienced declines in native shrublands, resulting in overall mixed trends; for example, in the Owyhee Uplands (vol. 3, appendix 1, table 4).

Condition of special habitat features--The extent of cliffs and rocky areas in the basin has not changed since the historical period, but the habitat quality of some cliffs has declined because of human disturbances (Lehmkuhl and others 1997).

Other factors affecting the group--Human disturbance can affect bat nursery colonies by disrupting young

during the critical periods of growth and development. For spotted and pallid bats, nursery colonies are often inaccessible, and therefore disturbance potentials are low. The exception could occur if one or more rock climbing routes passed through a nursery colony and were visited frequently by climbers. Currently, no situation of this kind has been identified in the basin, but this may be due to a lack of monitoring rather than an absence of nursery colony-climber interactions.

Human activities can result in habitat degradation or disturbance at day roosts. Examples include road construction, dam building, mineral extraction, and the stabilizing of hazardous falling rocks above developments (Sarell and McGuinness 1993).

Direct contact with pesticides can cause illness or death in bats. Although most organochlorine pesticides that cause accumulation of chemicals up the food chain have been banned in the United States or their use highly restricted, the relatively short-lived organophosphates can provide high risks during application (Clark 1988). For example, a large die-off of bats was observed in Arizona after the application of methyl parathion, and was believed to be linked to direct contact with this chemical (Clark 1988).

Pesticides also can impact bat populations by reducing the availability of arthropods that serve as prey. Bats in group 28 are impacted by the spraying of forests and agricultural crops for insect pests.

Population status and trends--Population estimates for bat species in the basin are either unknown or very local in scale. Lehmkuhl and others (1997), however, reported that habitat conditions for most bat species have declined significantly from historical conditions because of the conversion of native vegetation to agriculture and urban, increased human disturbance of roosts, reduced large snag densities, decreased acreage and distribution of late-seral montane and lower montane forests, and reduced acreage and quality of riparian areas.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 28 with broader, ecosystem-based objectives for all other resources.

Issues--Our results and the conclusions drawn from published literature suggest the following issues are important for group 28:

1. Loss of native shrub vegetation.
2. Disturbances at nursery and day roosts.
3. Impacts of pesticides on bats and their prey.
4. Lack of information on hibernacula, including locations, special features, and numbers of bats associated with them.
5. Lack of population trend data.

Potential strategies--The following strategies could be used to maintain and improve habitat for these bat species:

1. (To address issue no. 1) Maintain and improve the condition of native shrublands to provide foraging

areas.

2. (To address issue no. 2) Reduce human disturbances near known roosts.
3. (To address issue no. 3) Alleviate impacts of pesticides on bat populations.
4. (To address issue nos. 4 and 5) In cooperation with other state, Federal, and tribal agencies, establish a coordinated approach to search for hibernacula.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Identify areas of existing native shrubland that could be managed for long-term persistence of native shrub cover types.
2. (In support of strategy no. 1) Explore options under the CRP (Johnson and Igl 1995), or develop other incentive programs to encourage restoration of agricultural areas to native cover types. Focus on areas that would increase patch size or links with existing source habitat patches.
3. (In support of strategy no. 1) Restore degraded areas by appropriate mechanical treatments and with seedings of native shrub, grass, and forb species.
4. (In support of strategy no. 2) Monitor known nursery roosts for potential disturbances, and initiate seasonal closures of recreational activity where appropriate. For example, seasonal restrictions on rock climbing would be appropriate if climbing routes passed through spotted bat nursery colonies.
5. (In support of strategy no. 2) Provide access for bats when mines are permanently closed.
6. (In support of strategy no. 2) Conduct surveys for bat roosts and hibernacula before road construction, mineral extraction, or slope stabilization where such activities are scheduled to occur near cliffs or caves with potential roosts. Provide mitigation or seasonal restrictions of potentially disturbing activities within the appropriate planning documents.
7. (In support of strategy no. 3) Avoid pesticide use in areas of high bat foraging activity or near nursery colonies.
8. (In support of strategies no. 4) Use existing interagency cooperative agreements, or develop agreements where needed to conduct surveys for hibernacula.

GROUP 29--WESTERN BLUEBIRD

Results

Species ranges, source habitats, and special habitat features--Group 29 consists of migratory breeding habitat for western bluebirds. Within the basin, western bluebirds are distributed across eastern Oregon and Washington, northern and western Idaho, and northwestern Montana (fig. 87). They are present in all ERUs except the Upper Snake and Snake Headwaters.

Vol. 2, Figure 87--Ranges of species in group 29 within the basin.

Western bluebirds use open forest stands and woodlands in combination with shrub and grass habitats. Specific source habitats (vol. 3, appendix 1, table 1) include old forest, single-storied western white pine and ponderosa pine; old forest aspen; stand-initiation stages of most montane forest and lower montane forest community groups; juniper and white oak woodlands; the open-canopy low-medium shrub stage of most of the upland shrub community type; and native bunchgrasses and forbs. Additionally, burned pine forests created by stand-replacing fires, likely are source habitats (Saab and Dudley 1998). Burned habitats, however, were not identified for this analysis.

Juxtaposition of forested and open areas is a necessary characteristic of source habitats for western bluebirds because they typically nest in tree cavities and forage for insects in adjacent openings (DeGraaf and others 1991; vol. 3, appendix 1, table 2). Because juxtaposition of cover types is important for nesting western bluebirds, they are considered a "contrast" species, and a finer scale analysis is needed to fully evaluate the status of their source habitats.

Western bluebirds are secondary cavity-nesters, so snags are a special habitat feature (vol. 3, appendix 1, table 2). They will use old woodpeckers holes, natural cavities, and nest boxes (Brawn and Balda 1988, DeGraaf and others 1991). Their nests are located in open forests or at forest edges. In burned ponderosa pine forests of western Idaho, nesting western bluebirds favored partially salvage-logged compared to unlogged stands (0.44 nests per km surveyed [0.71 nests per mi] in logged vs. 0.16 nests per km [0.26 nests per mi] in unlogged) (Saab and Dudley 1998). Openings in partially logged, burned forests likely provided greater opportunities for aerial foraging by the bluebirds. In salvaged units, snag [>23 cm (9 in) d.b.h.] densities at bluebird nest sites ($n = 65$) averaged 65 ± 5.9 snags per ha (26.3 ± 2.4 snags per acre), and at nonnest random sites ($n = 180$) 31.4 ± 1.9 snags per ha (12.7 ± 0.8 snags/acre). Average diameter of nest trees in the burned forests of western Idaho was 34.8 ± 1.5 cm (13.7 ± 0.6 in).

Broad-scale changes in source habitats--Source habitats for western bluebirds declined strongly throughout most of the basin. Throughout the basin, source habitats for western bluebird had declined strongly in 50 percent of watersheds and moderately in another 25 percent of watersheds (figs. 88 and 89). The apparent strong negative trends were in seven ERUs: the Northern Cascades, Southern Cascades, Columbia Plateau, Blue Mountains, Northern Glaciated Mountains, Lower Clark Fork, and Upper Clark Fork (fig. 89). More moderate declining trends were projected for the Upper Klamath and Central Idaho Mountains (fig. 89), whereas there was little change in source habitats from historical to current in the Northern Great Basin and Owyhee Uplands (fig. 89).

Vol. 2, Figure 88--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 29.

Vol. 2, Figure 89--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 29, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Hann and others (1997, see table 3.139) reported ecologically significant basin-wide declines for four of the terrestrial communities that support components of western bluebird source habitats. Communities that declined significantly were early-seral lower montane forest, late-seral lower montane single-layer forest, upland shrublands, and upland herblands. Of the terrestrial communities providing source habitats for bluebirds, only upland woodlands showed a basin-wide significant increase from historical to current (table 3.139 in Hann and others 1997). Decreases in habitats important to western bluebirds were also significant at the level of individual ERUs. The upland herb community declined significantly in all 11 ERUs within the range of the western bluebird, early-seral lower montane forest and late-seral lower montane single-layer forest declined in 10 ERUs, upland shrub declined in 8 ERUs, and early-seral montane forest declined in 6 ERUs (tables 3.141 through 3.165 in Hann and others 1997). Late-seral single-layer montane forest declined in four ERUs while increasing in five ERUs, and upland woodlands declined in three ERUs while increasing in six ERUs. Our evaluation at the broad scale did not assess the distribution of foraging habitat in relation to that for nesting habitat. Additional analysis of the juxtaposition of foraging with nesting habitats is need at a finer scale of resolution. Results for source habitats shown here for both the current and historical time periods are likely overestimates as they do not take into account the need for juxtaposition of habitats.

Condition of special habitat features--Densities of large-diameter snags (>53 cm [21 in] d.b.h.) have declined basin-wide from historical to current levels (Hann and others 1997; Hessburg and others 1999; Quigley and others 1996). Trends in densities of smaller snags are variable (Hann and others 1997).

The scale of the analysis does not allow determination of change in the amount of edge or amount of edge habitat. Thus, this special habitat feature was not evaluated for changes in source habitats presented in the above results. Some levels of decrease in total habitat area may be associated with increases in edge habitat. Consequently, the large decreases reported here for western bluebird habitat may be somewhat mitigated by increases in edge as habitat blocks are harvested.

Other factors affecting the group--Some western bluebirds that breed in the basin migrate to California and Baja California in winter (DeGraaf and others 1991). Conditions on these wintering grounds could affect the status of populations in the basin. Western bluebirds respond positively to artificially constructed nest boxes in areas where the availability of cavities is limiting. In one study (Brawn and Balda 1988), bluebird densities increased from 8 to 31 pairs per 40 ha (100 acres) after the construction of nest boxes. Usurpation of nest cavities by Lewis' woodpeckers (Saab and Dudley 1995) could have negative effects on western bluebirds. Stress and elevated energetic costs could be associated with territorial encounters with Lewis' woodpeckers and potentially reduce reproductive success of western bluebirds.

Population status and trends--Saab and Rich (1997) reported that western bluebird populations in the basin were stable over the period 1968-94 based on BBS data. Stable population trends also have been reported for this western species throughout its range for the period 1966-96 (Sauer and others 1996). Specialized monitoring techniques may be needed for better estimates of bluebird population trends

(Saab and Rich 1997).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 29 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--Primary issues affecting source habitats of western bluebirds are as follows:

1. Reductions in snag densities.
2. Reductions in early- and late-seral montane and lower montane forests.
3. Possibly unsustainable conditions in late-seral montane and lower montane forests where there have been large transitions from shade-intolerant to shade-tolerant species.
4. Reductions and degradation of upland shrublands and herblands.

Potential strategies--Habitat for western bluebirds could be improved by implementing the following strategies:

1. (To address issue no. 1) Maintain large remnant trees and snags in all seral stages of montane and lower montane forests.
2. (To address issue no. 2) Maintain and restore early-seral and late-seral montane and lower montane forests where those types have been reduced in extent. Both the extent and pattern of these habitats are of concern because source habitats for western bluebirds are found in edge areas. Where possible, retention efforts for late-seral forests should be focused on areas where the potential for stand-replacing fires is low (USDA Forest Service 1996).
3. (To address issue no. 3) Restore fire regimes that maintain a natural mosaic of shrublands and forests in those ERUs and portions of ERUs where substantial habitat remains (for example, Northern Great Basin, Owyhee Uplands, southern portion of Columbia Plateau). In some areas, such strategies will result in temporary declines and periodic fluctuations in habitat abundance.
4. (To address issue no. 4) Restore upland shrub and herblands.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

- 1 (In support of strategy no. 1) Snag management practices could be designed to retain snags along forest edges in areas used by nesting western bluebirds, and artificial nest boxes could be used to help support western bluebird populations in areas where snags are not available as nesting structures.
2. (In support of strategy no. 1) In burned ponderosa pine-Douglas-fir forests selected for postfire salvage logging, retain about 65 snags per ha (26 per acre) of snags >23 cm (9 in) d.b.h.
3. (In support of strategy nos. 2 and 3) Use wildfire and prescribed fire to restore natural forest openings and enhance shrub understories to attract insect prey.

4. (In support of strategy no. 3) Accelerate development of mid-successional stages of ponderosa pine to old forests by silvicultural treatments of prescribed underburning and thinning of small-diameter trees (<25 cm [9 in] d.b.h.).
5. (In support of strategy no. 4) Discourage spread of exotic plants by minimizing human-associated disturbance activities.

GROUP 30--ASH-THROATED FLYCATCHER AND BUSHTIT

Results

Species ranges, source habitats, and special habitat features--Group 30 consists of the bushtit and ash-throated flycatcher. The bushtit is a year-long resident in the basin, whereas the ash-throated flycatcher is a summer migrant. For both the ash-throated flycatcher and the bushtit, the basin constitutes the northern edge of their ranges. Both species have similar distributions within the basin, occurring along the western and southern extent of the basin (fig. 90).

Vol. 2, Figure 90--Ranges of species in group 30 within the basin.

The bushtit and ash-throated flycatcher depend on a similar mix of source habitats (vol. 3, appendix 1, table 1), including mixed-conifer woodlands, juniper/sagebrush woodlands, Oregon white oak, and mountain mahogany. Cottonwood/willow in the old-forest multi-storied structural stage also is considered source habitat for the ash-throated flycatcher.

Ash-throated flycatchers nest in cavities (either natural, woodpecker-excavated, or human-made [nest boxes]) of taller trees and snags (Austin and Russell 1972, Dunning and Bowers 1990, Sharp 1992). Snags were identified as a special habitat feature for ash-throated flycatchers (vol. 3, appendix 1, table 2). Bushtits place their nests in tall shrubs. Both species forage on arthropods.

Broad-scale changes in source habitats--Source habitats for this group historically were distributed within the western and southern parts of the basin, and watersheds with habitat appeared to be disjunct (fig. 91A). Currently, source habitats are more abundant and in some areas more continuous in distribution (fig. 91B). The largest concentration of both current and historical habitats is within the southern part of the Columbia Plateau (figs. 91A, B). The watersheds with increases in source habitats were most often the same as or adjacent to watersheds that supported source habitats historically (figs. 91A, B).

Overall, source habitats for this group strongly increased within the basin. Over 60 percent of the watersheds in the basin had strongly increasing trends, whereas about 17 percent had decreasing trends (fig. 92). Nearly 50 percent or more of the watersheds in seven of the nine ERUs with greater than 1 percent of the area as source habitats had strongly increasing trends since the historical period (fig. 92). These were the Upper Klamath, Northern Great Basin, Columbia Plateau, Blue Mountains, Owyhee Uplands, Upper Snake, and Snake Headwaters. Only the Northern Cascades had a greater number of watersheds with decreasing rather than increasing amounts of source habitat (fig. 92). The Southern Cascades generally had no net trend (fig. 92). The amount of source habitat in the Northern Glaciated Mountains is minimal (<1 percent of the ERU) (vol. 3, appendix 1, table 3).

Vol. 2, Figure 91--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 30.

Vol. 2, Figure 92--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 30, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--The increasing trend in source habitats was attributed to increases in the juniper/sagebrush cover type (vol. 3, appendix 1, table 4). The extent of juniper/sagebrush woodlands has more than doubled in the basin, primarily because of excessive livestock grazing and fire suppression (Hann and others 1997).

Broad-scale trends in the other source habitat types, especially old-forest cottonwood-willow, Oregon white oak, and mountain mahogany, are difficult to determine at the 1-km² (0.4 mi²) scale of analysis because of small patch size or linear configuration of these cover types across the basin.

Condition of special habitat features--The trend and condition of nest cavities for ash-throated flycatchers are unknown. Presumably, as the number of juniper trees increases, the aging of these juniper will develop natural cavities as snags develop and older branches fall off.

Other factors affecting the group--The primary prey for these species during the breeding season is insects (Ehrlich and others 1988, Sharp 1992). Native understory shrubs and grasses provide important substrates for production of insects, and excessive grazing can reduce or eliminate many of these key substrates for insects.¹²

A common management action is to reduce the densities of juniper especially where encroachment of or densities of junipers have increased. Juniper removal may improve rangeland productivity and restore native biodiversity in some areas; however, management efforts to remove juniper trees would negatively affect source habitats for group 30.

Population status and trends--Data for ash-throated flycatchers and bushtits in the basin were insufficient to determine a population trend. Because both species have naturally low population numbers and narrow distributions, specialized monitoring techniques are required to estimate their numbers (Saab and Rich 1997).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 30 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--Primary issues affecting source habitats for ash-throated flycatchers and bushtits areas follows:

1. For ash-throated flycatchers, loss of trees with natural cavities or trees suitable for excavation by other species because of juniper removal.
2. Degradation and loss of native understory shrubs and grasses that provide substrates for arthropod prey.

Potential strategies--The issues identified above suggest the following broad-scale strategies would be effective in contributing to the long-term persistence of bushtits and ash-throated flycatchers:

1. (To address issue no. 1) Retain contiguous blocks of mature juniper/sagebrush, especially in areas containing old junipers with cavities and hollow centers for potential nest sites of ash-throated flycatchers. Old-growth specimens usually have round or flat tops as compared to young, actively growing individuals that have a symmetrical, cone-shaped top (Oregon Department of Fish and Wildlife 1994)
2. (To address issue no. 2) Protect and restore native understory shrubs and grasses in source habitats.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Consider site-specific ecological potential and response to management before removing juniper trees.
2. (In support of strategy no. 1) Retain junipers with cavities and hollow centers that are potential nest sites for ash-throated flycatchers.
3. (In support of strategy no. 1) Retain blocks of old-growth juniper during juniper control projects.
4. (In support of strategy no. 2) Restrict the use of herbicides, pesticides, and grazing in areas with contiguous blocks of source habitat that have intact native understories.
5. (In support of strategy no. 2) Restore native understories through seedings and plantings of native shrubs and grasses.
6. (In support of strategy no. 2) Minimize the likelihood of invasion of exotic vegetation by minimizing human-associated disturbances such as road building, motorized activity, grazing, and mining.

GROUP 31--FERRUGINOUS HAWK, BURROWING OWL, SHORT-EARED OWL, VESPER SPARROW, LARK SPARROW, WESTERN MEADOWLARK, and PRONGHORN

Results

Species ranges, source habitats, and special habitat features--Group 31 consists of breeding habitat for the migratory ferruginous hawk, burrowing owl, vesper sparrow, lark sparrow, and western meadowlark, and year-round habitat for the short-eared owl and pronghorn. The short-eared owl, vesper sparrow, and western meadowlark are the most widely distributed species within this group (fig. 93), occurring throughout the basin. Less widely distributed are the burrowing owl and lark sparrow, which are both

absent from the mountainous portions of central and northern Idaho (fig. 93). The ferruginous hawk uses less of the basin but is still widespread in the lower elevations (fig. 93). The least widely distributed species in this group is the pronghorn, which currently occupies most of the Northern Great Basin ERU, a large part of the Owyhee Uplands ERU, and small, disjunct areas over the southern half of the basin (fig. 93). In contrast, the historical range of the pronghorn included almost all of southern Idaho and eastern Oregon (fig. 93). Nelson (1925) stated that pronghorn historically occurred in Washington as well, but Yoakum (1978) disagreed. We have followed the recommendations of the latter author.

Vol. 2, Figure 93a, b--Ranges of species in group 31 within the basin.

Source habitats for this group include various shrub, grass, and herbaceous cover types (vol. 3, appendix 1, table 1). All seven species have source habitats in big sagebrush and fescue-bunchgrass cover types, six share low sagebrush, and five have source habitats in juniper/sagebrush, mountain big sagebrush, native forb, and wheatgrass bunchgrass types. Whereas particular plant species may differ geographically, a key feature of this group is their preference for open cover types with a high percentage of grass and forbs in the understory. All species use the shrub component of the vegetation directly for nest sites, perch sites, or hiding cover. Pronghorn move into areas of higher shrub cover during winter. The ferruginous hawk is the only species that will use trees, especially junipers, which provide preferred nest sites in some geographic areas.

Burrowing owls depend on burrows and natural cavities in lava flows or rocky areas for nest sites; thus, burrows are a special habitat feature for this species (vol. 3, appendix 1, table 2). Burrows are almost always provided by burrowing mammals such as ground squirrels, marmots, prairie dogs, coyotes, and badgers, and the use of an area by owls may be closely tied to populations of these mammals (Haug and Oliphant 1990, Rich 1986, Thomsen 1971).

Populations (White and Thurow 1985) and productivity (Bechard and Schmutz 1995, Schmutz and Hungle 1989, Steenhof and Kochert 1985) of the ferruginous hawk fluctuate in response to prey population densities. Similarly, breeding populations of the short-eared owl are nomadic, and high densities of breeding birds may occur when rodent densities are high (Marti and Marks 1989). Thus, the status of all three raptors in this group is rather closely tied to the status of various mammal populations. Notably, these three raptor species are more tolerant of degraded shrubsteppe habitats with exotic vegetation than are other species in this group.

Significant correlations were documented between the coverage of grass and the densities of western meadowlark ($r = 0.62$, $P < 0.001$) and lark sparrow ($r = 0.37$, $P < 0.05$) (Wiens and Rotenberry 1981). Similar correlations occurred for the coverage of litter and these songbird species $r = 0.36$, $P < 0.05$ and $r = 0.34$, $P < 0.05$, respectively).

Pronghorn may depend on free water during summers of dry years when they cannot meet water requirements from succulent forbs (Beale and Smith 1970, Clemente and others 1995). In most years, however, availability of free water probably does not affect pronghorn habitat use (Deblinger and Alldredge 1991).

Broad-scale change in source habitats--Historically, source habitats for this group were widely available throughout the basin, but particularly in the Northern Great Basin, Columbia Plateau, Owyhee Uplands, and Upper Snake ERUs (fig. 94A). The most contiguous shrubsteppe habitat occurs at lower elevations, and source habitats for this group become less extensive at higher elevations. This is demonstrated by

the narrow band of watersheds with 25 to 50 and 0 to 25 percent of area in source habitats within higher elevation ERUs (fig. 94B).

The projected extent of decreasing and strongly decreasing trends in source habitats was dramatic (fig. 94C). The Columbia Plateau and Upper Snake ERUs were dominated by decreasing trends, the latter having no watersheds with increasing trends. In contrast, large, contiguous portions of the Northern Great Basin and Owyhee Uplands ERUs, in areas of higher elevation and precipitation, show a stable trend and continue to provide source habitats for this group.

Basin-wide, 54 percent of the watersheds had moderately or strongly declining trends in source habitats (fig. 95). The Columbia Plateau ERU historically provided the most watersheds with source habitats for this group (fig. 95), but over 72 percent of the watersheds in that ERU had moderately or strongly declining trends. The second most important ERU, the Owyhee Uplands, had stable trends in about 81 percent of its watersheds, but another 19 percent were moderately or strongly declining. The number of watersheds with moderately and strongly declining trends in source habitats outnumbered those with increasing trend in all other ERUs (fig. 95) except the Central Idaho Mountains.

Vol. 2, Figure 94--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 31.

Vol. 2, Figure 95--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 31, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--The single largest loss in cover types within the basin has been the decline in big sagebrush (Hann and others 1997). Habitat losses were also significant for fescue-bunchgrass and wheatgrass bunchgrass (Hann and others 1997). This loss was most striking in the Columbia Plateau and Upper Snake ERUs (figs. 94C and 95). Other notable reductions include the near complete loss of source habitats in the Upper Clark Fork and Lower Clark Fork ERUs.

In the Columbia Plateau, major losses from historical conditions occurred in big and mountain sagebrush types, which declined by nearly half and over three-fourths, respectively (vol. 3, appendix 1, table 4). Native grass cover types also were heavily impacted, with a three-fourths decline in wheatgrass bunchgrass, and a nearly total loss of fescue-bunchgrass (Hann and others 1997). In the lower elevations of the Owyhee Uplands, big sagebrush was reduced by 25 percent (Hann and others 1997). Fescue-bunchgrass types had significant negative declines in nine ERUs (Hann and others 1997). Nearly all of the native forb cover type, source habitats for five of these species, was converted to other cover types (Hann and others 1997). Native forbs were projected to have covered a small portion of the basin historically but likely provided important local breeding habitats within larger blocks of more xeric vegetation.

In the Central Idaho Mountains ERU, nearly 33 percent of the watersheds had strongly increasing trends (fig. 95). This was attributed to large relative increases in juniper/sagebrush, juniper woodlands, and low sagebrush, all of which covered only a small fraction of the unit. A similar situation resulted in strongly increasing trends in the Northern Cascades, Blue Mountains, Northern Great Basin, and Snake Headwaters ERUs (fig. 95; vol. 3, appendix 1, table 4; Hann and others 1997). Any increases in wheatgrass bunchgrass or native forb cover types (vol. 3, appendix 1, table 4) should be viewed with caution because these cover types can be dominated by exotic vegetation, which is not considered source habitat for species of this group.

Several factors contributed to large-scale losses of sagebrush and fescue-bunchgrass habitats; foremost was conversion to agriculture. Agricultural lands have increased significantly in every ERU in the basin (vol. 3, appendix 1, table 4). In fact, the largest transitions among terrestrial communities from the historical to current periods were that of upland shrubland and upland hermland to agriculture (Hann and others 1997). This transition explains much of the pattern evident in figure 94C.

A second factor contributing to loss of sagebrush habitat was conversion of shrubsteppe vegetation to exotic forbs and annual grass. Notable portions of the Owyhee Uplands and Upper Snake ERUs underwent a conversion from upland shrubland to exotic hermland (Hann and others 1997). Conversion of native vegetation to exotics was augmented by the increased frequency of wildfire and by improper grazing (Quigley and others 1996, USDA Forest Service 1996).

Condition of special habitat features--Burrowing owls rely on burrows provided by burrowing mammals for nest sites (Haug and Oliphant 1990, Rich 1986, Thomsen 1971). Populations of many burrowing mammals have declined as a result of various pest control programs, which may have reduced nest site availability for burrowing owls. No special habitat features were identified for other members of this group.

Other factors affecting the group--Losses of native perennial grass and forb understories within the sagebrush types, associated with intensive livestock grazing, cheatgrass invasions, and noxious weed invasions, are microhabitat changes that could not be evaluated by our broad-scale analysis. Because species in group 31 favor grass or shrub-grass types for nesting, foraging, or hiding, we know that the grass component of historical shrublands was important (for example, Wiens and Rotenberry 1981, Marti and Marks 1989). Removal of grass cover by livestock potentially has detrimental effects on the short-eared owl (Marti and Marks 1989). Finer scale analysis is needed to determine the extent of this problem because the broad-scale data may show source habitats in upland shrub types, where the shrubs are present but the understory is gone. The presence of livestock also may attract brown-headed cowbirds and subsequently increase the incidence of brood parasitism (Robinson and others 1995). The western meadowlark and vesper sparrow are common cowbird hosts, whereas the lark sparrow is only occasionally parasitized (Ehrlich and others 1988).

Ferruginous hawks prefer trees for nest sites, particularly junipers (Jasikoff 1982), and are most common in the juniper/sagebrush ecotone (Powers and others 1973, Smith and Murphy 1973, Thurow and others 1980). Expansion of juniper woodlands and juniper/sagebrush in the basin as a result of fire suppression likely has benefitted the species.

Fields of hay and cereal grains attract vesper sparrows (Perritt and Best 1989) and western meadowlarks (Lanyon 1994) for nesting, where nests, young, or adults may be destroyed during harvest. Short-eared owls and lark sparrows also are likely affected by this process. These fields function as sinks for local populations.

Species in this group evolved in shrubsteppe habitats where microbiotic crusts were broadly distributed (see Kaltenecker and Wicklow-Howard 1994). Microbiotic, or cryptogamic, crusts consist of lichens,

bryophytes, algae, microfungi, cyanobacteria, and bacteria growing on or just below the soil surface in arid and semiarid environments (Kaltenecker and Wicklow-Howard 1994); these crusts developed without large herds of grazing ungulates (St. Clair and Johansen 1993). In addition, these crusts are projected to have been widely distributed throughout the source habitats for this group, particularly in the Northern Great Basin, Owyhee Uplands, and Upper Snake ERUs but also scattered in the Columbia Plateau ERU (Hann and others 1997, map 3.59). Increasing evidence suggests that microbiotic crusts improve soil stability, productivity, and moisture retention, moderate extreme temperatures at the soil surface, and enhance seedling establishment of vascular plants (Belnap and Gardner 1993, Harper and Pendleton 1993, Johansen and others 1993, St. Clair and others 1993), thus contributing to high ecological integrity of shrubsteppe habitats. Idaho BLM has recognized the potential importance of microbiotic crusts by proposing standards for rangeland health that include the maintenance of these crusts to ensure proper functioning and productivity of native plant communities (USDI Bureau of Land Management 1997). These crusts were widely destroyed by trampling during the excessive livestock grazing of the late 1800s and early 1900s (Daubenmire 1970, MacCracken and others 1983, Mack and Thompson 1982, Poulton 1955). Currently, high-intensity grazing and altered fire regimes modify shrubsteppe plant communities and threaten the maintenance and recovery of microbiotic crusts (Belnap 1995, Kaltenecker 1997, St. Clair and Johansen 1993).

Roads, human activities, and domestic dogs are known to impact ferruginous hawks, short-eared owls, burrowing owls (Bechard and Schmutz 1995, Green and Anthony 1989, Lokemoen and Duebbert 1976, Olendorff and Stoddart 1974, Ramakka and Woyewodziec 1993, Schmutz 1984, White and Thurow 1985) and western meadowlarks (Lanyon 1994). Harassment of pronghorn by snowmachine and all-terrain vehicles stresses animals at all times of the year (Autenrieth 1978). Pronghorn also avoid sheep dogs (Yoakum and O'Gara 1990). Human disturbance might be especially significant for those species that are attracted to features of the agricultural-shrubland or agricultural-grassland contact zones, that is, burrowing owl, short-eared owl, and pronghorn.

Recreational shooting of marmots and ground squirrels impacts burrowing owls because the owls are accidentally or deliberately shot, whereas more general illegal shooting impacts short-eared owls (Marti and Marks 1989). Pesticide use leads to direct mortality in burrowing owls, short-eared owls (Marti and Marks 1989), and western meadowlarks (Griffin 1959) and an indirect loss in burrowing owls through a reduction in the populations of burrowing mammals.

Pronghorn movement is restricted or completely impeded by net-wire and other fences that prevent them from crossing beneath the lower strand (Helms 1978, Oakley and Riddle 1974, Yoakum 1980). Roads are readily crossed by pronghorn, but snow accumulating in roadside ditches also might present barriers to movement during winter (Bruns 1977).

Population status and trends--Based on BBS data summarized for the basin (Saab and Rich 1997), significant declines were reported for the period 1966-94 for western meadowlark (-0.8 percent per yr, $\underline{n} \geq 14$, $P < 0.10$) and lark sparrow (-2.9 percent per yr, $\underline{n} \geq 14$, $P < 0.05$). Saab and Rich (1997) identified western meadowlark and lark sparrow as two of 15 species that are of high concern to management under all future management themes for the basin. Vesper sparrow, burrowing owl, and ferruginous hawk had stable population trends within the basin for the same time period (Saab and Rich 1997). In physiographic region 89 (Columbia Plateau), which corresponds to much of the range of this group within the basin, trends over the period 1966-95 (Sauer and others 1996) were positive for the ferruginous hawk (+6.3 percent per yr, $\underline{n} = 18$, $\underline{P} < 0.05$).

Burrowing owl populations are increasing across the West (+6.3 percent per yr; $\underline{n} = 116$, $\underline{P} < 0.001$; Sauer and others 1996). No detectable trend was found for the short-eared owl in the basin (Saab and Rich 1997) or in physiographic region 89 (Columbia Plateau; Sauer and others 1996). Marti and Marks (1989) reported that short-eared owl numbers were stable, with fluctuating populations.

Burrowing owls, short-eared owls, and ferruginous hawks are not adequately monitored by the BBS technique so apparent population trends, or the lack thereof, for these species may not be reliable (Saab and Rich 1997).

An estimated 99 percent of the continental pronghorn population was killed by indiscriminate hunting between 1850 and 1900, but numbers have increased dramatically since then in Idaho and Oregon (Yoakum 1968, 1978, 1986a; Yoakum and O'Gara 1990). Populations reached peaks in 1989 of 21,800 in Idaho and 22,650 in Oregon (O'Gara 1996). The most recent estimates (1995) are 12,500 in Idaho and 17,122 in Oregon (O'Gara 1996).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 31 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The condition of the habitat for group 31 can be summarized by the Composite Ecological Integrity Ratings (Quigley and others 1996, p. 122) that show most of the habitat to have a "low" rating. Fescues and bunchgrasses--critical habitat components for this group-- ". . . were irreversibly modified by extensive grazing in the late 1800s and early 1900s" (USDA Forest Service 1996, p. 51). Most of the current habitat for this group was classified into Rangeland Clusters 5 (generally corresponding to much of the Owyhee Uplands ERU) and 6 (generally the Northern Great Basin, Owyhee Uplands, and Upper Snake ERUs) where the primary risk to ecological integrity is "continued declines in herbland and shrubland habitats" (Quigley and others 1996, p. 112, 114). Further, Rangeland Cluster 6 has the additional risk of being ". . . highly sensitive to overgrazing and exotic grass and forb invasion" (Quigley and others 1996, p. 114). These widespread and overriding issues provide a clear statement of the problems facing this group over the long term.

Primary issues areas follows:

1. Permanent and continued loss of large acreage of shrubsteppe and fescue-bunchgrass habitat because of agricultural conversion, brush control, and cheatgrass invasion.
2. Soil compaction and loss of the microbiotic crust.
3. Adverse effects of human disturbance. For the burrowing owl, a primary issue is the loss of nesting burrows through poisoning and recreational shooting of burrowing mammals. For ground-nesting birds, the issue is nest mortality in agricultural fields from farm machinery during spring weed control and early harvests. For pronghorn, a primary issue is disruption of movement patterns because of fence constructions that inhibit passage. For all species in group 31, the issue is general disruption of breeding activity and movements because of human intrusion.

Potential strategies--

1. (To address issue no. 1) Identify and conserve large remaining areas (contiguous habitat >1000 ha [2,470 acres]) of shrubsteppe vegetation where ecological integrity is still relatively high, and manage to promote their long-term sustainability. Large contiguous blocks of public land in the Northern Great Basin and Owyhee Uplands are the most obvious sites. These generally include the subbasins in Rangeland Cluster 5 (Quigley and others 1996). These areas will provide long-term habitat stability for populations and provide the anchor points for restoration, corridor construction, and other landscape-level management.

2. (To address issue no. 1) Restore the grass and forb components of the shrubsteppe cover types to approximate historical levels throughout the basin.
3. (To address issue no. 2) Restore the microbiotic crust in ERUs where potential for redevelopment is high; that is, in areas near propogule sources that have suitable soil, vegetation, and climatic characteristics [see Belnap 1993, Belnap 1995, Kaltenecker 1997, Kaltenecker and Wicklow-Howard 1994]. ERUs with highest potential for redevelopment include the Northern Great Basin, Owyhee Uplands, Upper Snake, and, to a lesser extent, the Columbia Plateau (map 3.59 in Hann and others 1997).
4. (To address issue no. 3) Maintain burrows for nesting and roosting by burrowing owls. Reduce mortality of ground-nesting birds in agricultural areas. Construct fences in pronghorn range that allow pronghorn passage. Minimize the adverse effects of human intrusion.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Identify large areas of high ecological integrity to be managed for sustainability by analyzing current vegetation, precipitation patterns, elevation, temperature (Klemmedson and Smith 1964, Morrow and Stahlman 1984, Stewart and Hull 1949) and the presence of priority species in this group. These sites most likely will be successful on large areas of Federal land managed by BLM. Evaluation criteria for protection or enhancement include maintaining or increasing the size of smaller patches, preventing further habitat fragmentation, and protecting or increasing the size and integrity of corridors among patches, all in connection with the location of core areas.
2. (In support of strategy no. 1) Explore options under the CRP (Johnson and Igl 1995), or develop other incentive programs, to encourage restoration of agricultural areas to native cover types. Focus on areas that would increase patch size or links with existing source habitat patches.
3. (In support of strategy no. 2) Use fire prevention and suppression to retard the spread of cheatgrass in areas that are susceptible to cheatgrass invasion but currently are dominated by native grass species. Planting of fire-resistant vegetation through "green stripping" is being experimentally tested (Pellant 1994) and may be used to protect existing vegetation.
4. (In support of strategy no. 2) Restore selected areas of cheatgrass monocultures through seeding and other manipulations (Allen 1995, Daubenmire 1970, Evans and Young 1978, Hosten and West 1994, Kennedy 1994, Monsen and McArthur 1995, Ogg 1994, Whisenant 1995, Yoakum 1986b), where such restoration would increase the size of existing shrubsteppe patches or provide links between patches.
5. (In support of strategy no. 2) Restore native vegetation by appropriate treatments and seedings of native shrub, grass, and forb species.
6. (In support of strategy no. 2) Design livestock grazing systems to promote an abundance of forbs and grasses in the understory (Yoakum 1980).
7. (In support of strategy no. 3) Encourage the redevelopment of microbiotic crust by reducing or eliminating livestock grazing in areas where restoration of microbiotic crusts is encouraged (Mack and Thompson 1982, St. Clair and others 1993). Explore the use of ground-based and aerial soil inoculation to increase the speed and extent of dispersal of the organisms that create microbiotic crust (Belnap 1993).

8. (In support of strategy no. 4) Allow burrowing mammals such as ground squirrels and marmots to persist or expand to provide nesting burrows for burrowing owls (Coulombe 1971; Gleason and Johnson 1985; Rich 1984, 1986). Provide artificial burrows for burrowing owls where burrowing mammals must be controlled (Trulio 1995).
9. (In support of strategy no. 4) Modify agricultural practices to minimize direct mortality of nesting birds by delaying hay mowing until young birds are fledged (Clark 1975, Rodenhouse and others 1995, Vickery 1996). Avoid surface tillage for spring weed control. An alternative is to use the "undercutting" method, which is much less detrimental to meadowlarks (Rodgers 1983).
10. (In support of strategy no. 4) Control, reduce, or eliminate pesticide applications in and around agricultural areas, especially in the Columbia Plateau ERU where source habitats are small and virtually all surrounded by agricultural lands (USDA Forest Service 1996). The Upper Snake ERU, and to a lesser extent the Owyhee Uplands, also have relatively many miles of interface with agricultural lands.
11. (In support of strategy no. 4) Avoid construction of net-wire and similar fences in pronghorn habitat or in pronghorn migration routes (Oakley and Ridle 1974). Modify existing fences and construct new fences in pronghorn range with the following specifications (these are standard policy on BLM lands occupied by pronghorns): bottom wire at least 41 cm (16 in) from the ground and smooth, not barbed; next wire up is 66 cm (26 in) from the ground; top wire is 91 cm (36 in) from the ground (Yoakum 1980).
12. (In support of strategy no. 4) Protect pronghorn winter ranges and fawning areas from intrusion by snow machines and all-terrain vehicles (Autenrieth 1978) through timed access control and area closures. Minimize access roads and, where possible, locate them on the periphery of the pronghorn use areas (Autenrieth 1978). Provide artificial nesting structures in areas away from human disturbance to attract ferruginous hawks to safer sites (Apple 1994, Niemuth 1992, Schmutz 1984). Protect burrowing owl nesting sites from disturbance by domestic dogs (Green and Anthony 1989, Martin 1983).

GROUP 32--PREBLE'S SHREW, UINTA GROUND SQUIRREL, WHITE-TAILED ANTELOPE SQUIRREL, WYOMING GROUND SQUIRREL, WASHINGTON GROUND SQUIRREL, STRIPED WHIPSNAKE, LONGNOSE SNAKE, GROUND SNAKE, MOJAVE BLACK-COLLARED LIZARD, AND LONGNOSE LEOPARD LIZARD

Results

Species ranges, source habitats, and special habitat features--Group 32 consists of year-round habitat for the residents in this group: Preble's shrew, Uinta ground squirrel, white-tailed antelope squirrel, Wyoming ground squirrel, Washington ground squirrel, striped whipsnake, longnose snake, ground snake, Mojave black-collared lizard, and longnose leopard lizard.

Mammals--Little is known about the Preble's shrew, but they may be widely distributed in the basin (fig. 96), based on records from the area's borders (Cornely and others 1992, Zeveloff and Collett 1988). Among the four species of ground squirrels, the Uinta is restricted to the upper end of the Snake River drainage in the Snake Headwaters, Upper Snake, and Central Idaho Mountains ERUs (fig. 96). The range of the white-tailed antelope squirrel occurs in the Northern Great Basin and Owyhee Uplands ERUs and is nearly distinct from that of the Uinta ground squirrel (fig. 96). Two subspecies of the Wyoming ground squirrel occur in the basin, *Spermophilus elegans nevadensis* that overlaps with the

antelope squirrel in the Owyhee Uplands, and *Spermophilus elegans aureus* that overlaps with the Uinta ground squirrel in northeastern Idaho (fig. 96). Finally, both the current and historical (fig. 96) range of the Washington ground squirrel is allopatric with the other three species, being confined almost entirely to the northern part of the Columbia Plateau ERU. The current range of the latter species is reduced and disjunct compared to the historical period.

Vol. 2, Figure 96a, b, c--Ranges of species in group 32 within the basin.

Reptiles--The striped whipsnake is widely distributed at lower elevations in Washington, Oregon, and Idaho (fig. 96). Narrowly distributed and largely sympatric, the longnose snake and ground snake occur only in the Owyhee Uplands (fig. 96). The Mojave black-collared lizard has a distribution similar to the previous two species but has an additional portion of its range in the Northern Great Basin (fig. 96). Finally, the longnose leopard lizard is found largely in the Owyhee Uplands but has disjunct populations in the Northern Great Basin, Upper Snake, Columbia Plateau, and Southern Cascades ERUs.

Source habitats for group 32 include several shrub, grass, and herbaceous cover types (vol. 3, appendix 1, table 1). All 10 species have source habitats in big sagebrush, mountain big sagebrush, fescue-bunchgrass, and wheatgrass bunchgrass types. Ten species also have source habitats in low sage, whereas eight share juniper/sagebrush or mountain mahogany.

The striped whipsnake uses cliffs and talus where they occur in source habitats; these are special habitat features for this species (vol. 3, appendix 1, table 2). Preble's shrew requires a good understory of forbs and grasses and a dense overstory of sagebrush; it is associated with more mesic sites near ephemeral and perennial streams (Ports and George 1990). Down logs provide important foraging and hiding cover (vol. 3, appendix 1, table 2). Washington ground squirrels prefer deeper soils with less clay at 10 cm (4 in) and at 50 cm (20 in) compared to unoccupied sites (Betts 1990).

Talus slopes, canyon rims, and shadscale habitats are preferred over other types by ground snakes and collared lizards (Diller and Johnson 1982, Whitaker and Maser 1981). Collared lizards similarly prefer rock outcrops and sparse vegetation (Sanborn and Loomis 1979). Striped whipsnakes are much more apt to be encountered on canyon rims than on mid-slopes or in canyon bottoms (Gerber and others 1997).

Broad-scale changes in source habitats--Historically, source habitats for this group were projected to occur throughout the basin, with greatest concentrations in the Northern Great Basin, Columbia Plateau, Owyhee Uplands, and Upper Snake ERUs (fig. 97A). Substantial amounts of source habitats also occurred in the Blue Mountains, Northern Glaciated Mountains, Central Idaho Mountains, and Upper Klamath ERUs. Only the most mountainous and forested regions did not support members of this group.

The extent of decreasing and strongly decreasing trends in source habitats was dramatic (fig. 97C), particularly for the state of Washington, the northern half of Oregon, and the upper Snake River drainage. Nine ERUs had declining trends for most watersheds, whereas only two ERUs (Northern Great Basin and Owyhee Uplands) showed stable trends. The only noteworthy source habitat increases were in the Central Idaho Mountains (fig. 98).

Basin-wide, 56 percent of the watersheds showed a moderately or strongly declining trend in source habitats (fig. 98). The Columbia Plateau ERU historically provided the most watersheds with source habitats for this group (fig. 98). But over 83 percent of the watersheds in that ERU had moderately or

strongly declining trends and only about 5 percent were increasing. In the Blue Mountains, nearly 84 percent of the watersheds had moderately or strongly declining trends (fig. 98), and <4 percent were increasing. The Upper Snake ERU had no watersheds with increasing trends (fig. 98) and over 67 percent with moderately or strongly declining trends. In the Owyhee Uplands, over 81 percent of watersheds had stable trends, and 17 percent had moderately or strongly declining trends (fig. 98).

Vol. 2, Figure 97--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 32.

Vol. 2, Figure 98--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 32, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Declines in source habitats were primarily due to reductions in the amount of big sagebrush, fescue-bunchgrass, wheatgrass bunchgrass, and interior ponderosa pine (Hann and others 1997). These losses were most striking in the Columbia Plateau and Upper Snake ERUs (fig. 97B; vol. 3, appendix 1, table 4). In the Columbia Plateau, big and mountain sagebrush types declined by nearly half and three-quarters, respectively, from historical conditions. Wheatgrass bunchgrass declined by three-fourths and fescue-bunchgrass was nearly eliminated (Hann and others 1997) in the Columbia Plateau.

Large-scale losses of sagebrush and native bunchgrass habitats were primarily due to conversion to agriculture. Basin-wide, the largest transitions among terrestrial communities from the historical to current periods were that of upland shrubland and upland herbland to agricultural (Hann and others 1997).

Another factor contributing to loss of sagebrush habitat is conversion of shrubsteppe vegetation to exotic forbs and annual grass. Substantial portions of the Owyhee Uplands and Upper Snake ERUs have undergone conversions from upland shrubland to exotic herbland (Hann and others 1997). Noteworthy increases in this cover type have occurred in all major shrubsteppe ERUs. Conversion of native vegetation to exotics is augmented by the increased frequency of wildfire and by improper grazing (Braun and others 1976, Daubenmire 1970, Evans and Young 1978, Quigley and others 1996, USDA Forest Service 1996).

Any increases in wheatgrass bunchgrass or native forb cover types (vol. 3, appendix 1, table 4) should be viewed with caution because these cover types can be dominated by exotic vegetation, which is not considered source habitat for species of this group. Additionally, in some cases the wheatgrass bunchgrass cover type was misclassified as an upland herbland group instead of an early-seral forest group that was created as a result of timber harvest or recent large-scale wildfires (see Hann and others 1997).

Relatively large increases have occurred in the source habitats of juniper woodlands (tripled), mountain mahogany (tripled), juniper/sagebrush (doubled), and low sage (one-third increase) in the Central Idaho

Mountains (Hann and others 1997) (vol. 3, appendix 1, table 4).

Condition of special habitat features--The availability of mesic sites used by the Preble's shrew has declined as part of the general and widespread decline in riparian habitat conditions throughout the basin (Lee and others 1997, Quigley and others 1997).

Cliffs and talus habitat for the striped whipsnake, although difficult to measure at the scale of this analysis, were estimated to be in much the same condition now as historically.

Other factors affecting the group--Poisoning and other eradication potentially affect populations of all four species of ground squirrels. Ground squirrels also are popular targets for recreational shooting. The Mojave black-collared lizard, longnose leopard lizards, and longnose snakes use small-mammal burrows for cover (Beck and Peterson 1995, Brown and others 1995, Nussbaum and others 1983, Pough 1973), and therefore could be indirectly affected by both poisoning and shooting. The effect of these factors on these species in the basin is unknown.

Accidental and deliberate mortality of snakes potentially increases with increased roading and traffic in the basin. Although the three species of snakes in this group may not be as frequently killed by vehicles as are some more common species (such as gopher snake and western rattlesnake), increasing human access to source habitats will predictably result in more deliberate killing of snakes. Currently, large areas of the Owyhee Uplands ERU support moderate to high road densities (see figs. 21 and 22 and "Species and Groups Affected by Factors Associated with Roads" in vol. 1).

The typical small size of Washington ground squirrel colonies makes them vulnerable to extirpation (Tomich 1982). Source habitats for this species were estimated to have undergone the fourth greatest decline among 91 broad-scale species of focus analyzed in this report (vol. 1, table 7). Washington ground squirrels may benefit from corridors of vegetation created by cultivation that allow exchange among colonies and general dispersal (Betts 1990).

Four of the reptilian species of this group (Mojave black-collared lizard, longnose leopard lizard, longnose snake, and ground snake), are located in isolated disjunct areas within the basin that make them vulnerable to extirpation.

Areas dominated by dense stands of cheatgrass or other exotic plants may preclude use by longnose leopard lizards (Stebbins 1985), longnose snakes (Beck and Peterson 1995), and collared lizards. In the Owyhee Uplands, areas with low vegetative cover and high amounts of bare ground or rock have the highest lizard densities (Whitaker and Maser 1981). In a study of off-road vehicle and grazing effects in the Mojave Desert in California, leopard lizards were found only in plots unused by off-road vehicles (compared with moderately and heavily used plots), and were absent from grazed plots (Busack and Bury 1974).

Because reptiles are increasingly popular as pets, all reptile species in this group, but particularly the lizards, are potentially affected by collecting (Lehmkuhl and others 1997). This impact will increase as the human population in the basin increases.

Soil compaction caused by livestock grazing could negatively affect both the longnose snake and ground snake. These burrowers benefit from loose, sandy, and friable soils (Beck and Peterson 1995, Nussbaum and others 1982).

Species in this group evolved in shrubsteppe habitats, where microbiotic crusts were broadly distributed (see Kaltenecker and Wicklow-Howard 1994). Microbiotic, or cryptogamic, crusts consist of lichens, bryophytes, algae, microfungi, cyanobacteria, and bacteria growing on or just below the soil surface in

arid and semiarid environments (Kaltenecker and Wicklow-Howard 1994), and they developed without large herds of grazing ungulates (St. Clair and Johansen 1993). These crusts are projected to have been widely distributed throughout the source habitats for this group, particularly in the Northern Great Basin, Owyhee Uplands, and Upper Snake ERUs but also scattered in the Columbia Plateau ERU (Hann and others 1997, map 3.59). Increasing evidence indicates that microbiotic crusts improve soil stability, productivity, and moisture retention; moderate extreme temperatures at the soil surface; and enhance seedling establishment of vascular plants (Belnap and Gardner 1993, Harper and Pendleton 1993, Johansen and others 1993, St. Clair and others 1993), thus contributing to high ecological integrity of shrubsteppe habitats. Idaho BLM has recognized the potential importance of microbiotic crusts by proposing standards for rangeland health that include the maintenance of these crusts to ensure proper functioning and productivity of native plant communities (USDI BLM 1997). These crusts were widely destroyed by trampling during the excessive livestock grazing period of the late 1800s and early 1900s (Daubenmire 1970, MacCracken and others 1983, Mack and Thompson 1982, Poulton 1955). Currently, high-intensity grazing and altered fire regimes modify shrubsteppe plant communities and threaten the maintenance and recovery of microbiotic crusts (Belnap 1995, Kaltenecker 1997, St. Clair and Johansen 1993).

Population status and trends--Quantified population trends are not available for any of these species. The Washington ground squirrel has experienced range contraction (figs. 96), with 23 colonies in Washington and 12 in Oregon disappearing from 1980 to 1989. This area includes most of the colonies in the northern part of the basin (Betts 1990). This decline is wholly consistent with known habitat loss.

Lehmkuhl and others (1997) projected an historical decline in populations of the Mojave black-collared lizard as a result of the cumulative effects of habitat loss because of agricultural conversion, exotic weed invasion, and reservoir development.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 32 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The condition of the habitat for group 32 can be summed up by the Composite Ecological Integrity Ratings (Quigley and others 1996) that show most of the habitat to have a "low" rating. Most of the current habitat for this group is classified into Rangeland Clusters 5 (generally corresponding to much of the Owyhee Uplands ERU) and 6 (generally the Northern Great Basin, Owyhee Uplands, and Upper Snake ERUs), where the primary risk to ecological integrity is "continued declines in herbland and shrubland habitats" (Quigley and others 1996). Further, Rangeland Cluster 6 has the additional risk of being ". . . highly sensitive to overgrazing and exotic grass and forb invasion" (Quigley and others 1996, p. 123). These widespread and overriding issues provide a clear statement of the problems facing this group over the long term. The results of our habitat trend analysis, combined with other literature cited here, suggest the following issues are of high priority for group 32:

1. Permanent and continued loss of large areas of shrubsteppe and fescue-bunchgrass habitat to agricultural conversion, brush control, cheatgrass invasion, and expansion of juniper woodlands and mountain mahogany.
2. Increased soil compaction and loss of the microbiotic crust.
3. Reduction in burrow availability for lizards and snakes.

4. Human-caused mortality and capture of reptiles for pets.
5. Loss of downed logs.
6. Loss of surface water and riparian vegetation.

Potential strategies--The following strategies could be used to reverse broad-scale declines in source habitats. These strategies should be applied basin-wide:

1. (To address issue no. 1) Identify and conserve remaining large areas of shrubsteppe, fescue-bunchgrass, wheatgrass bunchgrass, and other source cover types where ecological integrity is still relatively high (Gray and Rickard 1989, Rickard and Poole 1989, Schuler and others 1993, Smith 1994, Yoakum 1980). Large contiguous blocks of Federal land in the Northern Great Basin and Owyhee Uplands are the most obvious sites to consider. These generally include the subbasins in Rangeland Cluster 5 (Quigley and others 1996). However, native shrublands that currently exist on military lands in Washington State (Rickard and Poole 1989, Schuler and others 1993, Smith 1994) also are important. These core areas will provide long-term habitat stability for populations and provide the anchor points for restoration, corridor construction, and other landscape-level management.
2. (To address issue no. 1) Minimize further spread of juniper woodlands, juniper/sagebrush, and mountain mahogany that have expanded as a result of fire suppression, particularly in the Central Idaho Mountains and the Columbia Plateau.
3. (To address issue no. 2) Reduce causes of soil compaction, particularly within source habitats of the longnose snake and ground snake. This factor may be important in the Owyhee Uplands ERU in particular. Restore microbiotic crusts in ERUs with potential for redevelopment (that is, areas near propogule sources, and with suitable soil, vegetation, and climatic characteristics [see Belnap 1993, 1995; Kaltenecker 1997; Kaltenecker and Wicklow-Howard 1994]): Northern Great Basin, Owyhee Uplands, and Upper Snake ERUs and, to a lesser extent, the Columbia Plateau ERU (Hann and others, map 3.59).
4. (To address issue no. 3) Maintain and restore small-mammal populations to provide burrows for the collared lizard, longnose leopard lizard, longnose snake, and ground snake.
5. (To address issue no. 4) Determine the impact of the capture of reptiles, especially lizards, for pets. Take action as necessary to allow wild populations to persist.
6. (To address issue no. 4) Reduce the direct and indirect effects of human disturbance on populations of species within group 32.
7. (To address issue no. 5) Increase the number of downed logs in the basin.
8. (To address issue no. 6) Improve the condition of riparian systems throughout the basin.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Identify large areas of high ecological integrity to be managed for long-term protection by analyzing current vegetation, precipitation patterns, elevation, temperature (Klemmedson and Smith 1964, Morrow and Stahlman 1984, Stewart and Hull 1949) and the presence of priority species in this group. These sites are most likely to be successful on large areas of Federal

land managed by BLM. Apply special management designations as necessary to protect these sites for the long term.

2. (In support of strategy no. 1) Explore options under the CRP (Johnson and Igl 1995), or develop other incentive programs, to encourage restoration of agricultural areas to native cover types. Focus on areas that would increase patch size or links with existing source habitat patches.
3. (In support of strategy no. 1) Avoid further loss of sagebrush cover through burning, plowing, seeding, and other brush "control" methods where sagebrush cover types are below historical levels.
4. (In support of strategy no. 1) Avoid further conversion of sagebrush and native grasslands to agricultural lands through policy and land management allocations. If conversion cannot be avoided, then tracts slated for conversion will have less impact if located so as to (a) minimize further fragmentation of shrubsteppe throughout the basin; (b) avoid further reducing the size of smaller, isolated patches, particularly in the Columbia Plateau ERU; and c) avoid conversion in areas that currently occur in large blocks of moderate Composite Ecological Integrity (Quigley and others 1996), primarily in the Owyhee Uplands and Northern Great Basin ERUs.
5. (In support of strategy no. 1) Use fire prevention and suppression to retard the spread of cheatgrass in areas that are susceptible to cheatgrass invasion but currently are dominated by native grass species. Planting of fire-resistant vegetation through "green stripping" (Pellant 1994) should be examined for its value to protect existing vegetation as well as allow degraded sites a chance to recover.
6. (In support of strategy no. 1) Restore selected areas of cheatgrass monocultures through seeding and other manipulations (Allen 1995, Daubenmire 1970, Evans and Young 1978, Hosten and West 1994, Kennedy 1994, Monsen and McArthur 1995, Ogg 1994, Whisenant 1995, Yoakum 1986b) where such restoration would increase the size of existing shrubsteppe patches or provide links among patches.
7. (In support of strategy no. 1) Restore native vegetation by appropriate mechanical treatments and seedings of native shrub, grass, and forb species.
8. (In support of strategy no. 2) Apply wildland fire and grazing practices that arrest the advances of juniper woodlands in areas that historically did not support this vegetation type.
9. (In support of strategy no. 3) Reduce or eliminate livestock grazing in critical habitat for the ground and longnose snakes if soil compaction is found to contribute to population declines. Encourage the redevelopment of microbial crust by reducing or eliminating livestock grazing (St. Clair and others 1993, Mack and Thompson 1982). Explore the use of ground-based and aerial soil inoculation to increase the speed and extent of dispersal of the organisms that create microbial crust (Belnap 1993).
10. (In support of strategy no. 4) Allow burrowing mammals such as ground squirrels and marmots to persist or expand to provide burrows for the lizards in this group and for the longnose snake.
11. (In support of strategy nos. 5 and 6) Minimize accidental and deliberate killing of snakes by vehicle and by humans on foot. Road densities, which provide an index to the potential for disturbance, reveal that the Owyhee Uplands, Northern Great Basin, and northern part of the Columbia Plateau ERUs are least susceptible to disturbance (Quigley and others 1996). Determine the direct effect of recreational shooting of ground squirrels on populations in this group. Effects may be serious only in local situations where the demand for this recreation and access to squirrels coincide. Washington ground squirrels are especially vulnerable because of their limited distribution and known losses to

date. Avoid poisoning or otherwise controlling ground squirrel populations. Encourage and enforce laws that protect reptiles from collection.

12. (In support of strategy no. 8) Maintain strips of trees and snags along riparian corridors. Restore and enhance riparian and shoreline vegetation around permanent and seasonal water sources.

GROUP 33--BREWER'S SPARROW, LARK BUNTING, SAGE SPARROW, SAGE THRASHER, SAGE GROUSE, PYGMY RABBIT, AND SAGEBRUSH VOLE

Results

Species ranges, source habitats, and special habitat features--Group 33 includes breeding habitat for the migratory Brewer's sparrow, lark bunting, sage sparrow, and sage thrasher, summer and winter range for the sage grouse, and year-round habitat for the pygmy rabbit and sagebrush vole. The basin encompasses a substantial portion of the entire range of all species in this group, with the exception of the lark bunting, which is peripheral to the basin occurring only in the southeastern part of the basin (fig. 99). Both the pygmy rabbit and sage grouse (current range) have notable gaps in their distribution, with significant disjunct populations primarily in the Columbia Plateau ERU. The current range of the sage grouse also has disjunct populations occurring in the Upper Klamath and Snake Headwaters ERUs. In comparison, the historical range of the sage grouse (fig. 99) was substantially more extensive and included portions of the Columbia Plateau, Blue Mountains, Northern Glaciated Mountains, Lower Clark Fork, and the Upper Clark Fork ERUs, where the species does not occur today.

Vol. 2, Figure 99a, b--Ranges of species in group 33 within the basin.

The seven species in this group have source habitats in two structural stages of big sagebrush and mountain big sagebrush: open canopy, low-medium shrub, and closed canopy, low-medium shrub (vol. 3, appendix 1, table 1). Four of the species (pygmy rabbit, sagebrush vole, sage grouse, and sage sparrow) also have source habitats in both structural stages of low sagebrush. Other habitats of importance are juniper/sagebrush (Brewer's sparrow, sage sparrow, sage thrasher) and the closed herb structural stage of big sagebrush (Brewer's sparrow, lark bunting, sage sparrow, and sage thrasher). Habitats used by only a single species in the group include mountain mahogany (Brewer's sparrow), salt desert shrub (sage sparrow), and herbaceous wetlands (sage grouse).

A special habitat feature for sage grouse during the brood-rearing period is riparian vegetation, especially wet meadows with forbs (vol. 3, appendix 1, table 2). Native forbs provide spring and summer food for hens and broods (Autenrieth and others 1982, Call 1979, Oakleaf 1971, Peterson 1970, Roberson 1986, Savage 1969, Wallestad and others 1975). Herbaceous vegetation is also important to sagebrush voles (Hall 1928) and pygmy rabbits (Lyman 1991), which augment their sagebrush diet with forbs and grasses. An understory composed of native grasses is believed important for most species in group 33 (Bock and Bock 1987, Connelly and others 1991, Cooper 1868, Dobler and others 1996, Gregg 1991, Hall 1928, Mullican and Keller 1986).

Broad-scale changes in source habitats--Source habitats for group 33 were historically widespread and continuous over much of the planning area (fig. 100A), particularly in the Columbia Plateau, Northern Great Basin, Owyhee Uplands, and Upper Snake ERUs. Brewer's sparrow might have been the most

abundant bird species in the basin.

Basin-wide, nearly 48 percent of the watersheds showed a moderately or strongly declining trend in habitat, and declines exceeded increases in every ERU (fig. 101). Extensive habitat reductions were estimated in the Columbia Plateau and Upper Snake ERUs, with moderate declines in the Owyhee Uplands (figs. 100 and 101). Strongly increasing trends in habitat, however, were apparent in about 20 percent of watersheds in the Central Idaho Mountains and Columbia Plateau ERUs (fig. 101). Only the Northern Great Basin ERU has changed little from historical conditions (figs. 100 and 101).

Vol. 2, Figure 100--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 33.

Vol. 2, Figure 101--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 33, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--The single largest loss in cover types within the basin was the decline in big sagebrush (Hann and others 1997). Large-scale loss of sagebrush habitat was attributed to several factors. The first factor was conversion to agriculture. Agricultural lands have increased significantly in every ERU in the basin (Hann and others 1997). In fact, the largest transition of any terrestrial community was from upland shrubland to agriculture (Hann and others 1997). The ERUs with the biggest changes were the Columbia Plateau and Upper Snake. The former is now nearly half agricultural lands, whereas the latter is nearly one-third. These ERUs have had the greatest degree of conversion among all ERUs in the basin. Agriculture also now occupies over a tenth of the Owyhee Uplands ERU. Only the Northern Great Basin ERU has been relatively free of agricultural conversions.

A second factor contributing to loss of sagebrush habitat was conversion of shrubsteppe vegetation to exotic forbs and annual grass. Significant increases in this cover type occurred in all the major sagebrush ERUs. Exotic forbs and annual grass now occupy small portions of the Northern Great Basin, Columbia Plateau, and Owyhee Uplands ERUs, and over a tenth of the Upper Snake ERU (Hann and others 1997).

Increases in source habitats in the Central Idaho Mountains and Columbia Plateau ERUs were attributed to expansions of juniper/sagebrush and mountain mahogany cover types (Hann and others 1997).

Habitat condition for group 33 can be described by the Composite Ecological Integrity Ratings (Quigley and others 1996) that show most of the habitat to have a "low" rating. Most of the current habitat for this group was classified into Rangeland Clusters 5 (generally corresponding to much of the Owyhee Uplands ERU) and 6 (generally the Northern Great Basin, Owyhee Uplands and Upper Snake ERUs), where the primary risk to ecological integrity is continued losses of herbland and shrubland habitats (Quigley and

others 1996). Further, Rangeland Cluster 6 is vulnerable to overgrazing and to exotic grass and forb invasions (Quigley and others 1996).

Condition of special habitat features--Wet meadows and riparian vegetation, cover types used for brood-rearing by sage grouse, have declined substantially since historical times (Lee and others 1997, Quigley and others 1996).

Bare ground is an important foraging substrate for sage sparrows and sage thrashers (Rotenberry and Wiens 1980). Brewer's sparrows, however, forage mostly in sagebrush. The value of bare ground to the other bird species in this group and the sagebrush vole is unknown. Because pygmy rabbits choose tall, dense sage for their burrows and foraging sites, we assume that vegetative cover that provides protection from predators is important (Lyman 1991) and that areas of bare ground would be avoided.

Other factors affecting the group--Roading (Quigley and others 1996) has contributed to increased human disturbance in ERUs most important for sage grouse. Moderate road densities (0.4 to 1.0 km per km² [0.7 to 1.7 mi per mi²]) are typical for the Northern Great Basin ERU, the Owyhee Uplands ERU, and the Upper Snake ERU. Roads and associated human disturbance can be especially harmful to sage grouse during the lekking and wintering periods. Habitat loss caused by roads is a direct effect.

The quality of soil may be very important to the two burrowing species in this group (sagebrush vole and pygmy rabbit) because the soil must be capable of sustaining burrows. Weiss and Verts (1984) determined that burrow sites for pygmy rabbits are found in areas where soils are significantly deeper and looser than adjacent soils. Grazing, if not managed properly, has the potential to damage pygmy rabbit habitat (Washington Department of Wildlife 1993b).

Voles seldom use compacted or rocky soil (Maser and others 1974) and may be absent from areas that have suffered soil erosion as a result of heavy livestock grazing (Maser and Strickland 1978).

Heavy livestock grazing could negatively impact other species in group 33 by altering the structure and composition of the soil and removing native herbaceous understory vegetation. Thus, areas that are currently judged to be source habitat because of the presence of sagebrush cover may not be currently suitable because of changes in soil or understory vegetation that cannot be mapped at the broad scale. Additionally, changes in natural wildfire regimes have contributed to invasions of exotic vegetation in native sagebrush habitats.

Species in this group evolved in shrubsteppe habitats, where microbiotic crusts were broadly distributed (see Kaltenecker and Wicklow-Howard 1994). Microbiotic, or cryptogamic, crusts consist of lichens, bryophytes, algae, microfungi, cyanobacteria, and bacteria growing on or just below the soil surface in arid and semiarid environments (Kaltenecker and Wicklow-Howard 1994); these crusts developed without large herds of grazing ungulates (St. Clair and Johansen 1993). In addition, these crusts are projected to have been widely distributed throughout the source habitats for this group, particularly in the Northern Great Basin, Owyhee Uplands, and Upper Snake ERUs but also scattered in the Columbia Plateau ERU (Hann and others 1997, map 3.59). Increasing evidence indicates that microbiotic crusts improve soil stability, productivity, and moisture retention, moderate extreme temperatures at the soil surface, and enhance seedling establishment of vascular plants (Belnap and Gardner 1993, Harper and Pendleton 1993, Johansen and others 1993, St. Clair and others 1993), thus contributing to high ecological integrity of shrubsteppe habitats. Idaho BLM has recognized the potential importance of microbiotic crusts by proposing standards for rangeland health that include the maintenance of these crusts to ensure proper functioning and productivity of native plant communities (USDI Bureau of Land Management 1997). These crusts were widely destroyed by trampling during the excessive livestock grazing of the late 1800s and early 1900s (Daubenmire 1970, MacCracken and others 1983, Mack and Thompson 1982, Poulton 1955). Currently, high-intensity grazing and altered fire regimes modify shrubsteppe plant communities

and threaten the maintenance and recovery of microbiotic crusts (Belnap 1995, Kaltenecker 1997, St. Clair and Johansen 1993).

Little information is available on effects of landscape patterns on species in this group. Brewer's sparrows are known to have small territories, and individual pairs will occupy small patches of suitable habitat placed within a matrix of unsuitable vegetation. Sage thrashers also appear to use discontinuous, patchy habitats surrounded by other types, but rarely occur as single pairs; the probability of habitat occupancy increases with shrub patch size (Knick and Rotenberry 1995). Sage sparrows seem to be both area sensitive and more social (Rich 1981) than the previous two species. Individual pairs essentially never occur alone. The species does not occupy small patches of habitat, and large patches of seemingly suitable habitat may be unoccupied. Thus, sage sparrows occur in large expanses of shrubsteppe where many pairs share adjacent territories (Knick and Rotenberry 1995) and apparently do not use slopes of greater than a few percent.

Disjunct patches of sagebrush that were previously connected to other patches may now be unsuitable source habitat for sage grouse because wintering flocks have large home ranges. Grouse select winter use sites based on snow depth and topography (Connelly 1982, Hupp 1987, Robertson 1991) where sagebrush is accessible. Sagebrush heights of 25 to 30 cm (10 to 12 in) and canopy cover of 10 to 25 percent, regardless of snow cover, are important for winter use by sage grouse. Because seasonal movements differ among regions and populations, this effect needs to be assessed on a case-by-case basis.

Populations of pygmy rabbits historically occurred in five counties in Washington, but current records indicate that populations occur in isolated fragments in only one county (Douglas) (Washington Department of Wildlife 1993b). These small, disjunct populations are susceptible to extirpation by habitat degradation and loss, as well as catastrophic events such as fire, disease, flooding, or intense predation.

The sage sparrow, Brewer's sparrow, and lark bunting are not frequently parasitized by brown-headed cowbirds (Ehrlich and others 1988). Both sparrows apparently accept the eggs (Rich 1978). The sage thrasher also is parasitized but rejects cowbird eggs (Rich and Rothstein 1985). Sage grouse using agricultural areas may be adversely affected by pesticide applications (Blus and others 1989, Post 1951, Ward and others 1942).

Population status and trends--Quantitative population trend data are available only for the bird species in group 33. No information is available for the pygmy rabbit, only anecdotal notes are available for the sagebrush vole and, because the lark bunting is peripheral to the basin, sample sizes for this species are inadequate.

Historical reports indicate that the sagebrush vole was abundant in grasslands around Walla Walla in 1868 (Cooper 1868), although it has not been found there since. Currently, other subspecies of this vole occur in higher elevation grasslands in Utah and California where sagebrush does not occur. This suggests that the species may occur today largely in shrubsteppe habitats because the large grasslands, which it may actually prefer, no longer exist. Thus, the species probably experienced substantial population declines.

Brewer's sparrow has the most clear population trend, decreasing 1.3 percent per yr ($\bar{n} \geq 14$, $\bar{P} < 0.01$) over the period 1968-94 and 4.3 percent per yr ($\bar{n} \geq 14$, $\bar{P} < 0.01$) over the period 1984-94 (Saab and Rich 1997) in the basin. This sparrow also is declining in Idaho (6.3 percent per yr, 1966-95; $\bar{n} = 40$, $\bar{P} < 0.01$) and in physiographic region 89 (Columbia Plateau; 5.2 percent decline over the same period, $\bar{n} = 57$, $\bar{P} < 0.01$) (Sauer and others 1996). Among 15 Neotropical migrants in the basin, Brewer's sparrow, sage sparrow, sage thrasher, and lark bunting were designated as species of high concern to management under all future management themes for the basin (Saab and Rich 1997).

Population trends for the sage sparrow and sage thrasher are not consistent with the population declines demonstrated by Brewer's sparrows and sage grouse. The sage sparrow shows no trend in the basin (Saab and Rich 1997) and a nonsignificant decline of -1.0 percent per yr (1966 to 1995, $\bar{n} = 38$) in physiographic region 89 (Columbia Plateau; Sauer and others 1996). The sage thrasher also shows no trend in the basin (Saab and Rich 1997), a nonsignificant 1.1-percent per yr decline in Idaho ($\bar{n} = 28$), a 2.1-percent per yr increase in Oregon ($\bar{n} = 27$, $P < 0.01$), and a nonsignificant 0.8-percent per yr increase in physiographic region 89 (Columbia Plateau; $\bar{n} = 51$) over the period 1966-95 (Sauer and others 1996). Sage grouse populations have shown significant, steep declines since the 1940s in Idaho,¹³ Oregon (Crawford and Lutz 1985), and Washington (Tirhi 1995). The rates of decline in Idaho and Oregon are not significantly different.¹⁴ Moreover, the rate of decline in Washington appears to be similar to that in Idaho and Oregon, thereby suggesting common, widespread factors affecting these populations. A complicating factor is that sage grouse in this geographic area may exhibit population cycles with a periodicity of around 10 years (Rich 1985, Willis and others 1993b). Thus, apparent trends over short periods should be regarded with caution. Populations in Washington were heavily impacted by habitat loss before surveys were established. Remaining populations now exist as isolated remnants (Tirhi 1995).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 33 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues---The results of our habitat trend analysis suggest the following issues are of high priority for group 33:

Primary issues are as follows:

1. Permanent and continued loss of large areas of shrubsteppe habitat to agricultural conversion, brush control, and cheatgrass invasion.
2. Soil compaction, erosion, and loss of microbiotic crust.
3. Continued degradation of wet meadow and riparian vegetation adjacent to springs, seeps, and streams by improper grazing and, in some areas, spring development to provide livestock water supplies.
4. Adverse effects of human disturbance.

Potential strategies--The following strategies could be used to reverse broad-scale declines in source habitats:

1. (To address issue no. 1) Identify and conserve large remaining areas of shrubsteppe vegetation where ecological integrity is still relatively high (Gray and Rickard 1989, Rickard and Poole 1989, Schuler and others 1993, Smith 1994, Yoakum 1980). Basin-wide, maintain or restore 15 to 25 percent sagebrush cover with heights of 36 to 79 cm (14 to 31 in) (Autenrieth 1981, Connelly and others 1991, Fischer 1994, Gregg 1991, Klebenow 1969, Patterson 1952, Peterson 1970, Wakkinen 1990, Wallestad 1975). In sage grouse winter range, maintain a mosaic of sagebrush height and cover classes to allow access to sagebrush with canopy cover of 10 to 25 percent and heights of 25 to 30 cm (10 in to 12 in) regardless of snow cover (Connelly 1982, Hupp 1987, Robertson 1991).
2. (To address issue no. 1) Restore native grass and forb understories to historical levels, where

restoration potential exists, and retard the spread of nonnative vegetation.

3. (To address issue no. 2) Reduce and eliminate soil compaction and erosion to benefit both pygmy rabbits and sagebrush voles.
4. (To address issue no. 2) Restore microbiotic crusts in ERUs with potential for redevelopment (that is, areas near propogule sources, and with suitable soil, vegetation, and climatic characteristics [see Belnap 1993, Belnap 1995, Kaltenecker 1997, Kaltenecker and Wicklow-Howard 1994]): the Northern Great Basin, Owyhee Uplands, Upper Snake, and, to a lesser extent, the Columbia Plateau (Hann and others, map 3.59).
5. (To address issue no. 3) Restore vegetation around springs, seeps, streams, meadows, and other riparian areas.
6. (To address issue no. 4) Minimize the adverse effects of human disturbance.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Identify sites of high ecological integrity to be managed for long-term protection by analyzing current vegetation, precipitation patterns, elevation, temperature (Klemmedson and Smith 1964, Morrow and Stahlman 1984, Stewart and Hull 1949) and the presence of priority species in this group. These practices are most likely to be successful on large areas of Federal land managed by the BLM.
2. (In support of strategy no. 1) Explore options under the CRP (Johnson and Igl 1995), or develop other incentive programs, to encourage restoration of agricultural areas to native cover types. Focus on areas that would increase patch size or links with existing source habitat patches.
3. (In support of strategy no. 1) Avoid further loss of sagebrush cover through burning, plowing, seeding, and other brush "control" methods where sagebrush cover types are below historical levels.
4. (In support of strategy no. 1) Avoid further conversion of source habitats to agricultural lands, or strive to minimize the impacts of further conversions through landscape design, to minimize further fragmentation of shrubsteppe.
5. (In support of strategy no. 2) Use fire prevention and suppression to retard the spread of cheatgrass in areas that are susceptible to cheatgrass invasion but currently are dominated by native grass species. Planting of fire-resistant vegetation through "green stripping" (Pellant 1994) could be explored to evaluate its effectiveness in protecting existing native vegetation.
6. (In support of strategy no. 2) Restore selected areas of cheatgrass monocultures through seeding and other manipulations (Allen 1995, Daubenmire 1970, Evans and Young 1978, Hosten and West 1994, Kennedy 1994, Monsen and McArthur 1995, Ogg 1994, Whisenant 1995, Yoakum 1986b), where such restoration would increase the size of existing shrubsteppe patches or provide links among patches.
7. (In support of strategy no. 2) Plant perennial bunchgrasses or native forbs where these components of the habitat have been lost or reduced (Braun and others 1976, Daubenmire 1970, Evans and Young 1978, Yoakum 1986b). Criteria for enhancement include maintaining or increasing the size of smaller patches, preventing further habitat disassociation, and protecting or increasing the size and integrity of corridors among patches, all in connection with the location of sites with high ecological integrity as identified above.

8. (In support of strategy nos. 2, 3, and 4) Modify grazing systems or reduce grazing use where native perennial bunchgrasses have been depleted.
9. (In support of strategy no. 4) Encourage the redevelopment of microbiotic crust by reducing or eliminating livestock grazing (Mack and Thompson 1982, St. Clair and others 1993). Explore the use of ground-based and aerial soil inoculation to increase the speed and extent of dispersal of the organisms that create microbiotic crust (Belnap 1993, 1994).
10. (In support of strategy no. 5) Protect existing riparian, spring, and seep sites of high ecological integrity from degradation, restore degraded sites, restore historical water tables in nonfunctioning riparian systems, and eliminate or greatly reduce water diversions. Seeding of native forbs, in particular, may be desirable in certain mesic areas to improve sage grouse brood-rearing habitat.
11. (In support of strategy no. 6) Protect sage grouse leks from human disturbance by designating leks and winter concentration sites as special management areas closed to public access, avoiding the placement of new roads or the improvement of existing roads in important sage grouse areas, and closing existing roads in sensitive areas.
12. (In support of strategy no. 6) Control, reduce, or eliminate pesticide use around agricultural areas adjacent to sage grouse habitat (Blus and others 1989, Post 1951, Ward and others 1942). Avoid use of toxic organophosphorus and carbamate insecticides in sage grouse brood-rearing habitats.
13. (In support of strategy no. 6) Restrict organized recreational events in sage grouse nesting, brood-rearing, and wintering habitats at the appropriate times of year (Call 1979, Roberson 1986).

GROUP 34--KIT FOX AND BLACK-THROATED SPARROW

Results

Species ranges, source habitats, and special habitat features--Group 34 consists of two shrubland species, the kit fox and black-throated sparrow. Both species occur in the most southern shrublands of the basin, and the black-throated sparrow also is found in south-central Washington (fig. 102). The kit fox is a year-round resident of the basin, whereas the black-throated sparrow is a summer resident, migrating to southern portions of its range and Baja California for the winter. The basin represents the northern periphery of the continental distribution for these species, both of which are more commonly associated with desert shrublands of southwestern North America.

Vol. 2, Figure 102--Ranges of species in group 34 within the basin.

Source habitats for both species are big sagebrush and salt desert shrub, and the black-throated sparrow also uses mountain big sagebrush (vol. 3, appendix 1, table 1). Structural stages within these cover types are open- and closed-canopy stages of low-medium shrubs. In southeastern Oregon and northern Nevada, black-throated sparrows are found predominantly in sites with higher shrub cover, greater maximum shrub height, and greater shrub species diversity than used by another shrubsteppe species, the sage sparrow (Wiens and Rotenberry 1981).

A special habitat feature identified for the kit fox is the presence of burrows for den sites (vol. 3, appendix 1, table 2). Kit foxes often use the abandoned dens of other species, and most home ranges include several dens (Egoscue 1962). In addition to reproductive purposes, dens provide resting habitat that modifies the extremes of desert weather and furnishes protection from predators (Golightly and Ohmart 1983). No special habitat features have been identified for the black-throated sparrow.

Broad-scale changes in source habitats--Source habitats have undergone localized declines since historical times. Historically, source habitats were concentrated along the southeastern border of Oregon and southern border of Idaho, extending also into the portions of Nevada and Utah that are included in the basin (fig. 103A). Source habitats for the black-throated sparrow also occurred in south-central Washington. The current distribution of source habitats is roughly the same, but declines in habitat availability have occurred primarily in south-central Washington and south-central Idaho (fig. 103B).

The amount of source habitats was estimated as roughly the same as the historical extent in 65 percent of the watersheds in which these species occur, but 33 percent of the watersheds have exhibited declining trends (fig. 104). The greatest declines occurred in the Upper Snake ERU, where 29 of 55 watersheds had strongly declining trends (fig. 104). The Blue Mountains and Snake Headwaters ERUs also had strongly declining trends, but only three watersheds in each ERU provided source habitats historically, so the magnitude of change may not be significant. Habitat trends were mostly static in the Owyhee Uplands ERU, although 82 of the 256 watersheds with source habitats have declining trends (fig. 104).

Vol. 2, Figure 103--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 34.

Vol. 2, Figure 104--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 34, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--The principal cause for observed declines in habitat availability is the alteration of sagebrush and salt desert shrub to other cover types, primarily agriculture, urban, juniper/sagebrush, and exotic forbs-annual grass. In the Columbia Plateau ERU, nearly one half of the big sagebrush cover type was converted to croplands (Hann and others 1997). Virtually all broad-scale patches of mountain big sagebrush in the Columbia Plateau within the range of the black-throated sparrow were eliminated (vol. 3, appendix 1, table 4). In the Owyhee Uplands, the dominant cover type transition was from the big sagebrush cover type to croplands and exotic forbs-annual grass (Hann and others 1997). In the Upper Snake ERU, an estimated 41 percent of the sagebrush cover type was converted to croplands (Hann and others 1997).

Condition of special habitat features--No information is available to determine whether changes in availability of burrows for kit fox dens, or in soil conditions needed for burrow excavation, have occurred in the basin. Lack of suitable loose-textured soil for burrow construction may be a natural, limiting factor

for kit foxes in southeastern Oregon (Keister and Immell 1994). The soil surface there is predominantly desert pavement, whereas soils near Fallon, Nevada, where higher densities of kit foxes occur than in Oregon, are typically sandy (Keister and Immell 1994). Land uses that increase soil compaction or cause the destabilization of dunes may inhibit burrow establishment.

Other factors affecting the group--The black-throated sparrow seems to show a positive numerical response to moderate livestock grazing (Bock and others 1984, cited in Saab and others 1995).

Because the kit fox is a predator, population health is affected by the availability of small-mammal prey, which in turn is affected by vegetation composition and structure. Land uses that do not directly affect kit foxes may nevertheless affect prey availability. Livestock grazing can impact small-mammal abundance and diversity (Bock and others 1984; Hanley and Page 1982, as cited in Horning 1994).

Kit foxes are vulnerable to poisoned baits placed for destruction of coyotes (Orloff and others 1986). They are also susceptible to hunting and trapping, usually as a nontarget species (DeStefano 1990). Coyote predation is a major cause of kit fox mortality in the San Joaquin Valley of California (White and others 1994), and is a potential limiting factor of kit foxes in the basin.

Population status and trends--Population trend data are not available for the black-throated sparrow within the basin. The only statistically significant population trend for the black-throated sparrow is based on numbers recorded on all BBS routes in North America with black-throated sparrow occurrences between 1966 and 1995. This survey-wide trend indicated a 4-percent annual decline across the range of the species over the 28-yr period ($\bar{n} = 258$, $P < 0.05$; Sauer and others 1996). Occurrences of the black-throated sparrow on BBS routes within the basin are insufficient to conduct a statistically robust trend analysis (Saab and Rich 1997). Saab and Rich (1997), however, included the black-throated sparrow as one of 15 Neotropical migrants in the basin that are of high concern to management under all future management themes for the basin primarily because of its' association with just four cover type-structural stage combinations. We know of no estimates of kit fox numbers within the basin.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 34 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--Primary conservation issues for group 34 are as follows:

1. Loss of desert shrub habitat to other land uses and to shrub control programs.
2. Degradation of desert shrub habitat quality through exotic weed invasions.
3. Effect of adverse land uses on understory vegetation that supports kit fox prey base.
4. Lack of information on the location and status of kit fox dens.

Potential strategies--strategies for addressing the issues listed above include the following:

1. (To address issue no. 1) Maintain remaining native desert shrublands, especially in the Upper Snake ERU and in all watersheds within the Owyhee Uplands where strong negative trends have occurred.
2. (To address issue no. 2) Restore desired vegetation composition and structural attributes of

shrublands that no longer meet source habitat conditions.

3. (To address issue no. 3) Avoid land use practices that potentially affect kit fox prey by reducing the grass-forb component of shrub communities.
4. (To address issue no. 4) Locate and protect active dens of the kit fox.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (To address strategy no. 1) Identify areas of native shrubland vegetation with high ecological integrity, particularly within the Columbia Plateau and Upper Snake ERUs, and actively manage to promote their long-term sustainability.
2. (To address strategy no. 2) Use prescribed burns, shrub planting, and exotic weed control to restore degraded shrublands, but avoid burning areas susceptible to noxious weed invasion.
3. (To address strategy no. 3) Adjust or maintain grazing management plans to promote long-term persistence of the grass and forb components of shrub communities.
4. (To address strategy no. 4) Conduct surveys for kit fox burrows, and provide protective measures for active burrows in all relevant planning documents.

GROUP 35--LOGGERHEAD SHRIKE

Results

Species ranges and source habitats--Group 35 consists of breeding habitat for the loggerhead shrike. Range of the loggerhead shrike (fig. 105) includes most of the basin except for the mountainous portions of Idaho and Montana and the eastern slope of the Cascade Range. Outside the planning area, the species is widespread as a breeder or year-round resident in the United States and Mexico (Yosef 1996).

Vol. 2, Figure 105--Ranges of species in group 35 within the basin.

This shrike uses various woodland and shrub cover types including juniper, sagebrush, mountain shrub types, salt desert shrubs, and bitterbrush/wheatgrass (vol. 3, appendix 1, table 1). The common structural feature is a good component of woody vegetation in a landscape dominated by more open structure. Nests are typically placed in the taller woody vegetation, whereas the bird forages in open areas.

Broad-scale changes in source habitats--High percentages of contiguous watersheds with source habitats for the loggerhead shrike historically occurred in the Columbia Plateau, Northern Great Basin, Owyhee Uplands, and Upper Snake ERUs (fig. 106A).

Basin-wide, moderate and strong declines (44 percent of watersheds) in source habitats exceeded moderate and strong increases (24 percent), but over 30 percent of watersheds showed no estimated

change from the historical condition (fig. 107). Although declining trends in the Columbia Plateau seem to balance against increasing trends (fig. 107), these upward trends were due to large relative increases in vegetation that actually covered <8 percent of the ERU. The biggest losses occurred in the Upper Snake ERU (fig. 107), with over 57 percent of the watersheds showing strong decreases. In contrast, the Upper Klamath ERU was estimated to have nearly 62 percent of its watersheds strongly increasing in source habitats (fig. 107).

Vol. 2, Figure 106--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 35.

Vol. 2, Figure 107--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 35, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Among source habitats basin-wide, big sagebrush types have declined by one-third, the most serious habitat change for shrikes because of the total acreage affected (vol. 3, appendix 1, table 4; Hann and others 1997). Salt desert shrub and mixed-conifer woodlands also have declined substantially, one-third and one-half, respectively. Together, the latter declines affected only a small part of the basin (vol. 3, appendix 1, table 4; Hann and others 1997). The only other significant basin-wide changes have been increases in juniper/sagebrush, juniper woodlands, and mountain mahogany (Hann and others 1997). The latter three types combined, however, cover only a small percentage of the basin.

The largest changes have been in the Upper Snake and Columbia Plateau ERUs, where big sagebrush has declined by about 50 percent (vol. 3, appendix 1, table 4). In the Upper Clark Fork and Blue Mountains ERUs, mixed-conifer woodlands have declined by over four-fifths and one-half, respectively (Hann and others 1997). Declines in the Upper Clark Fork can be attributed to a near total loss of mixed-conifer woodlands, although this type historically only covered a small portion of the ERU. Increases in the southern Columbia Plateau are due to juniper/sagebrush, which more than doubled, and mountain mahogany, up nearly sixfold; these types together now are estimated to occupy nearly one-tenth of the ERU. Similarly, juniper/sagebrush in the Upper Klamath is estimated to have tripled, making the availability of source habitats there significantly greater (Hann and others 1997). Large increases in source habitats in the Northern Glaciated Mountains are most likely because of relatively large increases in mixed-conifer woodlands, though source habitat is limited in this ERU.

Large-scale loss of sagebrush habitats is due to several factors. The first factor is conversion to agriculture. Agricultural lands have increased significantly in every ERU in the basin (Hann and others 1997). In fact, the largest transition of any terrestrial community from historical to the current period was that of upland shrubland to agriculture (+9.0 percent), and the second largest was that from upland herbland to agriculture (+6.6 percent, Hann and others 1997). This transition, occurring in the fundamental source habitats for this group, explains much of the pattern evident in habitat trends for loggerhead shrike (fig. 106).

A second factor contributing to loss of sagebrush habitat is conversion of shrubsteppe vegetation to exotic forbs and annual grass. Increases in exotic cover types have occurred in all the major shrubsteppe ERUs. Substantial portions of the Owyhee Uplands and Upper Snake ERUs have undergone a conversion from upland shrubland to exotic herbland (Hann and others 1997).

The condition of the habitat for group 35 can be described by the Composite Ecological Integrity Ratings (Quigley and others 1996) that show most of the habitat to have a "low" rating. Most of the current habitat for this group was classified into Rangeland Clusters 5 (generally corresponding to much of the Owyhee Uplands ERU) and 6 (generally the Northern Great Basin, Owyhee Uplands and Upper Snake ERUs), where the primary risk to ecological integrity is continued losses of herbland and shrubland habitats (Quigley and others 1996). Further, Rangeland Cluster 6 also is vulnerable to overgrazing and to exotic grass and forb invasions (Quigley and others 1996).

Other factors affecting the group--Shrikes prefer tall plants for nest sites, often choosing particularly tall individual big sagebrush plants or, more generally, sites with tall average shrub heights (for example, >1 m [3 ft]) (Leu 1995, Sharp 1992, Yosef 1996). This type of sagebrush community is apt to be a big sagebrush site with deeper soils and a slightly more mesic moisture regime. These sites are precisely where agricultural conversion has most commonly occurred in the past and where future risks of conversion remain the greatest (Hann and others 1997).

Shrikes also prefer to hunt from elevated perches such as fence posts, utility lines, and woody vegetation (Bohall-Wood 1987, Gawlik and Bildstein 1993, Yosef and Grubb 1992), and to restrict their foraging to an area within 10 m of such perches (Chavez-Ramirez and others 1994). Their use of any area may correspond directly to the availability of such perches. Young shrikes prefer to forage on bare ground and sites with little vegetative cover (Leu 1995). Foraging opportunities for young shrikes may be severely reduced because shrubsteppe habitats with natural opening of bare ground have been altered by exotic grasses (for example, cheatgrass) and forbs, creating a continuous vegetative layer (see Leu 1995).

In a study area generally corresponding to the Northern Great Basin ERU, shrike densities were negatively correlated with the cover of grass and positively correlated with woody cover, bare ground, and vegetation height (Rotenberry and Wiens 1980). Shrike densities were negatively correlated with those of Brewer's sparrow and positively correlated with those of rock wrens. Among habitat variables, shrikes were positively associated with the cover of rock and shrubs, and with shrub species diversity (Wiens and Rotenberry 1981).

Species in this group evolved in shrubsteppe habitats, where microbiotic crusts were broadly distributed (see Kaltenecker and Wicklow-Howard 1994). Microbiotic, or cryptogamic, crusts consist of lichens, bryophytes, algae, microfungi, cyanobacteria, and bacteria growing on or just below the soil surface in arid and semiarid environments (Kaltenecker and Wicklow-Howard 1994); these crusts developed without large herds of grazing ungulates (St. Clair and Johansen 1993). In addition, these crusts were projected to have been widely distributed throughout the source habitats for this group, particularly in the Northern Great Basin, Owyhee Uplands, and Upper Snake ERUs, but also scattered in the Columbia Plateau ERU (Hann and others 1997, map 3.59). Increasing evidence indicates that microbiotic crusts improve soil stability, productivity, and moisture retention, moderate extreme temperatures at the soil surface, and enhance seedling establishment of vascular plants (Belnap and Gardner 1993, Harper and Pendleton 1993, Johansen and others 1993, St. Clair and others 1993), thus contributing to high ecological integrity of shrubsteppe habitats. Idaho BLM has recognized the potential importance of microbiotic crusts by proposing standards for rangeland health that include the maintenance of these crusts to ensure proper functioning and productivity of native plant communities (USDI Bureau of Land Management 1997). These crusts were widely destroyed by trampling during the excessive livestock grazing of the late 1800s and early 1900s (Daubenmire 1970, MacCracken and others 1983, Mack and Thompson 1982, Poulton

1955). Currently, high-intensity grazing and altered fire regimes modify shrubsteppe plant communities and threaten the maintenance and recovery of microbiotic crusts (Belnap 1995, Kaltenecker 1997, St. Clair and Johansen 1993).

Conversion of native vegetation to exotics is augmented by the propensity of annuals, such as cheatgrass, to spread with wildfire and with improper grazing (Braun and others 1976; Daubenmire 1970; Evans and Young 1978; Quigley and others 1996, p. 123). Some losses of salt desert shrubs are likely due to selective grazing of palatable forbs in this cover type, combined with more xeric conditions that make vegetative resilience low.

Losses of pasture and old fields for wintering habitat in the Southeastern United States have affected shrike populations (Brooks and Temple 1990, Gawlik and Bildstein 1993). Loss of pasture and prairie habitats for breeding in Canada and the Eastern United States are widely cited as causes for population declines in those regions (Yosef 1996). These habitat losses have not been identified as limiting factors for shrike populations in the basin.

Because shrikes often forage and nest along roads (Blumton 1989, Craig 1978, Flickinger 1995, Yosef 1996), vehicular collisions may be an important source of mortality. Automobiles accounted for 29 percent of the observed fall and winter mortality of loggerhead shrikes in Virginia (Blumton 1989). Shrikes also may have been affected by DDT in the past and may suffer sublethal effects of certain insecticides, although the evidence is weak (Anderson and Duzan 1978, Grubb and Yosef 1994, Yosef 1996). Cowbird parasitism of nests does not appear to be a factor affecting productivity of loggerhead shrikes (Yosef 1996).

Population status and trends--Populations of loggerhead shrikes have been declining significantly in the basin, with a trend of -2.7 percent per yr ($\bar{n} \geq 14$, $P < 0.05$) over the period 1968-94 (Saab and Rich 1997). The 1966-95 trend for BBS physiographic region 89 (Columbia Plateau) was -2.3 percent per yr ($\bar{n} = 41$, $P < 0.05$; Sauer and others 1996). Saab and Rich (1997) included the loggerhead shrike as one of 15 Neotropical migrants in the basin that are of high concern to management under all future management themes for the basin.

Patterns of widespread declines throughout its range (Yosef 1996) suggest that either (1) habitat losses throughout its breeding range in various types of breeding habitat are similar, or (2) additional, more extensive factors are impacting the species, such as pesticides or wintering ground problems. These possibilities do not diminish the losses of source habitats in the basin but suggest that widespread population declines may be at least partly the result of a more pervasive cause.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 35 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues---The results of our habitat trend analysis suggest the following issues are of high-priority for group 35:

1. Permanent and continued loss of large acreage of big sagebrush cover types to agricultural conversion, brush control, reduction of microbiotic crusts, and cheatgrass invasion.
2. Adverse effects of human disturbance.

Potential strategies--The following strategies could be used to reverse broad-scale declines in source habitats:

1. (To address issue no. 1) Identify and conserve large remaining areas (contiguous habitat >1000 ha [2,470 acres]) of shrubsteppe vegetation where ecological integrity is still relatively high (Gray and Rickard 1989, Rickard and Poole 1989, Schuler and others 1993, Smith 1994, Yoakum 1980). Sites resistant to cheatgrass domination because of their moisture regime (>30 cm [12 in]) in the Upper Snake, Owyhee Uplands, Northern Great Basin, and Columbia Plateau ERUs are of highest priority.
2. (To address issue no. 1) Restore microbiotic crusts in ERUs with potential for redevelopment (that is, areas near propagule sources, and with suitable soil, vegetation, and climatic characteristics [see Belnap 1993, Belnap 1995, Kaltenecker 1997, Kaltenecker and Wicklow-Howard 1994]): the Northern Great Basin, Owyhee Uplands, Upper Snake, and, to a lesser extent, the Columbia Plateau (Hann and others, map 3.59).
3. (To address issue no. 1) Retard the spread of cheatgrass in native shrubsteppe vegetation communities.
4. (To address issue no. 2) Minimize adverse effects of human disturbance.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Protect and restore corridors and habitat blocks in areas of shrubsteppe that support large, contiguous areas of high ecological integrity so as to optimize long-term conservation of shrikes. These practices are most likely to be successful on large tracts of Federal land managed by BLM.
2. (In support of strategy no. 1) Restore existing agricultural lands to native vegetation when possible. Sites where this might be especially useful are areas that were historically shrubsteppe and areas that would augment corridors among existing shrubsteppe patches.
3. (In support of strategy no. 1) Avoid further loss of sagebrush cover through burning, plowing, seeding, and other brush "control" methods where sagebrush cover types are well below historical levels.
4. (In support of strategy no. 1) Minimize the impacts of further agricultural conversions through landscape design. If conversion cannot be avoided, then tracts slated for conversion should be located to minimize further disassociation of shrubsteppe, to avoid reducing the size of isolated patches, and to avoid areas that are currently in large blocks of moderate Composite Ecological Integrity (Quigley and others 1996).
5. (In support of strategy no. 1) Restore native vegetation by appropriate mechanical treatments and seedings of native shrub, grass, and forb species (Allen 1995, Monsen and McArthur 1995, Whisenant 1995, Yoakum 1986b).
6. (In support of strategy no. 2) Encourage the redevelopment of microbiotic crust by reducing or eliminating livestock grazing (Mack and Thompson 1982, St. Clair and others 1993). Explore the use of ground-based and aerial soil inoculation to increase the speed and extent of dispersal of the organisms that create microbiotic crust (Belnap 1993, 1994).
7. (In support of strategy no. 3) Use fire prevention and suppression to retard the spread of cheatgrass in areas that are susceptible to cheatgrass invasion but currently are dominated by native grass

species. Explore the effectiveness of planting fire-resistant vegetation through "green stripping" (Pellant 1994) to protect existing vegetation as well as allow degraded sites a chance to recover.

8. (In support of strategy no. 3) Restore selected areas of cheatgrass monocultures through seeding and other manipulations (Allen 1995, Daubenmire 1970, Evans and Young 1978, Hosten and West 1994, Kennedy 1994, Monsen and McArthur 1995, Ogg 1994, Whisenant 1995, Yoakum 1986b), where such restoration would increase the size of existing shrubsteppe patches or provide links among patches.
9. (In support of strategy no. 4) Minimize access to roads and, where possible, locate them on the periphery of areas known to have good shrike populations. Avoid construction of new roads or improvement of old roads in shrike habitat. Plan habitat enhancement projects for sites away from heavily traveled roads.
10. (In support of strategy no. 4) Avoid insecticide spraying during shrike breeding season.

GROUP 36 --COLUMBIAN SHARP-TAILED GROUSE (SUMMER)

Results

Species ranges, source habitats, and special habitat features--Columbian sharp-tailed grouse is a year-round resident that is distributed patchily in mesic shrubland and grassland types of the Upper Snake, Snake Headwaters, Central Idaho Mountains, Northern Glaciated Mountains, and Columbia Plateau ERUs (fig. 108). Only trends in summer habitat are evaluated here, because winter cover types (primarily riparian and upland shrub) occur in naturally small patches that could not be analyzed at the broad scale. During the late 1980s, early 1990s, 1996, and 1997, populations were augmented in Montana within the Northern Glaciated Mountains ERU and reintroduced in Oregon within the Blue Mountains ERU.

Vol. 2, Figure 108--Ranges of species in group 36 within the basin.

Summer source habitats of Columbian sharp-tailed grouse include open-canopied big, mountain, and low sagebrush cover types, wheatgrass and fescue bunchgrasses, herbaceous wetlands, upland or mountain shrub cover types of chokecherry-serviceberry-rose, and shrub wetland cover types (vol. 3, appendix 1, table 1) (Marks and Saab Marks 1987a, Meints and others 1992, Saab and Marks 1992). Within these habitats, sharptails only use areas where the annual precipitation is at least 30 cm (12 in) (Meints and others 1992), and where the topography is flat to rolling (<30 percent slope) (Saab and Marks 1992). During spring and summer, sagebrush and grasslands provide nesting and brood-rearing habitat, whereas mountain (upland shrub) and riparian shrubs are used for escape cover. Fall and winter habitats are primarily mountain shrub and riparian vegetation. Following those seasonal changes in habitat use, herbaceous vegetation and associated arthropods provide food for sharptails during spring and summer, whereas fruits and buds of woody vegetation, insects, and agricultural crops are consumed by grouse during fall and winter (Giesen and Connelly 1993).

During spring and summer in western Idaho, nesting and brood-rearing microhabitats used by sharptails are characterized by moderate vegetative cover (≥ 60 percent), high structural diversity, and a high diversity of native herbaceous vegetation (Marks and Saab Marks 1987a, Saab and Marks 1992). Native

perennials arrowleaf balsamroot and bluebunch wheatgrass were especially important nesting and brood-rearing cover during a drought year when many exotic annuals dried up and provided no cover (Saab and Marks 1992). Additionally, selected microhabitats in western Idaho were least modified by livestock grazing and near escape cover of mountain shrubs and riparian vegetation. Grouse broods in eastern Idaho preferred CRP lands over native shrublands or agricultural fields during summer (Sirotnak and others 1991). Seedlings on CRP lands provide nesting cover and are often good sources of food if the seedlings include alfalfa, *Tragopogon* species, and *Lactuca* species. Height of nest-brood cover was identified as a critical microhabitat feature and averaged 25 ± 16 cm (10 ± 6.3 in) in eastern Idaho (Meints and others 1992).

When native shrubland is used for nesting in Idaho, most nests are placed beneath a shrub (Marks and Saab Marks 1987a, Meints 1991). Thus, shrubs are a special habitat feature for this species (vol. 3, appendix 1, table 2). Shrub density at nests in eastern Idaho averaged 11,000 shrubs per ha (2.5 acres) compared to 5,000 shrubs per ha (2.5 acres) at independent, randomly located sites (Meints 1991). In a native grassland of northwestern Montana, preliminary data indicated that nests were placed beneath wheatgrass and fescue bunchgrasses.¹⁵

Spring and summer movements are typically within 1.0 to 2.5 km (0.63 to 1.6 mi) of dancing grounds (lek sites) (Saab and Marks 1992). Summer home ranges averaged 187 ± 114 ha (462 ± 282 acres) in western Idaho and 90 percent of all locations were within 1.2 km (0.75 mi) of a dancing ground (lek site) (Saab and Marks 1992). Nests have been located <100 m (328 ft) (Marks and Saab Marks 1987a) to >3 km (1.9 mi) (Meints 1991) from lek sites, with most females nesting ≤ 1.6 km (1.0 mi) from the lek where they were trapped (Marks and Saab Marks 1987a, Meints 1991, Oedekoven 1985).

Winter habitat requirements seem more restricted than in other seasons (Giesen and Connelly 1993). Columbian sharp-tails in western Idaho wintered almost exclusively in mountain shrub or riparian cover types, the only cover types that provided food and escape cover regardless of snow depth (Marks and Saab Marks 1988). Fruits of Douglas hawthorn and buds of serviceberry and chokecherry were the main winter foods. Winter grouse locations in eastern Idaho averaged 90 m (295 ft) to riparian cover (Meints 1991). Movements of sharp-tails between breeding and wintering areas varied from 2.6 km (1.6 mi) in western Idaho (Marks and Saab Marks 1987a) to 20 km (12.5 mi) in southeastern Idaho (Meints 1991). Columbian sharp-tails apparently move farther to wintering habitats in regions lacking a broad distribution of winter food resources (Giesen and Connelly 1993).

Broad-scale change in source habitats--Historically, source habitats for Columbian sharp-tailed grouse were broadly distributed in eastern Washington and eastern Oregon, except in the Northern and Southern Cascades ERUs (fig. 109A). Historical source habitats were also in western portions of the Central Idaho Mountains, in the southern Owyhee Uplands, southern Snake Headwaters, and eastern portions of the Upper Snake and Snake Headwaters ERUs (fig. 109A).

The current distribution of source habitats is extremely limited and highly disjunct compared to historical patterns (fig. 109B). The western half of the Snake Headwaters and eastern Upper Snake ERUs currently provide the most contiguous habitat within the current range (figs. 108, 109B). In contrast, other remaining populations are restricted to small and isolated portions of the Central Idaho Mountains, Northern Glaciated Mountains, Columbia Plateau, Blue Mountains, and Lower Clark Fork ERUs (fig. 109B). Breeding populations reintroduced to northeastern Oregon in the early 1990s occupy small areas near Enterprise in the Blue Mountains, and augmentations were conducted near Eureka, Montana, in the Northern Glaciated Mountains during the late 1980s and early 1990s (fig. 108).

Strong declines in source habitats were projected in over 60 percent of watersheds throughout the basin, whereas increases in habitat occurred in only 6 percent of watersheds (figs. 109C and 110). Eight of 11 ERUs with historical source habitats had strongly decreasing trends. The Northern Glaciated Mountains

experienced the greatest declines, where 94 percent of the watersheds had strong decreases in source habitats (fig. 110).

Vol. 2, Figure 109--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 36.

Vol. 2, Figure 110--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 36, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--The open-canopy low-medium structural stage of mountain big sagebrush and big sagebrush experienced some of the greatest absolute declines on an ERU basis. The combined absolute decline for the open-canopy low-medium structural stage of these two sagebrush types declined in the Upper Snake (-40 percent), Owyhee Uplands (-20 percent), Columbia Plateau (-13 percent), Snake Headwaters (-7 percent), and Northern Great Basin (-2 percent) (vol. 3 appendix 1, table 4). In these open-canopied cover types, in the absence of fire, shrubs and trees eventually invade much of the area that was occupied by grasses and forbs.

In addition, large-scale losses of sagebrush habitats were attributed primarily to agricultural development. Agricultural lands have increased substantially in all ERUs within the basin (vol. 3, appendix 1, table 4). The largest conversions of terrestrial communities from historical to current levels were those of upland shrubland to agriculture and from upland herbland to agriculture (Hann and others 1997). These conversions were widespread within the historical range of sharptails and, in part, explained the broad-scale changes in their source habitats (fig. 109C).

Mountain shrub (chokecherry-serviceberry-rose) and shrub wetland terrestrial community groups are key components of sharp-tailed grouse habitat during late summer, fall, and winter. These cover types naturally occur in small patches and were difficult to map at the scale of this analysis. Therefore, accurate information was not available on habitat trends in mountain shrub and shrub wetlands.

Condition of special habitat features--Mesic sagebrush lands, mountain shrub (chokecherry-serviceberry-rose) communities, and riparian vegetation are special habitat features used by sharptails. Loss and degradation of these features, as a result of livestock grazing and agricultural conversions, were identified as factors contributing to the widespread population declines in Columbian sharp-tailed grouse within the basin (Marks and Saab Marks 1987a, 1988; Meints and others 1992; Saab and Marks 1992; Tirhi 1995). Additionally, losses of native perennial grasses and forb understories of the mesic sagebrush zones, because of livestock grazing and exotic grass invasions, are microhabitat features that could not be examined by the broad-scale analysis.

Other factors affecting the group--Livestock grazing is the dominant land use in occupied Columbian sharp-tailed grouse habitat. Habitat degradation by high-intensity livestock grazing (also by native

ungulates) results in reductions or losses of native perennial grasses and forbs, necessary for grouse nesting and brood-rearing cover. Excessive grazing can alter the native vegetation by allowing invasions of exotic plants, including cheatgrass, medusahead, and mustards. Additionally, deciduous trees and shrubs, which are critical for sharptail escape cover and for winter food (Marks and Saab Marks 1987a, 1987b, 1988; Meints 1991; Tirhi 1995), may be reduced by intensive cattle browsing during late summer (Kovalchik and Elmore 1992).

Loss of lands managed under the CRP is potentially another factor influencing Columbian sharptails. In eastern Idaho, CRP lands provide important feeding, nesting, brood-rearing, and relatively mild winter habitat (Ulliman 1995). In Washington, however, CRP lands receive little use by sharptails (Schroeder 1994). Although some females nest in CRP and other idle croplands, the most successful nests in Washington were built in native habitats of sagebrush or forbs mixed with grass (Schroeder 1994).

Herbicides and pesticides have been identified as potential threats to sharptails (Giesen and Connelly 1993). Herbicide spraying has negative effects on the species because of losses in herbaceous and woody vegetation that is used for nesting, brood-rearing, and wintering habitat. Pesticide spraying may have negative impacts by directly killing young or by reducing or eliminating insects used for food.

Fire can either enhance or degrade sharp-tail habitat, depending on the cover type, timing, frequency, intensity, size of burn (Giesen and Connelly 1993), soils, and precipitation. Many species of deciduous shrubs (for example chokecherry and rose) resprout after fire. In contrast, most sagebrush species do not resprout and may be eliminated by fires. Exotic vegetation can invade following fire, depending on the soils and precipitation.

Human disturbances related to the expansion of residential developments, increases in road densities, and associated recreational activities will likely exacerbate losses of suitable habitat within the historical range of Columbian sharp-tailed grouse (Giesen and Connelly 1993, Tirhi 1995).

Population status and trends--Sharptails currently occupy <5 percent of their historical range in the basin. The BBS data summarized for western North America indicate that population trends declined by an average of -7.7 percent annually between 1966 and 1995 ($n = 39$, $P < 0.05$; Sauer and others 1996).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 36 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The following issues were identified by our analysis of source habitat trends and from the findings of other studies on Columbian sharp-tailed grouse:

1. Fragmentation and loss of mesic shrubsteppe and steppe habitats by conversion to agriculture.
2. Alteration of shrubsteppe and steppe habitats by invasions of exotic forbs and grasses.
3. Degradation and loss of cover types within the shrubsteppe, steppe, mountain shrub, herbaceous wetlands, and shrub wetland community groups by excessive livestock grazing.
4. Loss of sagebrush cover because of burning, herbicide spraying, and other brush control methods.
5. Human disturbance of leks and wintering populations because of increased roading and human

presence.

6. Increased application of pesticides in and near agricultural areas.
7. Loss of CRP lands by conversion back to active croplands.
8. Isolated and disjunct populations vulnerable to extinction by stochastic events (that is, demographic, environmental, or genetic stochasticity).

Potential strategies--The issues identified above suggest the following broad-scale strategies for the long-term persistence of Columbian sharp-tailed grouse:

1. (To address issue no. 1) Basin-wide, identify areas of high ecological integrity, mesic shrubsteppe vegetation and manage to promote their long-term sustainability.
2. (To address issue no. 2) Restore shrubsteppe and steppe habitats that have been altered by medusahead grass, cheatgrass, and exotic mustards, and focus on areas that would increase patch size or links with existing source habitat patches.
3. (To address issue no. 2) Protect shrubsteppe habitats against wildfire in areas vulnerable to invasion by exotic vegetation.
4. (To address issue no. 3) Reduce habitat degradation by livestock grazing in cover types within shrubsteppe, mountain shrub, riparian, grassland, and herbaceous wetland terrestrial community groups that are currently occupied by sharptails, with a high potential of being recolonized by sharptails, or that have been identified for reintroductions.
5. (To address issue no. 4) Maintain sagebrush and mountain shrub cover, and increase these shrublands in areas where substantial losses have occurred because of brush control, especially in locations currently occupied by sharptails, with a high potential of being recolonized by sharptails, or in locations that have been identified for reintroductions.
6. (To address issue no. 7) Maintain CRP lands that are currently occupied by sharptails, lands that have a potential of being used by sharptails, or near locations that have been identified for reintroductions.
7. (To address issue no. 8) Expand the current range of Columbian sharptails within their historical habitats.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy nos. 1 and 4) Establish special management areas for the maintenance of high-integrity shrublands where livestock grazing would be eliminated or restricted. Manage areas of at least 200 ha (494 acres) for summer nesting and brood-rearing habitat, and suitable wintering habitats should be within 2.6 to 20 km (1.6 to 12.5 miles) of summer areas.
2. (In support of strategy no. 2) Restore degraded shrubsteppe, mountain shrub, and shrub wetland habitats by plantings of native shrub and herbaceous vegetation, and by prescribed fire (in areas not vulnerable to invasion by exotic plants).
3. (In support of strategy no. 2) Develop methods through ongoing or new research to restore

shrubsteppe habitats altered by medusahead, cheatgrass, and exotic mustards.

4. (In support of strategy no. 3) Plant native vegetation that is naturally resistant to wildfire, and actively suppress wildfires in areas that are susceptible to postfire invasions of exotic vegetation.
5. (In support of strategy no. 4) Remove or explicitly control the timing and intensity of grazing to improve the ecological condition of degraded rangelands in locations occupied by sharptails, with a high potential of being recolonized by sharptails, or that have been identified for reintroductions.
6. (In support of strategy no. 5) Eliminate brush control for sagebrush and mountain shrubs in those areas currently occupied or with a high potential of being recolonized by sharptails, including the Snake Headwaters, Upper Snake, Central Idaho Mountains, Blue Mountains, and Columbia Plateau ERUs.
7. (In support of strategy no. 6) Promote the continuation and development of the CRP program, whereby private landowners are encouraged to reduce soil erosion and establish perennial cover, especially in the Upper Snake and Snake Headwaters ERUs.
8. (In support of strategy no. 7) Acquire lands that are currently occupied by sharptails but are not specifically managed for the grouse.
9. (In support of strategy no. 7) Reintroduce and augment sharp-tailed grouse populations after habitat enhancement.

GROUP 37--GRASSHOPPER SPARROW, CLAY-COLORED SPARROW, AND IDAHO GROUND SQUIRREL

Results

Species ranges, source habitats, and special habitat features--Group 37 consists of breeding habitat for the grasshopper sparrow and clay-colored sparrow, and year-round habitat for the Idaho ground squirrel. The breeding range of the grasshopper sparrow (fig. 111) includes most of the basin except for the Northern Great Basin, Upper Klamath, Southern Cascades, and Northern Cascades ERUs. The breeding range of the clay-colored sparrow (fig. 111), on the other hand, is restricted to the Northern Glaciated Mountains, Upper Clark Fork, and Snake Headwaters ERUs. Within the basin, ranges of these two sparrow species overlap only in Montana and Wyoming. The Idaho ground squirrel is endemic to the basin and has a limited range that overlaps with the grasshopper sparrow within the eastern portion of the Blue Mountains ERU and western portion of the Central Idaho Mountains ERU (fig. 111).

Vol. 2, Figure 111--Ranges of species in group 37 within the basin.

Fescue-bunchgrass is the one cover type shared by all three species (vol. 3, appendix 1, table 1). Open-canopied mountain big sagebrush is source habitat used by the grasshopper sparrow and Idaho ground squirrel. Additionally, the open-canopied big sagebrush is source habitat for the ground squirrel. The clay-colored sparrow also has source habitats in chokecherry-serviceberry-rose and native forb cover types.

Neither sparrow has a clear preference for any special habitat features, but the clay-colored may be attracted to sites that have dense shrubs in a matrix of more open grasslandlike vegetation (Janes 1983). Idaho ground squirrels inhabit meadows, usually with shallow soils and small intrusions of deeper soil for nest burrows (USDA Forest Service and USDI Fish and Wildlife Service 1996).

Broad-scale changes in source habitats--Historically, source habitats for the sparrows in this group were widespread, but generally occupied <25 percent of most watersheds (fig. 112A). High percentages of contiguous watersheds with source habitats occurred in the northeast end and along the eastern edge of the Columbia Plateau ERU, and in the northern end of the Blue Mountains ERU. In the rest of the basin, however, large, contiguous source habitats of high ecological integrity were small and scattered. Nonetheless, the sparrows likely occupied relatively small patches of suitable habitat throughout their historical ranges.

Habitat loss has been obvious as both contiguous areas of source habitats and watersheds with relatively less habitat have greatly diminished (fig. 112B). The Columbia Plateau and Blue Mountains ERUs had strongly declining trends in source habitats for grasshopper sparrows. Similarly, the small but important source areas for the clay-colored sparrow in the northeastern portion of the basin and for Idaho ground squirrel in the center of the basin have decreased. Although much of the basin never had a high percentage of watersheds with source habitats, large acreages have been converted to landscapes with no habitat (fig. 112B).

Over 60 percent of the watersheds had strongly declining trends in source habitats basin-wide (fig. 113). Within the two ERUs that constitute the heart of the habitat for grasshopper sparrow, the Columbia Plateau and Blue Mountains, changes were markedly negative (fig. 113). Similarly, where the two sparrows occur together in the Northern Glaciated Mountains and Upper Clark Fork ERUs, trends were clearly declining (fig. 113). Source habitats for the ground squirrel were projected to have undergone the second greatest decline among 91 species evaluated (vol. 1, table 7). All three species in this group were in the habitat trend category with the greatest decrease in source habitats (vol. 1, table 7).

Vol. 2, Figure 112--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 37.

Vol. 2, Figure 113--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 37, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--The principal vegetation change corresponding to the negative trend in source habitats was in the fescue-bunchgrass cover type, which declined two-thirds from historical levels basin-wide (Hann and others 1997). The largest declines within the species ranges occurred in the Columbia Plateau and Northern Glaciated Mountains (>80 percent); Blue Mountains (75 percent); and Upper Clark Fork and Central Idaho

Mountains (60 percent; vol. 3, appendix 1, table 4). The decrease in fescue-bunchgrass amounted to over 5 percent of all changes in the basin, an amount exceeded only by the decrease in big sagebrush (Hann and others 1997).

The open-canopy low-medium structural stage of mountain big sagebrush and big sagebrush experienced some of the greatest absolute declines on an ERU basis. The combined absolute decline for the open-canopy low-medium structural stage of these two sagebrush types declined in the Upper Snake (-40 percent), Owyhee Uplands (-20 percent), Columbia Plateau (-13 percent), Snake Headwaters (-7 percent), and Northern Great Basin (-2 percent) (vol. 3 appendix 1, table 4). In these open-canopied cover types, in the absence of fire, shrubs and trees eventually invade much of the area that was occupied by grasses and forbs.

Basin-wide declines in mountain big sagebrush were substantial (Hann and others 1997) and resulted in critical losses of source habitats for the grasshopper sparrow and Idaho ground squirrel. Vegetation changes affecting Idaho ground squirrels may be difficult to discern for small meadows of sagebrush or native herbaceous cover types within ponderosa pine-dominated forests. This mosaic of habitats is not always detectable at the 1-km² (0.4-mi²) pixel size that was used for evaluating habitat trends in this effort.

Increases in the Central Idaho Mountains were due to the large relative increase in native forbs, although this cover type occupies only a small fraction of the ERU (vol. 3, appendix 1, table 4).

Conversion of upland shrubland to agriculture affected 9 percent of the basin (Hann and others 1997). Major conversions in the Columbia Plateau, Owyhee Uplands, and Blue Mountains had large effects on this group. The basin-wide loss of fescue-bunchgrass and wheatgrass-bunchgrass cover types was largely the result of conversion to agriculture. Transition of upland herbland to agriculture affected 7 percent of the basin, a conversion rate second only to that for upland shrubland (Hann and others 1997). Conversion in the Columbia Plateau and Blue Mountains was particularly high--up to 25 percent of upland shrublands. Basin-wide declines in mountain big sagebrush and native forbs also were attributed in part to agricultural conversion.

Habitat condition for group 37 can be described by the Composite Ecological Integrity Ratings (Quigley and others 1996) that show most of the habitat to have a "low" rating. Fescues and bunchgrasses, critical habitat components for this group, were irreversibly modified by high-intensity grazing in the late 1800s to early 1900s (USDA Forest Service 1996). Most of the current habitat for this group was classified into Rangeland Clusters 5 (generally corresponding to much of the Owyhee Uplands ERU) and 6 (generally the Northern Great Basin, Owyhee Uplands, and Upper Snake ERUs), where the primary risk to ecological integrity is continued losses of herbland and shrubland habitats (Quigley and others 1996). Further, Rangeland Cluster 6 is vulnerable to overgrazing and exotic grass and forb invasions (Quigley and others 1996).

Other factors affecting the group--For the Idaho ground squirrel, meadow habitats of sagebrush and herbaceous vegetation surrounded by pine forests are decreasing because of forest encroachment and human developments (USDA Forest Service and USDI Fish and Wildlife Service 1996). The juxtaposition of these habitats is difficult to evaluate at the scale of this analysis.

Early season mowing of hayfields causes major nest failures in grassland-nesting species (Knapton 1994, Smith 1963). Where hayfields and similar agricultural lands have replaced native source habitats or are now located adjacent to such habitats, those sites likely serve as significant population sinks, particularly for grasshopper sparrows.

Grasshopper sparrow populations temporarily decline immediately after grassland fires (Bock and Bock

1992). Birds likely avoid recently burned areas because of the lack of grass cover, and they are expected to return to burned sites after grasses are restored. This sparrow also avoids areas where shrub cover exceeds 35 percent (Bock and Bock 1992, Smith 1963). Thus, fire plays a beneficial role in habitat management for this species.

Although clay-colored sparrows are sympatric with grasshopper sparrows in some regions, clay-colored sparrows prefer the other end of the grass-shrub gradient, becoming more common with increases in shrub cover and patches of shrubs (Knapton 1979, 1994; Owens and Myers 1973). Thus, clay-colored sparrows also will respond negatively, in the short term, to burning and may require more time to return to prefire population densities while shrubs become reestablished after fire (Pylypec 1991).

Species in this group evolved in shrubsteppe habitats, where microbiotic crusts were broadly distributed (see Kaltenecker and Wicklow-Howard 1994). Microbiotic, or cryptogamic, crusts consist of lichens, bryophytes, algae, microfungi, cyanobacteria, and bacteria growing on or just below the soil surface in arid and semiarid environments (Kaltenecker and Wicklow-Howard 1994); these crusts developed in the absence of large herds of grazing ungulates (St. Clair and Johansen 1993). In addition, these crusts are projected to have been widely distributed throughout the source habitats for this group, particularly in the Northern Great Basin, Owyhee Uplands, and Upper Snake ERUs but also scattered in the Columbia Plateau ERU (Hann and others 1997, map 3.59). Increasing evidence indicates that microbiotic crusts improve soil stability, productivity, and moisture retention, moderate extreme temperatures at the soil surface, and enhance seedling establishment of vascular plants (Belnap and Gardner 1993, Harper and Pendleton 1993, Johansen and others 1993, St. Clair and others 1993), thereby contributing to high ecological integrity of shrubsteppe habitats. Idaho BLM has recognized the potential importance of microbiotic crusts by proposing standards for rangeland health that include the maintenance of these crusts to ensure proper functioning and productivity of native plant communities (USDI Bureau of Land Management 1997). These crusts were widely destroyed by trampling during the excessive livestock grazing of the late 1800s and early 1900s (Daubenmire 1970, MacCracken and others 1983, Mack and Thompson 1982, Poulton 1955). Currently, high-intensity grazing and altered fire regimes modify shrubsteppe plant communities and threaten the maintenance and recovery of microbiotic crusts (Belnap 1995, Kaltenecker 1997, St. Clair and Johansen 1993).

Grazing may reduce or completely exclude grasshopper sparrow populations (Bock and Webb 1984, Saab and others 1995) because livestock remove grass, the main feature of a given site that attracts this species (Janes 1983).

The grasshopper sparrow may be area sensitive and more likely to occupy large tracts of habitat than small fragments (Samson 1980). Minimum area requirements in Maine are around 100 ha (247 acres) (Vickery and others 1994), and in Illinois are about 30 ha (74 acres) (Herkert 1994).

Although brown-headed cowbirds parasitize nests of grasshopper sparrows, the impact is believed to be generally low because of the cryptic nature of the nests of sparrows (Vickery 1996). Cowbirds also parasitize nests of clay-colored sparrows, which may accept or reject the eggs. The overall impact on this species is not known but may be lower than in many species, as cowbird parasitism accounts for only 22 percent of egg loss (Knapton 1994).

Idaho ground squirrels are threatened by sport shooting or "plinking" (Moroz 1995). Several sites occupied by the ground squirrels are regularly visited by shooters for this purpose. When populations are small, this activity could have a critical, detrimental impact. Increases in human occupation in the basin likely has caused an increase in human disturbance.

Idaho ground squirrels may experience competition with Columbian ground squirrels (Moroz 1995, USDA Forest Service and USDI Fish and Wildlife Service 1996). Both species use similar habitats, but the

Idaho ground squirrel tends to inhabit more xeric areas that cannot support Columbian ground squirrels. Columbian ground squirrels are larger and require larger areas with deeper soils. Although the Idaho ground squirrel can use the same habitats for hibernation, it may be competitively forced into the drier areas with more shallow soils. The shallow soil areas are more prone to fluctuating water tables and freezing during harsh winters, causing overwinter mortality in Idaho ground squirrels (Moroz 1995).

Low population numbers of the Idaho ground squirrel, probably no more than 600 to 800 individuals, make the species vulnerable to (1) genetic drift, inbreeding, and attendant loss of viability; (2) catastrophic invasions of predators, parasites, or diseases; and (3) extirpation because of natural population fluctuations (Moroz 1995). Populations are small and often isolated by several kilometers (Yensen 1991).

Poisoning through the use of rodenticides may negatively affect populations. Predation by domestic cats also is a concern (USDA Forest Service and USDI Fish and Wildlife Service 1996).

Forest encroachment into meadows may be a threat to Idaho ground squirrels because of fire suppression and natural succession (Moroz 1995). Encroachment on meadows and replacement of open forest stands with dense stands of trees may have eliminated or reduced dispersal corridors (USDA Forest Service and USDI Fish and Wildlife Service 1996).

Population status and trends--Sample sizes for the clay-colored sparrow in the basin were insufficient to determine population trend (Saab and Rich 1997). The 1966-95 trend for BBS physiographic region 64 (Central Rocky Mountains) is +11.4 percent per yr ($\bar{n} = 17$, $\underline{P} < 0.05$), but the sample size is small (Sauer and others 1996).

Saab and Rich (1997) reported a stable population trend for the grasshopper sparrow in the basin but also stated that the species is not well monitored by the BBS technique and advised specialized monitoring. The trend for Washington is +7.5 percent per yr ($\bar{n} = 18$, $\underline{P} < 0.1$) and for physiographic region 89 (Columbia Plateau) is stable ($\bar{n} = 24$, $\underline{P} > 0.1$; Sauer and others 1996). Again, sample sizes are too small to provide definitive results.

Currently, Idaho ground squirrels occupy 13 small and 2 larger meadows on both public and private land. Many of the habitats are small, isolated, or in poor condition (USDA Forest Service and USDI Fish and Wildlife Service 1996). All but one of the populations number <200 individuals.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 37 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--The results of our habitat trend analysis suggest the following issues are of high priority for group 37:

1. Continued loss of large acreage of fescue-bunchgrass and mountain big sagebrush cover types.
2. Loss of microbiotic crusts.
3. Undesired changes in shrub:grass ratios because of changes in historical fire regimes.
4. Direct mortality of ground nesting birds because of agricultural practices.

5. The disjunct nature of remaining habitat for grasshopper sparrow populations.
6. Loss of meadow habitat because of forest encroachment and human developments.
7. Loss of dispersal corridors for Idaho ground squirrel from replacement of open forest stands with dense stands.
8. Vulnerability to extinction of small, isolated populations of ground squirrels because of poisoning, shooting, predation, disease, or natural fluctuations.
9. Displacement from habitat because of interspecific competition.

Potential strategies--The following strategies could be used to reverse broad-scale declines in source habitats:

1. (To address issue nos. 1 and 5) Identify and conserve remaining large areas of mountain big sagebrush and fescue-bunchgrass vegetation where ecological integrity is still relatively high (Bock and others 1993, Gray and Rickard 1989, Rickard and Poole 1989, Schuler and others 1993, Smith 1994, Yoakum 1980). The remaining blocks of habitat in the eastern Blue Mountains and southern Central Idaho Mountains ERUs (fig. 112) may serve as focal points for protection. For the clay-colored sparrow, only the small watersheds in the Upper Clark Fork and Northern Glaciated Mountains ERUs (fig. 112) can be expected to contribute to source habitats.
2. (To address issue no. 1) Restore native perennial bunchgrasses and avoid further depletion because of improper grazing (Braun and others 1976, Daubenmire 1970, Evans and Young 1978). Priority areas for the grasshopper sparrow are the eastern Blue Mountains and southern Central Idaho Mountains ERUs (fig. 113). For the clay-colored sparrow, priority areas are the Upper Clark Fork and Northern Glaciated Mountains ERUs.
3. (To address issue no. 2) Restore microbiotic crusts in ERUs with potential for redevelopment (that is, areas near propagule sources, and with suitable soil, vegetation, and climatic characteristics [see Belnap 1993, 1995; Kaltenecker 1997; Kaltenecker and Wicklow-Howard 1994]): the Northern Great basin, Owyhee Uplands, and Upper Snake ERUs and, to a lesser extent, the Columbia Plateau ERU (Hann and others 1997, map 3.59).
4. (To address issue no. 3) Use fire to obtain desired shrub:grass ratios. Enhance development of shrub communities, particularly mountain sagebrush and chokecherry-serviceberry-rose, in the Upper Clark Fork and Northern Glaciated Mountains ERUs. Maintain dense grassland cover in the eastern Blue Mountains and southern Central Idaho Mountains ERUs.
5. (To address issue no. 4) Minimize direct mortality of ground nesting birds in agricultural areas.
5. (To address issue no. 5) Maintain and restore the largest areas of native grassland habitats.
7. (To address issues nos. 6 and 7) Maintain meadows and corridors currently used by the squirrel. Restore potentially suitable meadows within the range of the species. Stop or reverse forest encroachment into meadows.
8. (To address issue no. 8) Prevent direct human-caused mortality of Idaho ground squirrels.
9. (To address issue no. 8) Restore populations of the Idaho ground squirrel.

10. (To address issue no. 9) Explore the removal of Columbian ground squirrels from adjacent habitats.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Use landscape planning to avoid further reductions in the size of large blocks of mountain big sagebrush and fescue-bunchgrass within each watershed, particularly in the Blue Mountains and Central Idaho Mountains ERUs, where sizeable blocks of source habitats are available.
2. (In support of strategy no. 1) Explore options under the CRP (Johnson and Igl 1995), or develop other incentive programs, to encourage restoration of agricultural areas to native cover types. Focus on areas that would increase patch size or links with existing source habitat patches.
3. (In support of strategy nos. 2 and 3) Modify grazing systems or reduce grazing use where native perennial bunchgrasses have been depleted. The elimination of grazing may encourage the redevelopment of microbiotic crust (Mack and Thompson 1982, St. Clair and others 1993).
4. (In support of strategy no. 3) Explore the use of ground-based and aerial soil inoculation to increase the speed and extent of dispersal of the organisms that create microbiotic crust (Belnap 1993, 1994).
5. (In support of strategy nos. 1, 3, and 4) Develop a prescribed burning program designed to increase native grass cover and reduce shrub cover (Vickery 1996) on limited acreages and in concert with strategy no. 1. For example, summer burns, which correspond to the period of increased natural lightning strikes, may be more beneficial for maintaining source habitats than burns at other times of the year (Shriver and others 1996); extensive, hot burns in shrubsteppe habitats are probably less beneficial than cooler, more controlled burns that leave some shrub cover (Bock and Bock 1987). In clay-colored sparrow habitats, fire control will allow development of the shrub component that this species prefers (Knapton 1994).
6. (In support of strategy no. 5) Where possible, avoid early season mowing of hayfields and other agricultural lands (Rodenhouse and others 1995, Vickery 1996). Defer mowing on publicly owned lands and develop incentives for private land owners (Vickery 1996). Avoid creating hayfields and similar crop fields adjacent to, or in the general area of, natural nesting habitats.
7. (To address strategy no. 6) A breeding site of 100 to 200 pairs in an area of source habitats 800 to 1400 ha (1,330 to 2,330 acres) is recommended to sustain a population of grasshopper sparrows (Delany and others 1995). Avoid fragmenting existing source habitats below this size and work to protect and restore other sites to at least this standard.
8. (In support of strategy no. 7) Maintain meadow and meadow-corridor habitats within ponderosa pine cover types for Idaho ground squirrels. Retard conifer invasion of meadows by thinning young trees from stands, prescribed burning, and controlled grazing (Moroz 1995). Replant with native grasses.
9. (In support of strategy no. 7) Develop livestock grazing practices that retain grass seed-heads available to ground squirrels (Moroz 1995).
10. (In support of strategy no. 7) Create new meadow habitats at suitable locations with various deep and shallow soils. Expand existing meadow habitats through practices in issue no. 6, with attention to corridors that could provide dispersal habitats for existing populations of Idaho ground squirrels.
11. (In support of strategy no. 8) Avoid use of rodenticides in occupied habitats.

12. (In support of strategy no. 8) Control recreational uses such as off-road vehicles, roadside turnouts, and camping within meadow complexes occupied by the squirrel. Encourage the public to avoid shooting, poisoning, or trapping the squirrel. Close important ground squirrel areas to discharge of firearms. Inform the public about this endemic Idaho species.
13. (In support of strategy no. 9) Reintroduce Idaho ground squirrels into suitable habitats.
14. (In support of strategy no. 10) Determine if removal or reduction of Columbian ground squirrel populations will provide more habitat for the Idaho ground squirrel.

GROUP 38--BLACK ROSY FINCH AND GRAY-CROWNED ROSY FINCH

Results

Species ranges, source habitats, and special habitat features--Group 38 consists of the black rosy finch and the gray-crowned rosy finch, summer residents of alpine communities. The gray-crowned rosy finch occurs throughout the basin, whereas the black-crowned rosy finch is restricted to the eastern part of the basin (fig. 114). This analysis is focused on summer habitat only. Both finches winter in open habitats at lower elevations and occasionally are observed in towns.

Vol. 2, Figure 114--Ranges of species in group 38 within the basin.

Source habitats for group 38 are alpine tundra, barren rocky areas, and cliffs (vol. 3, appendix 1, table 1). Rosy finches nest primarily on cliffs in rocky crevices (French 1959), which are a special habitat feature used by these species. Both finches feed on seeds and insects (French 1959).

Broad-scale changes in source habitats--Source habitats coincide with the distribution of alpine tundra, both historically and currently (figs. 115A, B). The greatest amount of source habitat occurs in the Rocky Mountains in Montana (fig. 115B). No change in amount of source habitats was projected for this group (figs. 115C and 116).

Vol. 2, Figure 115--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 38.

Vol. 2, Figure 116--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 38, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Neutral trends in source habitats were attributed to insignificant changes in the amount of alpine tundra since historical times (Hann and others 1997). These projections were limited by the coarse resolution of the data. Hann and others (1997) suspected that finer resolution data would indicate long-term degradation of soils and changes in the composition of vegetation resulting from excessive domestic sheep grazing within alpine environments. Thus, the projected neutral trend should be interpreted as describing habitat extent but not habitat quality.

Condition of special habitat features--Changes in the abundance of rocks and cliffs have not been documented but likely are insignificant.

Other factors affecting species within the group--Potential overgrazing by sheep and human recreational activities in alpine tundra could have a negative effect on habitat suitability for these species (ICBEMP 1996g, Lehmkuhl and others 1997). Rock climbing could cause local disturbances of nest sites.

Population status and trends--Trend data for populations of the black rosy finch or the gray-crowned rosy finch are not available. Low population numbers and limited habitat contribute to conservation concerns for both species (ICBEMP 1996g, Marshall and others 1996).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 38 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--Results of our habitat trend analysis do not lead to any management issues at the broad-scale. Expert opinions (ICBEMP 1996g, Lehmkuhl and others 1997), however, suggest the following issues may be important for the long-term viability of rosy finches:

1. Declines in quality of alpine vegetation in the basin because of past and current sheep grazing and recreational activities.
2. Disturbance to cliff and rock nest sites.

Potential strategies--The primary strategy for addressing issue no. 1 is to minimize negative effects of grazing and recreational activities in alpine tundra habitat. Because of lack of information on the degree of impacts to rock and cliff nest sites, no strategies are proposed for issue no. 2.

Practices that support strategies--The following practices would be effective in implementing the strategy listed above:

1. Restrict human access and livestock use in heavily degraded areas of alpine tundra.
2. Modify grazing allotment plans and trail use regulations to prevent declines in good quality habitat.
3. Restore alpine areas that are in a degraded condition.

GROUP 39--LEWIS' WOODPECKER (RESIDENT POPULATION)

Results

Species ranges, source habitats, and special habitat features--Resident Lewis' woodpeckers are distributed in a small area of open woodlands in the northern end of the Southern Cascades and in southern portions of the Northern Cascades ERUs (fig. 117), along the eastern foothills of the Cascade Range. Birds use this area year round, unlike migratory Lewis' woodpeckers described in group 2 that use the basin only during the breeding season. Source habitats of the resident Lewis' woodpecker include oak woodlands (vol. 3, appendix 1, table 1), parklike pine-oak, burned pine-fir forests, and cottonwood groves (Galen 1989). These vegetation types apparently were most abundant, historically and currently, in a small area within the northern portion of the Southern Cascades ERU (fig. 118).

Vol. 2, Figure 117--Ranges of species in group 39 within the basin.

Unlike most woodpecker species, Lewis' woodpecker is an aerial insectivore and requires openings for their foraging maneuvers. This woodpecker breeds in habitats that provide abundant insects (see group 2 for a broader discussion on migratory Lewis' woodpeckers), and winters in areas where temperatures are warm enough to support flying insects and where acorns are abundant. Acorns are harvested in the fall and stored for winter use. Birds overwinter within the basin where a reliable acorn supply is available (Galen 1989).

Because this species has weak excavator morphology (Spring 1965), Lewis' woodpeckers typically require large snags in an advanced state of decay or trees with soft sapwood for ease of cavity excavation (Bock 1970, Raphael and White 1984, Saab and Dudley 1995, Tobalski 1997). Additionally, Lewis' woodpeckers usurp occupied cavities (Saab and Dudley 1995) or reuse old cavities created by strong excavators (that is, hairy woodpecker, black-backed woodpecker and Northern flicker) or nest in natural cavities of trees (Bock 1970, Galen 1989, Saab and Dudley 1995, Tashiro-Vierling 1994, Vierling 1997).

Nest tree species of resident birds in the basin were primarily Oregon white oak and ponderosa pine, and less commonly Douglas-fir and cottonwood (Galen 1989). Snag and tree diameters used for nesting are generally larger and more heavily decayed than that expected based on availability of such snags (see group 2 for description of source habitats). In north-central Oregon, tree diameters of 23 nests in Oregon white oak ranged from 31.8 to 99 cm (12.5 to 39 in) and averaged 55.9 cm (22 in), and tree height ranged from 3.0 to 15.2 m (10 to 50 ft) and averaged 9.7 m (32 ft) (Galen 1989). Most of these nest trees, however, were living or had light decay. Heavily decayed trees, typical of nest trees elsewhere (see group 2 for source habitat description), were probably not necessary in north-central Oregon because nesting only occurred in preexisting cavities, and there was no evidence of Lewis' woodpeckers excavating new cavities (Galen 1989).

Nesting habitat in north-central Oregon was usually open pine-oak woodlands and burned coniferous forests (Galen 1989). Nests also were located in cottonwood groves and narrow oak groves adjacent to open areas. No nests were found in scrub-oak thickets along south-facing slopes, unburned coniferous forests, or clearcuts. Proximity to openness was considered a critical microhabitat feature for breeding habitat (Galen 1989). Open woodlands provide sufficient visibility and space for effective flycatching. Most nests (36 of 53) were located in areas with >75 percent open canopy. Snags were also an

important component of nesting habitat. Snags were used for perching during the breeding season and for acorn storage during winter.

Nesting densities of resident woodpeckers in Oregon differed from one breeding pair per 8 ha (20 acres) of woodland to one breeding pair per 16 ha (40 acres), depending on suitable snags, trees, and cavities available for nesting (Galen 1989). Nesting habitat required for one pair of Lewis' woodpeckers was estimated at 10 ha (25 acres) of open pine-oak, oak or cottonwood when these woodlands are adjacent to open areas of equal or greater size (Galen 1989).

Wintering habitat of resident Lewis' woodpeckers in the basin was associated with nest trees used during the breeding season (Galen 1989). Nearly 90 percent of 46 nests showed signs of wintering woodpeckers. Acorns were stored in nest trees or in adjacent snags, and oaks were nearby.

In foothills habitat of southeastern Colorado, acorns were the primary winter food source (Vierling 1997). Acorn crops were higher at occupied winter sites than at random sites. Availability of storage sites for mast was a critical feature of winter habitat (Vierling 1997). Storage trees were significantly taller (\bar{x} = 17.5 m versus 10.9 m [57.8 ft vs. 36 ft]) and of larger diameter (\bar{x} = 104.8 cm versus 61.7 cm [41.3 in versus 24.3 in]) than random trees (Vierling 1997). Crevices in dead and decaying trees, and the deep furrowed bark of cottonwoods, were important characteristics of acorn storage sites.

Broad-scale changes in source habitats--No apparent broad-scale changes occurred in breeding and wintering source habitats of resident Lewis' woodpeckers (figs. 118A, B; 119).

Vol. 2, Figure 118--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 39.

Vol. 2, Figure 119--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 39, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Areal extent of oak woodlands, the only source habitats used for this group, was not estimated to have changed using the large pixel size of this analysis (vol. 3, appendix 1, table 4; figs. 118A, B). This cover type is limited within the basin and has had few threats of logging activities but greater threats by development and firewood cutting. Changes in oak woodlands may not be discernable where oaks occur in small stands or where they occur within conifer stands. Nearby pine-oak, burned pine forests, and cottonwood woodlands used by this resident population were not evaluated in the broad-scale analysis. Thus, a broad-scale analysis for this group has limited application.

Condition of special habitat features--Abundance of large, heavily decayed snags for nesting and acorn storage may have declined in the range of resident Lewis' woodpeckers within the basin. Densities of

large-diameter snags (>53 cm [21 in] d.b.h.) have declined basin-wide from historical to current levels (Hann and others 1997; Hessburg and others 1999; Quigley and others 1996).

Oak mast-producing trees are critical for overwinter survival of Lewis' woodpeckers in the basin. A 500-yr-old Oregon white oak attains large diameters (58 to 89 cm d.b.h. [23 to 35 in]) on generally dry slopes that offer slower growing conditions (Galen 1989). Destruction of these old and mature trees by clearing for pastures and firewood cutting could jeopardize resident Lewis' woodpeckers.

Open woodlands that allow foraging maneuvers have probably decreased as a result of fire control practices. Historically, oak woodlands in Washington were maintained by frequent wildfires, and through controlled burning by early inhabitants (Ryan and Carey 1995). Oak woodlands currently are threatened by encroachment of ponderosa pine and Douglas-fir. Fire control also likely has reduced understory shrubs and associated arthropods that provide food during the breeding season. Additionally, understory shrubs may have been altered by disturbances of grazing practices and recreational activities.

Other factors affecting the group--Road densities have significantly increased throughout the basin (Hann and others 1997, Quigley and others 1996), allowing greater human access into forested regions and subsequent increases in snag removal for firewood. Salvage logging is another threat to snags that provide potential nest sites (Marshall and others 1996). Prolonged human presence at or near nest sites may cause abandonment (Bock 1970); however, stable populations coexist with park development and heavy tourist use during the breeding season in British Columbia (Siddle and Davidson 1991).

Chlorinated hydrocarbons, particularly DDT, which were formerly used as pesticides in fruit orchards and gardens, could have potential negative effects on Lewis' woodpeckers (Tobalski 1997) because these woodpeckers sometimes nest in agricultural settings (Sorensen 1986, Tashiro-Vierling 1994). Elevated energetic costs and stress may be associated with high rates of territorial encounters with European starlings and this could reduce reproductive success even if Lewis' woodpecker dominates the interaction (Siddle and Davidson 1991). Altered fire regimes and subsequent changes in the structure and composition of lower montane forests (Hann and others 1997) could reduce suitable oak woodlands for breeding and wintering Lewis' woodpeckers. Large cottonwoods, used for nesting and acorn storage, are threatened by altered hydrologic regimes, grazing practices, and urban development (Marshall and others 1996).

Population status and trends--No population trends are available for the resident Lewis' woodpeckers that occupy the eastern foothills of Mount Hood. Breeding Bird Surveys (BBS) for the entire basin indicate that population trends have been stable during 1968-94 (Saab and Rich 1997), but any relation to the resident population is not known. Trend data generated by the BBS may be more adequate for monitoring populations of resident Lewis' woodpeckers than migratory populations (see group 2, "Population status and trends"). Dramatic cycles of population abundance related to local changes in habitat (Bock 1970) may not apply to resident birds that will use acorns as a year-round food source, supplemented by insects during the breeding season.

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 39 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--

1. Exclusion of fire in parklike oak and pine-oak woodlands and subsequent decreases in natural forest

openings and shrubby understories because of invasions by conifers (Marshall and others 1996).

2. Losses of large oak trees for mast production, as a result of firewood cutting, fire control, and pasture development.
3. Decline in availability of large, heavily decayed ponderosa pine for nesting and acorn storage sites.
4. Losses of large cottonwoods used for nesting and acorn storage (Marshall and others 1996).
5. Increase in application of agricultural insecticides.

Potential strategies--The issues identified above suggest the following broad-scale strategies for the long-term persistence of resident Lewis' woodpeckers in the northern portion of the Southern Cascades ERU.

1. (To address issue no. 1) Return natural fire regimes to oak and pine-oak woodlands.
2. (To address issue nos. 2, 3, and 4) Retain large (> 30 cm d.b.h. [12 in]), old snags and trees of Oregon white oak, ponderosa pine, and cottonwoods (Galen 1989).
3. (To address issue nos. 3 and 4) Protect acorn storage sites in wintering areas (Galen 1989, Marshall and others 1996).
4. (To address issue no. 4) Maintain existing old-growth cottonwood forests and manage young forests for the long-term sustainability of cottonwood/riverine systems.
5. (To address issue no. 5) Avoid use of toxic chlorinated hydrocarbons and organophosphorus insecticides near Lewis' woodpecker nesting and wintering sites.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Maintain parklike oak and pine-oak woodlands by using silvicultural treatments of prescribed fire and thinning of small diameter ponderosa pine (<30 cm [12 in]).
2. (In support of strategy no. 2) Retain all Oregon white oak and ponderosa pine trees or snags over 3 m (10 feet) tall and >30 cm (12 in) d.b.h. (Galen 1989). Management of 10-ha (25-acre) units having approximately 25 percent canopy cover will likely provide nesting habitat for one pair of Lewis' woodpeckers (see Galen 1989).
3. (In support of strategy no. 3) Control fuel wood permits for removal of oaks, pines, or cottonwood used for winter storage sites. Minimize the density of roads open to motorized vehicles. Close roads after timber harvest activities, and maintain short periods during which such roads are open to reduce removal of snags along roads. In addition or as an alternative to road management, actively enforce fuel wood regulations to minimize removal of snags.
4. (In support of strategy no. 4) Survey and map existing old forests of cottonwoods and reference their locations in land management planning documents. Monitor conditions of cottonwood stands to ensure that sufficient seedling or vegetative regeneration, or both, is occurring. Identify factors limiting regeneration so that appropriate corrective measures can be taken. For example, return natural hydrologic regimes to portions of large river systems that support cottonwood riparian woodlands (that is, the Columbia River).

5. (In support of strategy no. 5) Establish zones with no use of toxic agricultural insecticides near Lewis' woodpecker breeding and wintering habitats.

GROUP 40--BROWN-HEADED COWBIRD

Results

Species ranges, source habitats, and special habitat features--Group 40 consists of the brown-headed cowbird, a migrant summer breeder found throughout the basin (fig. 120). The cowbird is considered a contrast species (vol. 3, appendix 1, table 2) because it requires a juxtaposition of contrasting vegetative structure to meet all aspects of its ecology. Foraging areas are in disturbed sites near livestock, and breeding areas generally are in forests and riparian areas where passerine densities are high (Robinson and others 1995). Source habitats for the brown-headed cowbird are the agricultural community type (vol. 3, appendix 1, table 1), and the presence of livestock is a special habitat feature. Additionally, the cowbird is dependent on the presence of active bird nests for parental care of their offspring. Nest parasitism by cowbirds has been documented for over 220 bird species, primarily passerine species, and at least 144 species have fledged cowbird young (Friedmann and Kiff 1985).

Vol. 2, Figure 120--Ranges of species in group 40 within the basin.

Although not mappable at the broad-scale of our analysis, horse corrals and pack stations in lower montane and montane community groups also provide source habitats. Associated breeding sites are located as far as 7 km (4.3 mi) (Rothstein and others 1987) from livestock areas, where cowbirds congregate to forage. Because of the presence of livestock areas, the distribution of source habitats is much greater than estimated by our broad-scale analysis.

Broad-scale changes in source habitats--Source habitats for the cowbird were probably not present in the basin historically (fig. 121A). Source habitats are now present in all ERUs and are particularly widespread in the Columbia Plateau and Upper Snake (fig. 121B). The trend in habitat availability has been strongly increasing basin-wide (figs. 121C and 122).

Vol. 2, Figure 121--Percentage of area identified as source habitats, historically and currently, and the relative change in percentage of area of source habitats from historical to current periods for group 40.

Vol. 2, Figure 122--Percentage of watersheds within five trend categories of relative change in source habitats from historical to current periods for group 40, basin-wide and by ecological reporting units.

Interpreting Results

Composition and structure of vegetation associated with changes in source habitats--Increases in source habitats were primarily attributed to the conversion of native vegetation to agriculture. The establishment of the cropland-hay-pasture cover type occurred on sites previously dominated by the fescue-bunchgrass, big sagebrush, and native forb cover types (Hann and others 1997). Agriculture now covers >10 percent of the land area in five ERUs: Columbia Plateau (estimated 44 percent), Blue Mountains (estimated 17 percent), Northern Glaciated Mountains (estimated 12 percent), Owyhee Uplands (estimated 12 percent), and Upper Snake (estimated 33 percent; vol. 3, appendix 1, table 4).

Condition of special habitat features--The presence of livestock is strongly associated with agricultural land uses throughout the basin. Livestock areas suitable for cowbird foraging, therefore, have probably increased in proportion to the estimated increase in area used for agriculture. Moreover, livestock areas in the lower montane and montane community groups likely have increased from historical conditions because of the location of pack stations adjacent to wilderness areas, parks, and rural expansion into forested areas.

Other factors affecting the group--Because cowbirds rely on other bird species to raise their young, they are affected by the same factors that govern breeding success of their selected hosts. About 50 percent of cowbird eggs are lost to normal nest-related mortality such as weather and predation (Nice 1957). Additional losses depend on the behavioral responses of the host, including egg rejection, egg burial, and nest desertion (Friedmann 1929).

Microsite conditions affect cowbird densities and parasitism rates. Cowbird numbers and parasitism rates are higher in proximity to internal forest openings, near powerline corridors, in small versus large woodlots, and near streams (Robinson and others 1995). Forest fragmentation and high edge density are conducive to successful breeding by cowbirds (Robinson and others 1995).

Population status and trends--Cowbirds have undergone a dramatic range expansion across North America, both eastward and westward. Expansion into eastern forests occurred in the late 1700s; this expansion was brought about by forest clearing and increases in agriculture and livestock uses. Colonization westward into Washington and Oregon began a century later (Rothstein 1994); this range expansion was likely associated with the clearing of lands for agricultural and livestock uses. Population trends were stable basin-wide from 1966 to 1994 (Saab and Rich 1997). Within Oregon, BBS data suggested that populations have been decreasing by 4 percent annually from 1966 to 1995 ($\bar{n} = 88$; $\bar{p} < 0.05$; Sauer and others 1996).

Management Implications

The following issues, strategies, and practices may be useful to managers as a starting point for integration of potential resource objectives for group 40 with broader, ecosystem-based objectives for all other resources on FS- and BLM-administered lands in the basin.

Issues--Issues primarily relate to the effect of nest parasitism by cowbirds on host species.

1. Reductions in nest success of host species, particularly state species of concern with known high parasitism rates.
2. Continued invasion of cowbirds into lower montane and montane community groups through the aid of small, remote livestock areas.

Potential strategies--

1. (To address issue no. 1) Minimize livestock concentrations in proximity to known source habitats for state and federally listed sensitive avian species.
2. (To address issue no. 1) Reduce parasitism rates on state species of concern.
3. (To address issue no. 2) Reduce opportunities for cowbird establishment in lower montane and montane community groups.

Practices that support strategies--The following practices would be effective in implementing the strategies listed above:

1. (In support of strategy no. 1) Consider the proximity of state species of concern before locating livestock-handling facilities on Federal land. Consider relocation of livestock facilities if such facilities exist in areas deemed important for recovery of an avian species of concern.
2. (In support of strategy no. 2) Intensively trap and remove cowbirds near nests of selected species of concern with high parasitism rates (Robinson and others 1995).
3. (In support of strategy no. 3) Delay annual establishment of livestock corrals within the lower montane and montane community groups during the early breeding season when cowbirds are actively seeking host nests (Kie 1991, Sanders and Flett 1989).
4. (In support of strategy no. 4) Consolidate remote livestock areas into fewer sites.

ABBREVIATIONS

Centimeter	(cm)
Hectare	(ha)
Inch	(in)
Kilometer	(km)
Meter	(m)
Mile	(mi)
Year	(yr)

REFERENCES

- Anon. 1989. Goshawk breeding habitat in lowland Britain. *British Birds*. 82: 56-67.
- Adams, Layne G.; Bailey, James A. 1983. Winter forages of mountain goats in central Colorado. *Journal of Wildlife Management*. 47: 1237-1243.
- Allen, Edith B. 1995. Restoration ecology: limits and possibilities in arid and semiarid lands. In: Roundy, Bruce A.; McArthur, E. Durant; Haley, Jennifer S.; Mann, David K., comps. *Proceedings, wildland shrub and arid land restoration symposium; 1993 October 19-21; Las Vegas, NV*. Gen. Tech. Rep. INT-315. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 7-15.
- Allendorf, F.W.; Harris, R.B.; Metzgar, L.H. 1991. Estimation of effective population size of grizzly bears by computer simulation. In: Dudley, E.C., ed. *The unity of evolutionary biology: Proceedings of the fourth international congress of systematic and evolutionary biology*. Vol. 2. 1990 July. College Park, MD: University of Maryland. Portland, OR: Dioscorides Press: 650-654.
- Anderson, W.L.; Duzan, R.E. 1978. DDE residues and eggshell thinning in loggerhead shrikes. *Wilson Bulletin*. 90(2): 215-220.
- Apple, L. 1994. Moving time. *Wyoming Wildlife*. 58(8): 18-23.
- Arthur, S.M.; Krohn, W.B.; Gilbert, J.R. 1989. Home range characteristics of adult fishers. *Journal of Wildlife Management*. 53(3): 674-679.
- Aubrey, K.B.; Houston, D.B. 1992. Distribution and status of the fisher in Washington. *Northwestern Naturalist*. 73: 69-79.
- Austin, G.T.; Russell, S.M. 1972. Interspecific aggression of ash-throated flycatchers on Cassin's sparrows. *Condor*. 74(4): 481.
- Autenrieth, R. 1978. Guidelines for the management of pronghorn antelope. In: *Proceedings of the 8th biennial pronghorn antelope workshop; 1978 May 2-4; Jasper, AB*. Jasper, AB: Alberta Recreation, Parks and Wildlife, Fish and Wildlife Division: 472-525.
- Autenrieth, R.E. 1981. Sage grouse management in Idaho. *Wildlife Bull.* 9. Boise, ID: Idaho Department of Fish and Game. 239 p.
- Autenrieth, Robert; Molini, William; Braun, Clait, eds. 1982. Sage grouse management practices. *Tech. Bull.* 1. Twin Falls, ID: Western States Sage Grouse Committee. 42 p.
- Bailey, T.N.; Bangs, E.E.; Portner, M.F. [and others]. 1986. An apparent overexploited lynx population on the Kenai Peninsula, Alaska. *Journal of Wildlife Management*. 50: 279-290.
- Baldwin, P.H. 1968. Predator-prey relationships of birds and spruce beetles. In: *Proceedings of the North Central Branch, Entomological Society of America; 1968 March 27-29; Denver, CO*. Lafayette, IN: Entomology Department, Purdue University. 23(2): 90-99.
- Banci, Vivian. 1994. Wolverine. In: Ruggiero, Leonard F.; Aubry, Keith B.; Buskirk, Steven W. [and others], tech eds. *The scientific basis for conserving forest carnivores: American marten, fisher, lynx*

- and wolverine in the Western United States. Gen. Tech. Rep. RM-254. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 7-37. Chapter 2.
- Bangs, Edward E.; Fritts, Steven H. 1996. Reintroducing the gray wolf to central Idaho and Yellowstone National Park. *Wildlife Society Bulletin*. 24(3): 402-413.
- Barclay, R.M.R. 1985. Long-versus short-range foraging strategies of hoary (*Lasiurus cinereus*) and silver-haired (*Lasionycteris noctivagans*) bats and the consequences for prey selection. *Canadian Journal of Zoology*. 63(11): 2507-2515.
- Barclay, R.M.R. 1986. The echolocation calls of hoary (*Lasiurus cinereus*) and silver-haired (*Lasionycteris noctivagans*) bats as adaptations for long- versus short-range foraging strategies and the consequences for prey selection. *Canadian Journal of Zoology*. 64(12): 2700-2705.
- Barclay, R.M.R. 1991. Population structure of temperate zone insectivorous bats in relation to foraging behavior and energy demand. *Journal of Animal Ecology*. 60: 165-178.
- Barnum, D.A. 1975. Aspects of western gray squirrel ecology. Pullman, WA: Washington State University. 58 p. M.S. thesis.
- Beale, D.M.; Smith, A.D. 1970. Forage use, water consumption, and productivity of pronghorn antelope in western Utah. *Journal of Wildlife Management*. 34(3): 570-582.
- Bechard, M.J.; Schmutz, J.K. 1995. Ferruginous hawk (*Buteo regalis*). In: Poole, A.; Gill, F., eds. *The birds of North America*. No. 172. Philadelphia, PA: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 20 p.
- Beck, Jonathan M.; Peterson, Charles R. 1995. Movements and habitat selection of the longnose snake (*Rhinocheilus lecontei*) in southwestern Idaho. Tech. Bull. 95-18. Boise, ID: U.S. Department of the Interior, Bureau of Land Management, Idaho State Office. 30 p.
- Belnap, J. 1995. Surface disturbances: their role in accelerating desertification. *Environmental Monitoring and Assessment*. 37: 39-57.
- Belnap, Jayne. 1993. Recovery rates of cryptobiotic crusts: inoculant use and assessment methods. *Great Basin Naturalist*. 53: 89-95.
- Belnap, Jayne. 1994. Potential role of cryptobiotic soil crusts in semiarid rangelands. In: Monsen, Stephen B.; Kitchen, Stanley G., eds. *Proceedings, ecology and management of annual rangelands; 1992 May 18-21; Boise, ID*. Gen. Tech. Rep. INT-313. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 179-189.
- Belnap, Jayne; Gardner, John S. 1993. Soil microstructure in soils of the Colorado Plateau: the role of the cyanobacterium *Microcoleus vaginatus*. *Great Basin Naturalist*. 53: 40-47.
- Benkman, C.W. 1990. Intake rates and the timing of crossbill reproduction. *Auk*. 107(2): 376-386.
- Benkman, Craig W. 1992. White-winged crossbill. In: Poole, A.; Gill, F., eds. *The birds of North America*. No. 27. Philadelphia, PA: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 20 p.

- Bentz, Jerry A.; Woodard, Paul M. 1988. Vegetation characteristics and bighorn sheep use on burned and unburned areas in Alberta. *Wildlife Society Bulletin*. 16(2): 186-193.
- Betts, B. 1996. Bat maternity tree roosts and forest management. *Natural Resource News*. Spring: 2-3.
- Betts, B.J. 1990. Geographic distribution and habitat preferences of Washington ground squirrels (*Spermophilus washingtoni*). *Northwestern Naturalist*. 71(2): 27-37.
- Black, Hal L. 1974. A north temperate bat community: structure and prey populations. *Journal of Mammalogy*. 55: 138-157.
- Blackford, J.L. 1955. Woodpecker concentrations in burned forest. *Condor*. 57(1): 28-30.
- Blair, Steve; Servheen, Greg; Gordon, Floyd [and others]. 1995. Saving all the pieces: the Idaho state conservation effort, white-headed woodpecker (*Picoides albolarvatus*) in Idaho. Boise, ID: Idaho Department of Fish and Game. 24 p.
- Block, W.M.; Brennan, L.A. 1987. Characteristics of Lewis woodpecker habitat on the Modoc Plateau, California. *Western Birds*. 18(4): 209-212.
- Blood, D.A. 1961. An ecological study of California bighorn sheep *Ovis canadensis californiana* (Douglas) in southern British Columbia. Vancouver, BC: University of British Columbia. 127 p. M.S. thesis.
- Blumton, A.K. 1989. Factors affecting loggerhead shrike mortality in Virginia. Blacksburg, VA: Virginia Polytechnical Institute and State University. 85 p. M.S. thesis.
- Blus, L.J.; Staley, C.S.; Henny, C.J. [and others]. 1989. Effects of organophosphorus insecticides on sage grouse in southeastern Idaho. *Journal of Wildlife Management*. 53(4): 1139-1146.
- Boal, C.W.; Mannan, R.W. 1994. Northern goshawk diets in ponderosa pine forests on the Kaibab Plateau. *Studies in Avian Biology*. 16: 97-102.
- Bock, C.E. 1970. The ecology and behavior of the Lewis woodpecker (*Asyndesmus lewis*). In: *Zoology*. 92/1970 May 25. Berkeley, CA: University of California Press. 91+ p.
- Bock, C.E.; Bock, J.H. 1974. On the geographical ecology and evolution of the three-toed woodpeckers, *Picoides tridactylus* and *Picoides arcticus*. *American Midland Naturalist*. 92(2): 397-405.
- Bock, C.E.; Bock, J.H. 1987. Avian habitat occupancy following fire in a Montana shrubsteppe. *Prairie Naturalist*. 19(3): 153-158.
- Bock, C.E.; Bock, J.H. 1992. Response of birds to wildfire in native versus exotic Arizona grassland. *Southwestern Naturalist*. 37(1): 73-81.
- Bock, C.E.; Bock, J.H.; Kenney, W.R.; Hawthorne, V.M. 1984. Responses of birds, rodents, and vegetation to livestock enclosure in a semidesert grassland site. *Journal of Range Management*. 37: 239-242.
- Bock, C.E.; Bock, J.H.; Smith, H.M. 1993. Proposal for a system of Federal livestock enclosures on public rangelands in the western United States. *Conservation Biology*. 7: 731-733.
- Bock, C.E.; Webb, B. 1984. Birds as grazing indicator species in southeastern Arizona. *Journal of Wildlife*

- Management. 48(3): 1045-1049.
- Bock, Carl E. 1969. Intra- vs. interspecific aggression in pygmy nuthatch flocks. *Ecology*. 50(5): 903-905.
- Bock, Carl E.; Lepthien, L.W. 1976. Synchronous eruptions of boreal seed-eating birds. *American Naturalist*. 110: 559-579.
- Bodie, W.L.; Hickey, W.O. 1980. Response of bighorns to a rest-rotation grazing plan in Central Idaho. Biennial symposium, Northern Wild Sheep and Goat Council. 2: 60-69.
- Bohall-Wood, P. 1987. Abundance, habitat use, and perch use of loggerhead shrikes in north central Florida. *Wilson Bulletin*. 99(1): 82-86.
- Boutin, S.C.; Krebs, J.; Sinclair, A.R.E.; Smith, J.N.M. 1986. Proximate causes of losses in a snowshoe hare population. *Canadian Journal of Zoology*. 64: 606-610.
- Braun, Clait E.; Baker, Maurice F.; Eng, Robert L. [and others]. 1976. Conservation committee report on effects of alteration of sagebrush communities on the associated avifauna. *Wilson Bulletin*. 88: 165-171.
- Brawn, J.D.; Balda, R.P. 1988. Population biology of cavity nesters in northern Arizona: do nest sites limit breeding densities? *Condor*. 90(1): 61-71.
- Brennan, L.A. 1990. What happened to the mountain quail in Idaho? *Quail Unlimited*. 9: 42-43, 69.
- Brennan, Leonard A. 1989. Report on a survey of mountain quail in eastern Oregon, southeastern Washington, western Idaho, and northern Nevada. Boise, ID: The Chukar Foundation. 58 p. Unpublished report. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.
- Brennan, L.A.; Block, W.M.; Gutierrez, R.J. 1987. Habitat use by mountain quail in northern California. *Condor*. 89: 66-74.
- Brittall, J.D.; Poelker, R.J.; Sweeny, S.J. [and others]. 1989. Native cats of Washington. Olympia, WA: Washington Department of Wildlife. 169 p. Unpublished report. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.
- Brooks, B.L.; Temple, S.A. 1990. Habitat availability and suitability for loggerhead shrikes in the upper Midwest. *American Midland Naturalist*. 123: 75-83.
- Brown, Herbert A.; Bury, R. Bruce; Darda, David M. [and others]. 1995. Reptiles of Washington and Oregon. Seattle, WA: Seattle Audubon Society. 176 p.
- Bruns, E.H. 1977. Winter behavior of pronghorns in relation to habitat. *Journal of Wildlife Management*. 41(3): 560-571.
- Bull, E.L. 1991. Summer roosts and roosting behavior of Vaux's swifts in old growth forests. *Northwest Naturalist*. 72(2): 78-82.

- Bull, E.L.; Hohmann, J.E. 1993. The association between Vaux's swifts and old growth forests in northeastern Oregon. *Western Birds*. 24(1): 38-42.
- Bull, E.L.; Holthausen, R.S. 1993. Habitat use and management of pileated woodpeckers in northeastern Oregon. *Journal of Wildlife Management*. 57(2): 335-345.
- Bull, E.L.; Holthausen, R.S.; Henjum, M.G. 1992. Roost trees used by pileated woodpeckers in northeastern Oregon. *Journal of Wildlife Management*. 56(4): 786-793.
- Bull, E.L.; Wright, A.L.; Henjum, M.G. 1990. Nesting habitat of flammulated owls in Oregon. *Journal of Raptor Research*. 24(3): 52-55.
- Bull, Evelyn L. 1980. Resource partitioning among woodpeckers in northeastern Oregon. Moscow, ID: University of Idaho. 109 p. Ph.D. dissertation.
- Bull, Evelyn L.; Henjum, Mark G. 1990. Ecology of the great gray owl. Gen. Tech. Rep. PNW-GTR-265. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 39 p.
- Bull, Evelyn L.; Partridge, Arthur D. 1986. Methods of killing trees for use by cavity nesters. *Wildlife Society Bulletin*. 14: 142-146.
- Bull, Evelyn L.; Peterson, Steven R.; Thomas, Jack W. 1986a. Resource partitioning among woodpeckers in northeastern Oregon. Res. Note. PNW-444. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 19 p.
- Bull, Evelyn L.; Thomas, Jack Ward; Horn, Kirk. 1986b. Snag management on National Forests in the Pacific Northwest - 1984. *Western Journal of Applied Forestry*. 1(2): 41-43.
- Bull, Evelyn L.; Twombly, Asa D.; Quigley, Thomas M. 1980. Perpetuating snags in managed mixed conifer forests of the Blue Mountains, Oregon. In: *Proceedings, management of western forests and grasslands for nongame birds; 1980 February 11-14; Salt Lake City, UT*. Gen. Tech. Rep. INT-86. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 325-336.
- Busack, Stephen D.; Bury, R. Bruce. 1974. Some effects of off-road vehicles and sheep grazing on lizard populations in the Mojave Desert. *Biological Conservation* 6: 179-183.
- Buskirk, Stephen W.; Powell, Roger A. 1994. Habitat ecology of American martens and fishers. In: Buskirk, S.W.; Harestad, A.; Raphael, M.; Powell, R.A., eds. *Martens, sables and fishers: biology and conservation*. Ithaca, NY: Cornell University Press: 283-296.
- Butts, T.W. 1992. *Lynx (Felis lynx) biology and management: a literature review and annotated bibliography*. Missoula, MT: U.S. Department of Agriculture, Forest Service, Northern Region 1 vol.
- Cade, B.S.; Hoffman, R.W. 1990. Winter use of Douglas-fir forests by blue grouse in Colorado. *Journal of Wildlife Management*. 54(3): 471-479.
- Cade, B.S.; Hoffman, R.W. 1993. Differential migration of blue grouse in Colorado. *Auk*. 110: 70-77.
- Call, Mayo W. 1979. Habitat requirements and management recommendations for sage grouse. Tech. Note. Denver, CO: U.S. Department of the Interior, Bureau of Land Management. 37 p.

- Carbyn, L.N.; Patriquin, D. 1983. Observations on home range sizes, movements and social organization of lynx (Lynx canadensis) in Riding Mountain National Park, Manitoba. *Canadian Field-Naturalist*. 97: 262-267.
- Carey, A.B. 1995. Sciurids in Pacific Northwest managed and old-growth forests. *Ecological Applications*. 5(3): 648-661.
- Carey, Andrew B. 1991. The biology of arboreal rodents in Douglas-fir forests. Gen. Tech. Rep. PNW-GTR-276. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 46 p. (Huff, Mark H.; Holthausen, Richard S.; Aubry, Keith B., tech. coords. Biology and management of old-growth forests).
- Cassirer, E. Frances; Oldenburg, Lloyd E.; Coggins, Victor L. [and others]. 1996. Overview and preliminary analysis of a bighorn sheep dieoff, Hells Canyon 1995-96. In: Proceedings of the 10th biennial symposium; 1996 Apr. 29-May 3; Silverthorne, CO. Cody, WY: Northern Wild Sheep and Goat Council. 10: 78-86.
- Caton, E. 1996. Effects of fire and salvage logging on the cavity-nesting bird community in northwestern Montana. Missoula, MT: University of Montana. 115 p. Ph.D. dissertation.
- Chadwick, Douglas H. 1973. Mountain goat ecology-logging relationships in the Bunker Creek drainage of western Montana. Missoula, MT: School of Forestry, University of Montana; final report; State of Montana Project No. W-120-R-3, 4. No. 91.01; 228 p. M.S. thesis..
- Chapin, T.G. 1995. Influence of landscape pattern and forest type on use of habitat by marten in Maine. Orono, ME: University of Maine. 131 p. M.S. thesis.
- Chavez-Ramirez, F.; Gawlik, D.E.; Prieto, F.G.; Slack, R.D. 1994. Effects of habitat structure on patch use by loggerhead shrikes wintering in a natural grassland. *Condor*. 96(1): 228-231.
- Christy, R.E.; West, S.D. 1993. Biology of bats in Douglas-fir forests. Gen. Tech. Rep. PNW-GTR-308. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 28 p.
- Chung-MacCoubrey, Alice L. 1996. Bat species composition and roost use in pinyon-juniper woodlands of New Mexico. In: Barclay, Robert M.R.; Brigham, R. Mark, eds. Bats and forests symposium; 1995 October 19-21; Victoria, BC. Victoria, BC: Ministry of Forests Research Program: 118-123.
- Clark, D.R., Jr. 1988. Environmental contaminants and the management of bat populations in the United States. Gen. Tech. Rep. RM-166. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 409-413.
- Clark, H.W. 1935. Fire and bird populations. *Condor*. 37: 16-18.
- Clark, Richard J. 1975. A field study of the short-eared owl, Asio flammeus (Pontoppidan), in North America. *Wildlife Monographs*. 47: 1-67.
- Clarkson, D.A.; Mills, L.S. 1994. Hypogeous sporocarps in forest remnants and clearcuts in southwest Oregon. *Northwest Science*. 68(4): 259-265.
- Clemente, F.; Valdez, R.; Holechek, J.L. [and others]. 1995. Pronghorn home range relative to permanent water in southern New Mexico. *Southwestern Naturalist*. 40(1): 38-41.

- Coffin, K.W.; Kujala, Q.J.; Douglass, R.J.; Irby, L.R. 1997. Interactions among marten prey availability, vulnerability, and habitat structure. In: Prouls, G.; Gryant, H.N.; Woodard, P.M., eds. *Martes: taxonomy, ecology, techniques, and management: Proceedings of the 2d international Martes symposium*; [dates of meeting unknown]; [meeting location unknown]. Edmonton, AB: The Provincial Museum of Alberta: 199-211.
- Coggins, V.L. 1988. The Lostine Mountain bighorn sheep die-off and domestic sheep. In: Proceedings of the 6th biennial symposium; 1988 Apr. 11-15; Banff, Alberta. Cody, WY: Northern Wild Sheep and Goat Council. 6: 66-76.
- Coggins, Victor L.; Matthews, Patrick E.; Van Dyke, Walt. 1996. History of transplanting mountains goats and mountain sheep - Oregon. In: Proceedings of the 10th biennial symposium; 1996 Apr. 29 - May 3; Silverthorne, CO. Cody, WY: Northern Wild Sheep and Goat Council. 10: 190-197.
- Connelly, J.W., Jr. 1982. An ecological study of sage grouse in southeastern Idaho. Pullman, WA: Washington State University. 84 p. Ph.D. dissertation.
- Connelly, J.W.; Wakkinen, W.L.; Apa, A.D.; Reese, K.P. 1991. Sage grouse use of nest sites in southeastern Idaho. *Journal of Wildlife Management*. 55: 521-524.
- Cooper, J.G. 1868. The fauna of Montana territory. *American Naturalist*. 2: 528-538.
- Copeland, Jeffrey P. 1996. Biology of the wolverine in central Idaho. Moscow, ID: University of Idaho. 138 p. M.S. thesis.
- Corn, J.G.; Raphael, M.G. 1992. Habitat characteristics at marten subnivean access sites. *Journal of Wildlife Management*. 56: 442-448.
- Cornely, John E.; Carraway, L.N.; Verts, B.J. 1992. *Sorex preblei*. *Mammalian species*. No. 416: American Society of Mammalogists: 1-3.
- Coulombe, H.N. 1971. Behavior and population ecology of the burrowing owl, *Speotyto cunicularia*, in the Imperial Valley of California. *Condor*. 73(2): 162-176.
- Craig, R.B. 1978. An analysis of the predatory behavior of the loggerhead shrike. *Auk*. 95(2): 221-234.
- Craig, T.H.; Trost, C.H. 1979. The biology and nesting density of breeding American kestrels and long-eared owls on the Big Lost River, southeastern Idaho. *Wilson Bulletin*. 91(1): 50-61.
- Craighead, J.J.; Craighead, F.C. 1956. Hawks, owls, and wildlife. Harrisburg, PA: Stackpole Company. 443 p.
- Crawford, J.A.; Lutz, R.S. 1985. Sage grouse population trends in Oregon, 1941-1983. *Murrelet*. 66(3): 69-74.
- Crocker-Bedford, D.C. 1990. Goshawk reproduction and forest management. *Wildlife Society Bulletin*. 18: 262-269.
- Daubenmire, R. 1970. Steppe vegetation of Washington. Bull. 62. Pullman, WA: State of Washington, Agricultural Experiment Station. 131 p.
- Dawson, William R. 1997. Pine siskin (*Carduelis pinus*). In: Poole, Alan; Gill, Frank B., eds. *The birds of*

- North America. No. 280: Philadelphia, PA: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 24 p.
- Deblinger, R.D.; Alldredge, A.W. 1991. Influence of free water on pronghorn distribution in a sagebrush/steppe grassland. *Wildlife Society Bulletin*. 19(3): 321-326.
- DeGraaf, R.M.; Scott, V.E.; Hamre, R.H. [and others]. 1991. Forest and rangeland birds of the United States: natural history and habitat use. *Agric. Handb.* 688. Washington, DC: U.S. Department of Agriculture. 625 p.
- Delany, M.F.; Moore, C.T.; Progulske, D.R., Jr. 1995. Territory size and movements of Florida grasshopper sparrows. *Journal of Field Ornithology*. 66: 305-309.
- Demarchi, R.A. 1965. An ecological study of the Ashnola bighorn winter ranges. Vancouver, BC: University of British Columbia. 103 p. M.S. thesis.
- DeSante, D.F.; Pyle, P. 1986. Distributional checklist of North American birds. Volume 1: United States and Canada. Lee Vining, CA: Artemisia Press.
- DeStefano, Stephen. 1990. Investigation of the status of kit foxes in southeastern Oregon. Tech. Rep. 90-5-01. Portland, OR: Oregon Department of Fish and Wildlife, Wildlife Diversity Program. 60 p.
- Diem, K.L.; Zeveloff, S.I. 1980. Ponderosa pine bird communities. In: DeGraaf, R.M.; Tilghman, N.G., eds. *Management of western forests and grasslands for nongame birds*. Gen Tech. Rep. INT-86. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 170-197.
- Diller, Lowell V.; Johnson, Donald R. 1982. Ecology of reptiles in the Snake River Birds of Prey Area. Boise, ID: U.S. Department of the Interior, Bureau of Land Management, Snake River Birds of Prey Research Project. 107 p.
- Dixon, R.; Saab, V. [In prep.]. Black-backed woodpecker. In: Poole, A.; Gill, F., eds. *The birds of North America*. Washington, DC: American Ornithologists' Union. Bimonthly. Issued jointly: Washington, DC: The American Ornithologists' Union; Philadelphia, PA: The Academy of Natural Sciences of Philadelphia. ISSN: 1061-5466.
- Dobkin, D.S.; Gettinger, R.D.; Gerdes, M.G. 1995. Springtime movements, roost use, and foraging activity of Townsend's big-eared bat (*Plecotus townsendii*) in central Oregon. *Great Basin Naturalist*. 55(4): 315-321.
- Dobler, F.C.; Eby, J.; Richardson, S.; Vander Haegen, M. 1996. Status of Washington's shrub-steppe ecosystem: extent, ownership, and wildlife/vegetation relationships. Res. Rep. Olympia, WA: Washington Department of Fish and Wildlife.
- Dodge, W.E. 1977. Status of the fisher (*Martes pennanti*) in the conterminous United States. U.S. Department of the Interior. Unpublished report. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.
- Doyle, F.I.; Smith, J.M.N. 1994. Population responses of northern goshawks to the 10-year cycle in numbers of showshoe hares. *Studies in Avian Biology*. 16: 122-129.
- Doyle, Frank I.; Smith, James M.N. 1994. Population responses of northern goshawks to the 10-year

- cycle in numbers of snowshoe hares. The northern goshawk: ecology and management: Proceedings of a symposium of the Cooper Ornithological Society; 1993 April 14-15; Sacramento, CA. In: *Studies in Avian Biology*. Cooper Ornithological Society. 16: 122-129.
- Duncan, J.R. 1992. Influence of prey abundance and snow cover on great gray owl breeding dispersal. Winnipeg, MB: University of Manitoba. 127 p. Ph.D. dissertation.
- Duncan, James R.; Hayward, Patricia H. 1994. Current management situation: great gray owls. In: Hayward, G.D.; Verner, J., tech. eds. 1994. Flammulated, boreal, and great gray owls in the United States: a technical conservation assessment. Gen. Tech. Rep. RM-253. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 159-175.
- Dunning, J.B., Jr.; Bowers, R.K., Jr. 1990. Lethal temperatures in ash-throated flycatcher nests located in metal fence poles. *Journal of Field Ornithology*. 61(1): 98-103.
- Eberhardt, L.L.; Knight, R.R. 1996. How many grizzlies in Yellowstone? *Journal of Wildlife Management*. 60(2): 416-421.
- Edminster, F.C. 1954. American game birds of field and forest. Their habits, ecology and management. New York: Charles Scribner's Sons. 490 p.
- Egoscue, Harold J. 1962. Ecology and life history of the kit fox in Tooele County, Utah. *Ecology*. 43(3): 481-497.
- Ehrlich, Paul R.; Dobkin, David S.; Wheye, Darryl. 1988. *The birder's handbook*. New York: Simon and Schuster. 785 p.
- Erickson, Janet L.; West, Stephen D. 1996. Managed forests in the western Cascades: the effects of seral stage on bat habitat use patterns. In: Barclay, Robert M.R.; Brigham, R. Mark, eds. *Bats and forests symposium*; 1995 Oct 19-21; Victoria, BC. Victoria, BC: Ministry of Forests Research Program: 215-227.
- Evans, H.F. 1960. A preliminary investigation of caribou in the northwestern United States. Bozeman, MT: Montana State University. 145 p. M.S. thesis.
- Evans, Raymond A.; Young, James A. 1978. Effectiveness of rehabilitation practices following wildfire in a degraded big sagebrush-downy brome community. *Journal of Range Management*. 31: 185-188.
- Feldhamer, G.A.; Klann, R.S.; Gerard, A.S.; Driskell, A.C. 1993. Habitat partitioning, body size, and timing of parturition in pygmy shrews and associated soricids. *Journal of Mammology*. 74(2): 403-411.
- Fischer, R.A. 1994. Sage grouse ecology final report. Boise, ID: Idaho Department of Fish and Game; job completion report; W-160-R-19; subproject 9. 91 p.
- Flickinger, E.L. 1995. Loggerhead fatalities on a highway in Texas. In: Yosef, R.; Lohrer, F.E., eds. *Shrikes (Laniidae) of the world: biology and conservation: 1st international shrike symposium*; 1993 January 11-15; Lake Placid, FL. In: *Proceedings of the Western Foundation of Vertebrate Zoology*; 6(1): 67-69.
- Flinn, P. 1956. Caribou of Idaho. Boise, ID: Idaho Department of Fish and Game. Unpublished report. On file with: Boise, ID: Idaho Department of Fish and Game, P.O. Box 25, Boise, ID 83707. 79 p.

- Foreyt, William J. 1994. Effects of controlled contact exposure between healthy bighorn sheep and llamas, domestic goats, mountain goats, cattle, domestic sheep, or mouflon sheep. Proceedings of the 9th biennial symposium. Northern Wild Sheep and Goat Council. 1994 May 2-6. Cranbrook, British Columbia. Thermopolis, WY: Northern Wild Sheep and Goat Council. 9: 7-14. Proceedings available from: Foundation for North American Wild Sheep, 720 Allen Avenue, Cody, WY 82414.
- Foreyt, William J.; Jessup, David A. 1982. Fatal pneumonia of bighorn sheep following association with domestic sheep. *Journal of Wildlife Diseases*. 18: 163-168.
- Fortin, C.; Cantin, M. 1994. The effects of trapping on a newly exploited American marten population. In: Buskirk, S.W.; Harestad, A.S.; Raphael, M.G.; Powell, R.A., eds. *Martens, sables, and fishers: biology and conservation*. Ithaca, NY: Cornell University Press: 179-191.
- Frederick, Glenn P. 1991. Effects of forest roads on grizzly bears, elk, and gray wolves: a literature review. Publ. R1-91-73. Libby, MT: Kootenai National Forest. 53 p.
- French, N.R. 1959. Life history of the black rosy finch. *Auk*. 76(2): 159-180.
- Friedmann, H.; Kiff, L.F. 1985. The parasitic cowbirds and their hosts. *Proceedings of the Western Foundation of Vertebrate Zoology*. 2(4): 227-266.
- Friedmann, J. 1929. *The cowbirds, a study in the biology of social parasitism*. Springfield, IL: C.C. Thomas. 421 p.
- Fritts, Steven H.; Mech, L. David. 1981. Dynamics, movements, and feeding ecology of a newly protected wolf population in northwestern Minnesota. *Wildlife Monographs No. 80*. 79 p.
- Fritts, Steven H.; Paul, William J.; Mech, L. David. 1985. Can relocated wolves survive? *Wildlife Society Bulletin*. 13(4): 459-463.
- Fuller, Todd K.; Keith, Lloyd B. 1980. Wolf population dynamics and prey relationships in northeastern Alberta. *Journal of Wildlife Management*. 44(3): 583-602.
- Galen, Christie. 1989. A preliminary assessment of the status of the Lewis' woodpecker in Wasco County, Oregon. Tech. Rep. 88-3-01. Portland, OR: Oregon Department of Fish and Wildlife, Nongame Wildlife Program.
- Garrett, K.L.; Raphael, M.G.; Dixon, R.D. 1996. White-headed woodpecker. In: Poole, A.; Gill, F., eds. *The birds of North America*. No. 252. Philadelphia, PA: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 24 p.
- Gawlik, D.E.; Bildstein, K.L. 1993. Seasonal habitat use and abundance of loggerhead shrikes in South Carolina. *Journal of Wildlife Management*. 57(2): 352-357.
- Genter, D.L. 1986. Wintering bats of the upper Snake River plain: occurrence in lava-tube caves. *Great Basin Naturalist*. 46: 241-244.
- Gerber, Mark F.; Munger, James C.; Ames, Aaron; Stewart, Jeff. 1997. Reptiles and amphibians in deep canyons: the Big Jacks and Little Jacks Creek drainages of Owyhee County, Idaho. Tech. Bull. 97-1. Boise, ID: U.S. Department of the Interior, Bureau of Land Management, Idaho State Office. 14 p.
- Getz, L.L. 1961. Hunting areas of the long-eared owl. *Wilson Bulletin*. 73(1): 79-82.

- Giesen, K.M.; Connelly, J.W. 1993. Guidelines for management of Columbian sharp-tailed grouse habitats. *Wildlife Society Bulletin*. 21(3): 325-333.
- Gleason, R.S.; Johnson, D.R. 1985. Factors influencing nesting success of burrowing owls (*Athene cunicularia*) in southeastern Idaho. *Great Basin Naturalist*. 45(1): 81-84.
- Goggans, R. 1986. Habitat use by flammulated owls in northeastern Oregon. Corvallis, OR: Oregon State University. 54 p. M.S. thesis.
- Goggans, Rebecca; Dixon, Rita D.; Seminara, Claire. 1988. Habitat use by three-toed and black-backed woodpeckers, Deschutes National Forest, Oregon. Nongame Rep. 87-3-02. Portland, OR: Oregon Department of Fish and Wildlife; Deschutes National Forest, Nongame Wildlife Program. 49 p.
- Golightly, R.T.; Ohmart, R.D. 1984. Metabolism and body temperature of two desert canids: coyotes and kit foxes. *Journal of Mammalogy*. 64(4): 624-635.
- Gray, R.H.; Rickard, W.H. 1989. The protected area of Hanford as a refugium for native plants and animals. *Environmental Conservation*. 16: 215-216, 251-262.
- Green, G.A.; Anthony, R.G. 1989. Nesting success and habitat relationships of burrowing owls in the Columbia Basin, Oregon. *Condor*. 91(2): 347-354.
- Green, Gerald I.; Mattson, David J.; Peek, James M. 1997. Spring feeding on ungulate carcasses by grizzly bears in Yellowstone National Park. *Journal of Wildlife Management*. 61(4): 1040-1055.
- Gregg, M.A. 1991. Use and selection of nesting habitat by sage grouse in Oregon. Corvallis, OR: Oregon State University. M.S. thesis.
- Griffin, Daude N. 1959. The poisoning of meadowlarks with insecticides. *Wilson Bulletin*. 71(2): 193.
- Grubb, T.C., Jr.; Yosef, R. 1994. Habitat-specific nutritional condition in loggerhead shrikes (*Lanius ludovicianus*): evidence from ptilochronology. *Auk*. 111(3): 756-759.
- Hall, E.R. 1928. Notes on the life history of the sagebrush meadow mouse (*Lagurus*). *Journal of Mammalogy*. 9: 201-204.
- Hamer, D.; Herrero, S. 1987. Grizzly bear food and habitat in the Front Ranges of Banff National Park, Alberta. *International Conference on Bear Research and Management*. 7:199-214.
- Hamilton, K.S.; Holl, S.K.; Douglas, C.L. 1982. An evaluation of the effects of recreational activity on bighorn sheep in the San Gabriel Mountains, California. *Transactions of the annual meeting of the Desert Bighorn Council*. 26: 50-55.
- Hanley, T.A.; Page, J.L. 1982. Differential effects of livestock use on habitat structure and rodent populations in Great Basin communities. *California Fish and Game*. 68(3): 160-174.
- Hann, Wendel J.; Jones, Jeffrey L.; Karl, Michael G. [and others]. 1997. Landscape dynamics of the basin. In: Quigley, Thomas M.; Arbelbide, Sylvia J., tech. eds. An assessment of ecosystem components in the interior Columbia basin and portions of the Klamath and Great Basins. Gen. Tech. Rep. PNW-GTR-405. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 337-1055. Chapter 3.

- Hargis, C.D. 1996. The influence of forest fragmentation and landscape pattern on American martens and their prey. Logan, UT: Utah State University. 154 p. Ph.D. dissertation.
- Hargis, Christina D.; McCarthy, Clinton; Perloff, Richard D. 1994. Home ranges and habitats of northern goshawks in eastern California. The northern goshawk: ecology and management: Proceedings of a symposium of the Cooper Ornithological Society; 1993 April 14-15; Sacramento, CA. In: Studies in Avian Biology. Cooper Ornithological Society; 16: 66-75.
- Harper, Kimball T.; Pendleton, Rosemary L. 1993. Cyanobacteria and cyanolichens: Can they enhance availability of essential minerals for higher plants? Great Basin Naturalist. 53: 59-72.
- Harrington-Tweit, Bill; Mattocks, Philip. W., Jr. 1985. The nesting season June 1-July 31: Northern Pacific Coast Region. American Birds. 39(5): 953-956.
- Hash, H.S. 1987. Wolverine. In: Novak, M.; Baker, J.A.; Obbard, M.E., eds. Wild furbearer management and conservation in North America. Toronto: Ontario Ministry of Natural Resources: 575-585.
- Haug, E.A.; Oliphant, L.W. 1990. Movements, activity patterns, and habitat use of burrowing owls in Saskatchewan. Journal of Wildlife Management. 54(1): 27-35.
- Hayward, G.D.; Escano, R.E. 1989. Goshawk nest-site characteristics in western Montana and northern Idaho. Condor. 91(2): 476-479.
- Hayward, G.D.; Hayward, P.H.; Garton, E.O. 1993. Ecology of boreal owls in the northern Rocky Mountains. Wildlife Monographs. 124: 1-59.
- Hayward, Gregory D. 1994a. Conservation status of boreal owls in the United States. In: Hayward, G.D.; Verner, J., tech. eds. 1994. Flammulated, boreal, and great gray owls in the United States: a technical conservation assessment. Gen. Tech. Rep. RM-253. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 139-153.
- Hayward, Gregory D. 1994b. Conservation status of great gray owls in the United States. In: Hayward, G. D, and Verner, J., tech. eds. 1994. Flammulated, boreal, and great gray owls in the United States: a technical conservation assessment. Gen. Tech. Rep. RM-253. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 202-206.
- Hayward, Gregory D. 1994c. Review of technical knowledge: boreal owls. In: Hayward, G.D.; Verner, J., tech. eds. 1994. Flammulated, boreal, and great gray owls in the United States: a technical conservation assessment. Gen. Tech. Rep. RM-253. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 92-127.
- Hayward, Gregory D. 1997. Forest management and conservation of boreal owls in north America. Journal of Raptor Research. 31(2):114-124.
- Heinemeyer, K.S. 1995. Conservation assessment for fisher (*Martes pennanti*) in Idaho. In: Idaho Department of Fish and Game, Nez Perce Tribe; Sawtooth National Forest. Forest carnivores in Idaho-Habitat conservation assessments (HCAs) and conservation strategies (CS). Idaho State Conservation Effort; Idaho Department of Fish and Game, Idaho Department of Parks and Recreation; Bureau of Land Management; U.S. Department of Agriculture, Forest Service, Northern Region and Intermountain Region; U.S. Department of the Interior, Fish and Wildlife Service: 28-41. Draft report dated April 21, 1995. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project,

USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.

- Hejl, S.J. 1994. Human-induced changes in bird populations in coniferous forests in western North America during the past 100 years. In: Studies in Avian Biology. Cooper Ornithological Society; 15: 232-246.
- Helms, B. 1978. Antelope and energy development. In: Proceedings of the 8th biennial pronghorn antelope workshop; 1978 May 2-4; Jasper, AB. Jasper, AB: Alberta Recreation, Parks and Wildlife, Fish and Wildlife Division: 206-215.
- Henny, C.J.; Blus, L.J.; Kaiser, T.E. 1984. Heptachlor seed treatment contaminates hawks, owls, and eagles of Columbia Basin, Oregon. Raptor Research. 18: 41-48.
- Herkert, J.R. 1994. The effects of habitat fragmentation on Midwestern grassland bird communities. Journal of Ecological Applications. 4: 461-471.
- Hessburg, Paul F.; Smith, Bradley G.; Kreiter, Scott D. [and others]. 1999. Historical and current forest and range landscapes in the interior Columbia River basin and portions of the Klamath and Great Basins. Part I: Linking vegetation patterns and landscape vulnerability to potential insect and pathogen disturbances. Gen. Tech. Rep. PNW-GTR-458. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 357 p. (Quigley, Thomas M., tech. ed.; Interior Columbia Basin Ecosystem Management Project: scientific assessment).
- Hicks, Lorin L.; Elder, James M. 1979. Human disturbance of Sierra Nevada bighorn sheep. Journal of Wildlife Management. 43(4): 909-915.
- Hines, James E. 1986. Social organization, movements, and home ranges of blue grouse in fall and winter. Wilson Bulletin. 98: 419-432.
- Hjeljord, O. 1973. Mountain goat forage and habitat preference in Alaska. Journal of Wildlife Management. 37(3): 353-362.
- Hodgman, T.P.; Harrison, D.J.; Katnik, D.D.; Elowe, K.D. 1994. Survival in an intensively trapped marten population in Maine. Journal of Wildlife Management. 58(4): 593-600.
- Hoffman, N. 1997. Distribution of *Picoides* woodpeckers in relation to habitat disturbance within the Yellowstone area. Bozeman, MT: Montana State University. 74 p. M.S. thesis.
- Hoffman, Stephen W.; DeRagon, William R.; Bednarz, James C. 1992. Patterns and recent trend in counts of migrant hawks in western North America. Washington, DC: U.S. Fish and Wildlife Service, Office of Migratory Bird Management. Unpublished report. Albuquerque, NM: Hawkwatch International. On file with: Hawkwatch International, P.O. 35706, Suite 206, Albuquerque, NM 87176.
- Hogstad, O. 1976. Sexual dimorphism and divergence in winter foraging behaviour of three-toed woodpeckers, *Picoides tridactylus*. Ibis. 118(1): 41-50.
- Holt, Denver W. 1997. The long-eared owl (*Asio otus*) and forest management: a review of the literature. Journal of Raptor Research. 32: 175-186.
- Horning, John. 1994. Grazing to extinction: endangered, threatened and candidate species imperiled by livestock grazing on western public lands. Washington, DC: National Wildlife Federation. 68 p.

- Hornocker, M.G.; Hash, H.S. 1981. Ecology of the wolverine in northwestern Montana. *Canadian Journal of Zoology*. 59(7): 1286-1301.
- Hosten, P.E.; West, N.E. 1994. Cheatgrass dynamics following wildfire on a sagebrush semidesert site in central Utah. In: Monsen, Stephen B.; Kitchen, Stanley G., eds. *Proceedings-ecology and management of annual rangelands; 1992 May 18-21; Boise, ID. Gen. Tech. Rep. INT-313*. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 56-68.
- Humphrey, S.R. 1975. Nursery roosts and community diversity of nearctic bats. *Journal of Mammalogy*. 56(2): 321-346.
- Humphrey, S.R.; Kunz, T.H. 1976. Ecology of a Pleistocene relict, the western big-eared bat (*Plecotus townsendii*), in the southern Great Plains. *Journal of Mammalogy*. 57: 470-494.
- Hupp, J.W. 1987. Sage grouse resource exploitation and endogenous reserves in Colorado. Fort Collins, CO: Colorado State University. 73 p. Ph.D. dissertation.
- Hutto, R.L. 1995. Composition of bird communities following stand-replacement fires in northern Rocky Mountain conifer forests. *Conservation Biology*. 9(5): 1041-1058.
- Idaho State Conservation Effort. 1995. Habitat conservation assessment and conservation strategy for the Townsend's big-eared bat. Draft unpublished report no. 1. Boise, ID: Idaho Department of Fish and Game, P.O. Box 25, Boise, ID 83707. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.
- Interagency Grizzly Bear Committee. 1986. Interagency grizzly bear guidelines. 99 p. Unpublished report. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.
- Interagency Grizzly Bear Committee. 1998. Grizzly update: news on grizzly bear recovery activities in the states of Wyoming, Montana, Idaho, and Washington from the Interagency Grizzly Bear Committee. 12 p. Unpublished report. On file with: Missoula, MT: U.S. Department of the Interior, Fish and Wildlife Service, Interagency Grizzly Bear Committee, University Hall, Room 309, University of Montana, Missoula, MT 59812.
- Interior Columbia Basin Ecosystem Management Project (ICBEMP). 1996a. Expert panel notes for amphibians and reptiles terrestrial habitat assessment, fall 1996, for the Interior Columbia Basin. Unpublished notes. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.
- Interior Columbia Basin Ecosystem Management Project (ICBEMP). 1996b. Expert panel notes for amphibian and reptile viability assessment, spring 1996, for the interior Columbia basin. Unpublished notes. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.
- Interior Columbia Basin Ecosystem Management Project (ICBEMP). 1996c. Expert panel notes for bats and small mammals terrestrial habitat assessment, fall 1996, for the interior Columbia basin. Unpublished notes. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of

the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.

Interior Columbia Basin Ecosystem Management Project (ICBEMP). 1996d. Expert panel notes for bats and small mammals viability assessment, spring 1996, for the interior Columbia basin. Unpublished notes. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.

Interior Columbia Basin Ecosystem Management Project (ICBEMP). 1996e. Expert panel notes for carnivore terrestrial habitat assessment, fall 1996, for the interior Columbia basin. Unpublished notes. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.

Interior Columbia Basin Ecosystem Management Project (ICBEMP). 1996f. Expert panel notes for mammalian carnivore and ungulate viability assessment, spring 1996, for the interior Columbia basin. Unpublished notes. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.

Interior Columbia Basin Ecosystem Management Project (ICBEMP). 1996g. Expert panel notes for passerine and other birds viability assessment, spring 1996, for the interior Columbia basin. Unpublished notes. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.

Interior Columbia Basin Ecosystem Management Project (ICBEMP). 1996h. Expert panel notes for raptor and upland game bird viability assessment, spring 1996, for the interior Columbia basin. Unpublished notes. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.

Interior Columbia Basin Ecosystem Management Project (ICBEMP). 1996i. Expert panel notes for ungulate terrestrial habitat assessment, fall 1996, for the interior Columbia basin. Unpublished notes. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.

Janes, S.W. 1983. Status, distribution, and habitat selection of the grasshopper sparrow in Morrow County, Oregon. *Murrelet*. 64: 51-54.

Jasikoff, Thomas M. 1982. Habitat suitability index models: ferruginous hawk. FWS/OBS-82/10.10. Washington, DC: U.S. Department of the Interior, Fish and Wildlife Service. 18 p.

Jensen, W.R.; Fuller, T.K.; Robinson, W. L. 1986. Wolf, *Canis lupus*, distribution on the Ontario-Michigan border near Sault Ste. Marie. *Canadian Field-Naturalist*. 100: 363-366.

Jewett, S.G.; Taylor, W.P.; Shaw, W.T.; Aldrich, J.W. 1953. *Birds of Washington State*. Seattle: University of Washington Press. 767 p.

Johansen, Jeffrey R.; Ashley, John; Rayburn, William R. 1993. Effects of range fire on soil algal crusts in

- semiarid shrub-steppe of the lower Columbia Basin and their subsequent recovery. *Great Basin Naturalist*. 53: 73-88.
- Johnson, D.H.; Igl, L.D. 1995. Contributions of the Conservation Reserve Program to populations of breeding birds in North Dakota. *Wilson Bulletin*. 107: 709-718.
- Johnson, R.E. 1977. An historical analysis of wolverine abundance and distribution in Washington, USA. *Murrelet*. 58: 13-16.
- Johnson, Rolf L. 1983. Mountain goats and mountain sheep of Washington. *Biol. Bull. No. 18*. Olympia, WA: Washington State Game Department. 196 p.
- Jones, Jeffrey L. 1991. Habitat use of fisher in north central Idaho. Moscow, ID: University of Idaho. 147 p. M.S. thesis.
- Jones, Jeffrey L.; Garton, Edward O. 1994. Selection of successional stages by fishers in north-central Idaho. In: Buskirk, S.W., Harestad, A.; Raphael, M.; Powell, R.A., eds. *Martens, sables, and fishers: biology and conservation*. Ithaca, NY: Cornell University: 377-387.
- Jonkel, Charles J.; Cowan, Ian McT. 1971. The black bear in the spruce-fir forest. *Wildlife Monograph*. No. 27. 57 p.
- Joslin, G. 1986. Mountain goat population changes in relation to energy exploration along Montana's Rocky Mountain Front. In: *Proceedings of the 5th biennial symposium. Northern Wild Sheep and Goat Council; 1986 Apr. 14-17; Missoula, MT. Thermopolis, WY: Northern Wild Sheep and Goat Council. 5:253-271. Proceedings available from: Foundation for North American Wild Sheep, 720 Allen Avenue, Cody, WY 82414.*
- Kaltenecker, J.H. 1997. The recovery of microbiotic crusts following postfire rehabilitation on rangelands of the western Snake River plain. Boise, ID: Boise State University. 99 p. M.S. thesis.
- Kaltenecker, J.; Wicklow-Howard, M. 1994. Microbiotic soil crusts in sagebrush habitats of southern Idaho. 48 p. (plus an appendix). Prepared for Eastside Ecosystem Management Project. Unpublished report. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.
- Kasworm, Wayne F.; Manley, T.L. 1990. Road and trail influences on grizzly bears and black bears in northwest Montana. *International Conference on Bear Research and Management*. 8: 79-84.
- Keister, George P.; Immell, David. 1994. Continued investigations of kit fox in southeastern Oregon and evaluation of status. *Tech. Rep. 94-5-01*. Portland, OR: Oregon Department of Fish and Wildlife, Wildlife Diversity Program. 33 p.
- Kennedy, Ann C. 1994. Biological control of annual grass weeds. In: Monsen, Stephen B.; Kitchen, Stanley G., eds. *Proceedings, ecology and management of annual rangelands; 1992 May 18-21; Boise, ID. Gen. Tech. Rep. INT-GTR-313*. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 186-189.
- Kennedy, Patricia L.; Ward, Johanna M.; Rinker, George A.; Gessaman, James A. 1994. Post-fledging areas in northern goshawk home ranges. *The northern goshawk: ecology and management: Proceedings of a symposium of the Cooper Ornithological Society; 1993 April 14-15; Sacramento, CA.*

- In: Studies in Avian Biology. Cooper Ornithological Society; 16: 75-82.
- Kie, J.G. 1991. Wildlife and livestock grazing alternatives in the Sierra Nevada. In: Transactions of the Western Section of the Wildlife Society; 1991 February 7-9; Sacramento, CA. Sacramento, CA: The Western Section of the Wildlife Society. 27: 17-29.
- Kieth, L.B.; Cary, J.R.; Rongstad, O.J.; Brittingham, M.C. 1984. Demography and ecology of a declining snowshoe hare population. Wildlife Monographs. 90. 43 p.
- Klebenow, D.A. 1969. Sage grouse nesting and brood habitat in Idaho. Journal of Wildlife Management. 33: 649-661.
- Klemmedson, James O.; Smith, Justin G. 1964. Cheatgrass (Bromus tectorum L.). The Botanical Review. 30: 226-262.
- Knapton, R.W. 1979. Optimal size of territory in the clay-colored sparrow, Spizella pallida. Canadian Journal of Zoology. 57(7): 1358-1370.
- Knapton, Richard W. 1994. Clay-colored sparrow (Spizella pallida). In: Poole, A.; Gill, F., eds. The birds of North America. No. 120. Philadelphia, PA: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 16 p.
- Knick, S.T.; Rotenberry, J.T. 1995. Landscape characteristics of fragmented shrubsteppe habitats and breeding passerine birds. Conservation Biology. 9(5): 1059-1071.
- Knick, Steven T.; Kasworm, Wayne. 1989. Shooting mortality in small populations of grizzly bears. Wildlife Society Bulletin. 17(1): 11-15.
- Knight, R.R.; Blanchard, B.M.; Eberhardt, L.L. 1988. Mortality patterns and population sinks for Yellowstone grizzly bears, 1973-1985. Wildlife Society Bulletin. 16:121-125.
- Koehler, C.E.; Barclay, R.M.R. 1988. The potential for vocal signatures in the calls of young hoary bats (Lasiurus cinereus). Canadian Journal of Zoology. 66(9): 1982-1985.
- Koehler, G.M. 1990. Population and habitat characteristics of lynx and snowshoe hares in north central Washington. Canadian Journal of Zoology. 68: 845-851.
- Koehler, G.M.; Brittell, J.D. 1990. Managing spruce fir habitat for lynx and snowshoe hares. Journal of Forestry. 88: 10-14.
- Koehler, Gary M; Aubry, Keith B. 1994. Lynx. In: Ruggiero, L.; Aubry, K.; Buskirk, S. [and others], tech. eds. The scientific basis for conserving forest carnivores: American marten, fisher, lynx and wolverine in the Western United States. Gen. Tech. Rep. RM-254. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 74-98. Chapter 4.
- Koplin, James R. 1969. The numerical response of woodpeckers to insect prey in a subalpine forest in Colorado. Condor. 71(4): 436-438.
- Kovalchik, B.L.; Elmore, W. 1992. Effects of cattle grazing systems on willow-dominated plant associations in central Oregon. In: Symposium on ecology and management of riparian shrub communities; 1991 May 29-31; Sun Valley, ID. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 111-119.

- Kuck, L. 1977. The impact of hunting on Idaho's Pahsimeroi mountain goat herd. In: Samuel, W.; MacGregor, W.G., eds. Proceedings First International Mountain Goat Symposium. 1: 114-125.
- Lanyon, W. E. 1994. Western meadowlark (*Sturnella neglecta*). In: Poole, A.; Gill, F., eds. The birds of North America. No. 104. Philadelphia, PA: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 20 p.
- Lauer, J.L.; Peek, J.M. 1976. Big game-livestock relationships on the bighorn sheep winter range, East Fork Salmon River, Idaho. Bull. No. 12. Moscow, ID: College of Forestry, Wildlife, and Range Science. 44 p.
- Lee, Danny C.; Sedell, James R.; Rieman, Bruce E. [and others]. 1997. Broadscale assessment of aquatic species and habitats. In: Quigley, Thomas M.; Arbelbide, Sylvia J., tech. eds. An assessment of ecosystem components in the interior Columbia basin and portions of the Klamath and Great Basins. Gen. Tech. Rep. PNW-GTR-405. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 1057-1496. Vol. 3. (Quigley, Thomas M., tech. ed. Interior Columbia Basin Ecosystem Management Project: scientific assessment).
- LeFranc, Maurice N. Jr.; Moss, Mary Beth; Patnode, Kathleen; Sugg, William C. III, eds. 1987. Grizzly bear compendium. Washington, DC: Interagency Grizzly Bear Committee and the National Wildlife Federation. 540 p.
- Lehmkuhl, John F.; Raphael, Martin G.; Holthausen, Richard S. [and others]. 1997. Historical and current status of terrestrial species and the effects of proposed alternatives. In: Quigley, Thomas M.; Lee, Kristine M.; Arbelbide, Sylvia J., tech. eds. Evaluation of EIS alternatives by the Science Integration Team. Gen. Tech. Rep. PNW-GTR-406. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 537-730. Vol. 2.
- Leopold, A.S.; Gutierrez, R.J.; Bronson, M.T. 1981. North American game birds and mammals. New York: Charles Scribner's Sons. 198 p.
- Lester, Amy N. 1980. Numerical response of woodpeckers and their effect on mortality of mountain pine beetles in lodgepole pine in northwestern Montana. Missoula, MT: University of Montana. 103 p. M.S. thesis.
- Leu, Matthias. 1995. The feeding ecology and the selection of nest shrubs and fledgling roost sites by loggerhead shrikes (*Lanius ludovicianus*) in the shrub-steppe habitat. Seattle: University of Washington. 90 p. M.S. thesis.
- Linden, H.; Wikman, M. 1983. Goshawk predation on tetraonids: availability of prey and diet of the predator in the breeding season. *Journal of Animal Ecology*. 52: 953-968.
- Linder, K.A. 1994. Habitat utilization and behavior of nesting Lewis' woodpeckers (*Melanerpes lewis*) in the Laramie Range, southeast Wyoming. Laramie, WY: University of Wyoming. 98 p. M.S. thesis.
- Linkhart, B.D. 1984. Range, activity, and habitat use by nesting flammulated owls in a Colorado ponderosa pine forest. Fort Collins, CO: Colorado State University. M.S. thesis.
- Lokemoen, J.T.; Duebbert, H.F. 1976. Ferruginous hawk nesting ecology and raptor populations in northern South Dakota. *Condor*. 78(4): 464-470.
- Lyman, R.L. 1991. Late quaternary biogeography of the pygmy rabbit (*Brachylagus idahoensis*) in

- eastern Washington. *Journal of Mammology*. 72(1): 110-117.
- MacArthur, R.A.; Johnson, R.H.; Geist, V. 1979. Factors influencing heart rate in free-ranging bighorn sheep: a physiological approach to the study of wildlife harassment. *Canadian Journal of Zoology*. 57: 2010-2021.
- MacCracken, J.G.; Alexander, L.E.; Uresk, D.W. 1983. An important lichen of southeastern Montana rangelands. *Journal of Range Management*. 36: 35-37.
- Mace, Richard D.; Waller, John S.; Manley, Timothy L. [and others]. 1996. Relationships among grizzly bears, roads and habitat in the Swan Mountains, Montana. *Journal of Applied Ecology*. 33: 1395-1404.
- Mack, R.N.; Thompson, J. N. 1982. Evolution in steppe with few large, hooved mammals. *American Naturalist*. 119: 757-773.
- Maj, M.; Hillis, M.; Ormiston, J. 1995. Range permit reissuance-flammulated owl. 5 p. Unpublished memo. On file with: U.S. Department of Agriculture, Forest Service, Northern Region, P.O. Box 7669, Missoula, MT 59807.
- Mannan, R.W. 1984. Summer area requirements of pileated woodpeckers in western Oregon. *Wildlife Society Bulletin*. 12(3): 265-268.
- Manning, R.W.; Knox-Jones, J. 1989. *Myotis evotis*. *Mammalian Species*. No. 329. Baltimore, MD: American Society of Mammalogists. 5 p.
- Marcot, Bruce G.; Castellano, Michael A.; Christy, John A. [and others]. 1997. Terrestrial ecology of the basin. In: Quigley, Thomas M.; Arbelbide, Sylvia J., tech. eds. An assessment of ecosystem components in the interior Columbia basin and portions of the Klamath and Great Basins. Gen. Tech. Rep. PNW-GTR-405. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 1497-1713. Vol. 3. (Quigley, Thomas M., tech. ed. Interior Columbia Basin Ecosystem Management Project: scientific assessment).
- Marcot, B.G., B. Wales, and R. Demmer. [In prep.]. Range maps of terrestrial species and communities in the interior Columbia River basin and northern portions of the Klamath and Great basins. Gen. Tech. Rep. PNW-GTR-XXX. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station.
- Marks, J.S. 1986. Nestsite characteristics and reproductive success of long-eared owls in southwestern Idaho. *Wilson Bulletin*. 98(4): 547-560.
- Marks, J.S.; Evans, D.L.; Holt, D. W. 1994. Long-eared owl (*Asio otus*). In: Poole, A.; Gill, F., eds. *The birds of North America*. No. 133. Philadelphia, PA: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 24 p.
- Marks, J.S.; Saab, V. 1987a. Habitat selection by Columbian sharp-tailed grouse in west-central Idaho. Boise, ID: U.S. Department of the Interior, Bureau of Land Management, Boise District. 115 p.
- Marks, J.S.; Saab Marks, V. 1987b. Influence of radio collars on survival of sharp-tailed grouse. *Journal of Wildlife Management*. 51(2): 468-471.
- Marks, J.S.; Saab Marks, V. 1988. Winter habitat use by Columbian sharp-tailed grouse in western Idaho. *Journal of Wildlife Management*. 52(4): 743-746.

- Marshall, D.B.; Chilcote, M.W.; Weeks, H. 1996. Species at risk: sensitive, threatened and endangered vertebrates of Oregon. 2d ed. Portland, OR: Oregon Department Fish and Wildlife. 1 vol.
- Marshall, David B. 1992. Status of the black-backed woodpecker in Oregon and Washington. In: Proceedings of the Blue Mountains biodiversity conference; 1992 May 26-29; Portland, OR. Portland, OR: Portland Audubon Society. 13 p.
- Marshall, J.T., Jr. 1988. Birds lost from a giant sequoia forest during fifty years. *Condor*. 90: 359-372.
- Marti, C.D.; Marks, J.S. 1989. Medium-sized owls. In: Proceedings of the western raptor management symposium and workshop; 1987 October 26-28; Boise, ID. National Wildlife Federation Science and Technical Series. 12: 124-133.
- Martin, Kevin D.; Schommer, Tim; Coggins, Victor. 1996. Literature review regarding the compatibility between bighorn and domestic sheep. In: Proceeding of the 10th biennial symposium. Northern Wild Sheep and Goat Council; 1996 Apr. 29-May 3; Silverthorne, CO. Thermopolis, WY: Northern Wild Sheep and Goat Council. 10: 72-77.
- Martin, S.J. 1983. Burrowing owl occurrence on white-tailed prairie dog colonies. *Journal of Field Ornithology*. 54(4): 422-423.
- Maser, C.; Hammer, E.W.; Brown, C. [and others]. 1974. The sage vole, Lagurus curtatus, in the Crooked River National Grassland, Jefferson County, Oregon: a contribution to its life history and ecology. *Saugetierkundliche Mitteilungen*. 22: 193-222.
- Maser, C.; Strickland, G.S. 1978. The sage vole, Lagurus curtatus, as an inhabitant of subalpine sheep fescue Festuca ovina communities on Steens Mountain: an observation and interpretation. *Northwest Science*. 52: 276-284.
- Matthews, Patrick E.; Coggins, Victor L. 1994. Status and history of mountain goats in Oregon. Proceedings of the 9th biennial symposium. Northern Wild Sheep and Goat Council; 1994 May 2-6; Cranbrook, British Columbia. Thermopolis, WY: Northern Wild Sheep and Goat Council. 9: 69-74. Proceedings available from: Foundation for North American Wild Sheep, 720 Allen Avenue, Cody, WY 82414.
- Mattson, D.J.; Blanchard, B.M.; Knight, R.R. 1991a. Food habits of Yellowstone grizzly bears, 1977-1987. *Canadian Journal of Zoology*. 69: 1619-1629.
- Mattson, D.J.; Reinhart, D.P. 1997. Excavation of red squirrel middens by grizzly bears in the whitebark pine zone. *Journal of Applied Ecology*. 34: 926-940.
- Mattson, David J.; Blanchard, Bonnie M.; Knight, Richard R. 1992. Yellowstone grizzly bear mortality, human habituation, and whitebark pine seed crops. *Journal of Wildlife Management*. 56(3): 432-442.
- Mattson, David J.; Gillin, Colin M.; Benson, Scott A.; Knight, Richard R. 1991b. Bear feeding activity at alpine insect aggregation sites in the Yellowstone ecosystem. *Canadian Journal of Zoology*. 69: 2430-2435.
- Mattson, David J.; Herrero, Stephen; Wright, Gerald; Pease, Craig M. 1996a. Designing and managing protected areas for grizzly bears: how much is enough? In: Wright, R.G., ed. National parks and protected areas : their role in environmental protection. Cambridge, MA: Blackwell Science: 133-164.

- Mattson, David J.; Herrero, Stephen; Wright, Gerald; Pease, Craig M. 1996b. Science and management of Rocky Mountain grizzly bears. *Conservation Biology*. 10(4): 1013-1025.
- Mattson, David J.; Knight, Richard R.; Blanchard, Bonnie M. 1987. The effects of developments and primary roads on grizzly bear habitat use in Yellowstone National Park, Wyoming. *International Conference on Bear Research and Management*. 7: 259-273.
- McCallum, D.A. 1994. Conservation status of flammulated owls in the United States. In: Hayward, G.D.; Verner, J., eds. *Flammulated, boreal, and great gray owls in the United States: a technical conservation assessment*. Gen. Tech. Rep. RM-253. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 74-79. Chapter 6.
- McCallum, D.A.; Gehlbach, F.R. 1988. Nest-site preferences of flammulated owls in western New Mexico. *Condor*. 90(3): 653-661.
- McCool, Stephen F.; Burchfield, J.A.; Allen, Stewart. 1997. Social assessment of the basin. In: Quigley, Thomas M.; Arbelbide, Sylvia J., tech. eds. *An assessment of ecosystem components in the interior Columbia basin including portions of the Klamath and Great Basins*. Gen. Tech. Rep. PNW-GTR-405. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 1872-2009. Chapter 7. Vol. 4. (Quigley, Thomas M., tech. ed. *Interior Columbia Basin Ecosystem Management Project: scientific assessment*).
- McEllin, S.M. 1979. Nest sites and population demographics of white-breasted and pygmy nuthatches in Colorado. *Condor*. 81(4): 348-352.
- McKelvey, Kevin S.; Aubry, Keith B.; Ortega, Yvette K. 1999. History and distribution of lynx in the contiguous United States. In: Ruggiero, Leonard F.; Aubry, Keith B.; Buskirk, Steven W. [and others]. *Ecology and conservation of lynx in the United States*. Gen. Tech. Rep. RMRS-GTR-30WWW. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station: 207-264. Chapter 8.
- McLean, D.D. 1930. The quail of California. *California Div. Fish and Game. Game Bull.* 2: 1-47.
- McLellan, B.N.; Shackleton, D.M. 1988. Grizzly bears and resource-extraction industries: effects of roads on behaviour, habitat use, and demography. *Journal of Applied Ecology*. 25: 451-460.
- McLellan, Bruce N.; Hovey, Fred W. 1995. The diet of grizzly bears in the Flathead River drainage of southeastern British Columbia. *Canadian Journal of Zoology*. 73: 704-712.
- McWhirter, Doug; Smith, Scott; Merrill, Evelyn; Irwin, Larry. 1992. Foraging behavior and vegetation responses to prescribed burning on bighorn sheep winter range. In: *Proceedings of the 8th biennial symposium. Northern Wild Sheep and Goat Council; 1992 Apr 27-May 1; Cody, WY. Thermopolis, WY: Northern Wild Sheep and Goat Council. 8: 264-278. Proceedings available from: Foundation for North American Wild Sheep, 720 Allen Avenue, Cody, WY 82414.*
- Meagher, Mary; Fowler, Sandi. 1989. The consequences of protecting problem grizzly bears. In: Bromley, M., ed. *Bear-people conflicts: Proceedings of a symposium on management strategies; [dates of meeting unknown]; [meeting location unknown]*. Yellowknife, NWT: Northwest Territories Department of Renewable Resources: 141-144.
- Mech, L.D. 1970. Implications of wolf ecology to management. In: Jorgensen, S.E.; Faulkner, L.E.; Mech, L.D., eds. *Proceedings of symposium on wolf management in selected areas of North America*;

[meeting dates unknown]; [meeting location unknown]. [Place of publication unknown]: U.S. Department of the Interior, Fish and Wildlife Service: 39-44.

- Mech, L.D. 1980. Age, sex, reproduction, and spatial organization of lynxes colonizing northeastern Minnesota. *Journal of Mammology*. 61: 261-267.
- Mech, L.D.; Fritts, S.H.; Radde, G.L.; Paul, W.J. 1988. Wolf distribution and road density in Minnesota. *Wildlife Society Bulletin*. 16: 85-87.
- Meints, D.R. 1991. Seasonal movements, habitat use, and productivity of Columbian sharp-tailed grouse in southeastern Idaho. Moscow, ID: University of Idaho. 74 p. M.S. Thesis.
- Meints, D.R.; Connelly, J.W.; Reese, K.P. [and others]. 1992. Habitat suitability index procedure for Columbian sharp-tailed grouse. *Idaho Forest Wildlife and Range Experiment Station Bulletin*. Moscow, ID: University of Idaho, College of Forestry and Range Sciences. (vi): 1-27.
- Merigliano, Michael. F. 1996. Ecology and management of the south fork Snake River cottonwood forest. *Tech. Bull.* 96-9. Boise, ID: U.S. Department of the Interior, Bureau of Land Management, Idaho State Office. 79 p.
- Merrill, Troy; Mattson, David J.; Wright, R. Gerald; Quigley, Howard B. 1999. Defining landscapes suitable for restoration of grizzly bears *Ursus arctos* in Idaho. *Biological Conservation*. 87(2): 231-248.
- Mikkola, H. 1983. *Owls of Europe*. Vermillion, SD: Buteo Books. 397 p.
- Milne, K.A.; Hejl, S.J. 1989. Nest-site characteristics of white-headed woodpeckers. *Journal of Wildlife Management*. 53(1): 50-55.
- Mladenoff, David J.; Sickley, Theodore A.; Haight, Robert G.; Wydeven, Adrian P. 1995. A regional landscape analysis and prediction of favorable gray wolf habitat in the northern Great Lakes region. *Conservation Biology*. 9(2): 279-294.
- Monsen, Stephen B.; McArthur, E. Durant. 1995. Implications of early Intermountain range and watershed restoration practices. In: Roundy, Bruce A.; McArthur, E. Durant; Haley, Jennifer S.; Mann, David K., eds. *Proceedings; wildland shrub and arid land restoration symposium; 1993 October 19-21; Las Vegas, NV. Gen. Tech. Rep. INT-315*. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 16-25.
- Moore, K.R.; Henny, C.J. 1983. Nest site characteristics of three coexisting accipiter hawks in northeastern Oregon. *Raptor Research*. 17: 65-76.
- Moroz, Paul. 1995. Idaho ground squirrel, *Spermophilus brunneus brunneus*, habitat conservation assessment (HCA), conservation strategy (CS). Idaho Interagency Conservation/Prelisting Effort. Idaho Department of Fish and Game; Idaho Department of Parks and Recreation; U.S. Department of the Interior, Fish and Wildlife Service and Bureau of Land Management; U.S. Department of Agriculture, Forest Service. 26 p. Draft unpublished report. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702..
- Morrison, Michael L. 1982. The structure of western warbler assemblages: ecomorphological analysis of the black-throated gray and hermit warblers. *Auk*. 99(3): 503-513.

- Morrow, Larry A.; Stahlman, Phillip W. 1984. The history and distribution of downy brome (Bromus tectorum) in North America. *Weed Science*. 32 (Supplement 1): 2-6.
- Mosconi, S.L.; Hutto, R.L. 1981. The effect of grazing on land birds of a western Montana riparian habitat. In: Peek, J.M.; Dalke, P.D., eds. Symposium on wildlife-livestock interactions; [date of meeting unknown]; [location of meeting unknown]. Moscow, ID: University of Idaho: 221-233.
- Mullican, T.R.; Keller, B.L. 1986. Ecology of the sagebrush vole (Lemmyscus curtatus) in southeastern Idaho. *Canadian Journal of Zoology*. 64(6): 1218-1223.
- Murray, D.L.; Boutin, S. 1991. The influence of snow on lynx and coyote movements: does morphology affect behavior? *Oecologia*. 88: 463-469.
- Murray, T.B. 1938. Upland game birds in Idaho and their future. *Univ. Idaho Bull*. 33: 55-60.
- Mussehl, T.W. 1963. Blue grouse brood cover selection and land use implications. *Journal of Wildlife Management*. 27: 547-556.
- Nagorsen, D.W.; Brigham, R.M. 1993. *Bats of British Columbia*. Vancouver, BC: UBC Press. 164 p.
- Nellis, Carl H.; Wetmore, Stephen P.; Keith, Lloyd B. 1972. Lynx-prey interactions in central Alberta. *Journal of Wildlife Management*. 36(2): 320-329.
- Nelson, Edward W. 1925. Status of the pronghorned antelope, 1922-1924. *Bull.* 1346. Washington, DC: U.S. Department of Agriculture. 64 p.
- Nero, R.W. 1980. *The great gray owl-phantom of the northern forest*. Washington, DC: Smithsonian Institution Press. 167 p.
- Newby, R.E.; McDonald, J.J. 1964. Range extension of the wolverine in Montana. *Journal of Mammalogy*. 45: 485-486.
- Newby, R.E.; Wright, P.L. 1955. Distribution and status of the wolverine in Montana. *Journal of Mammalogy*. 36: 248-253.
- Nice, M.M. 1957. Nesting success in altricial birds. *Auk*. 74(3): 305-321.
- Niemuth, N. 1992. Use of man-made structures by nesting ferruginous hawks in Wyoming. *Prairie Naturalist*. 24(1): 43.
- Noss, R.F.; LaRoe, E.T., III; Scott, J. M. 1995. *Endangered ecosystems of the United States: a preliminary assessment of loss and degradation*. Biological report 28. Washington, DC: National Biological Service.
- Noss, Reed F.; Quigley, Howard B.; Hornocker, Maurice G. [and others]. 1996. Conservation biology and carnivore conservation in the Rocky Mountains. *Conservation Biology*. 10(4): 949-963.
- Nussbaum, Ronald A.; Brodie, Edmund D., Jr.; Storm, Robert M. 1983. *Amphibians and reptiles of the Pacific Northwest*. Moscow, ID: University of Idaho Press. 332 p.
- Oakleaf, R.J. 1971. The relationship of sage grouse to upland meadows in Nevada. Federal Aid Project W-28-2. [Place of publication unknown]: Nevada Department of Fish and Game. 64 p.

- Oakley, C.; Riddle, P. 1974. The impact of a severe winter and fences on antelope mortality in south central Wyoming. In: Proceedings of the 6th antelope states workshop; 1974 February 19-21; Salt Lake City, UT. [Place of publication unknown]: [Publisher unknown]: 155-171.
- Oedekoven, O.O. 1985. Columbian sharp-tailed grouse population distribution and habitat use in south central Wyoming. Laramie, WY: University of Wyoming. 58 p. M.S. thesis.
- O'Farrell, M.J.; Studier, E.H. 1980. Myotis thysanodes. Mammalian Species. No. 137. Baltimore, MD: American Society of Mammalogists. 5 p.
- O'Gara, B.W. 1996. Estimates of pronghorn numbers. Unpublished report. On file with: School of Forestry, University of Montana, Missoula, MT 59812. 1 p.
- Ogg, Alex G. 1994. A review of the chemical control of downy brome. In: Monsen, Stephen B.; Kitchen, Stanley G., eds. Proceedings, ecology and management of annual rangelands; 1992 May 18-21; Boise, ID. Gen. Tech. Rep. INT-GTR-313. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 194-196.
- Ognev, S.I. 1935. Mammals of USSR and adjacent countries. Vol. 3: Carnivora (Fissipedia and Finnepedia). Jerusalem, Israel. 641 p. [Translated by Israel Program for Scientific Translations. 1962].
- Olendorff, R.R.; Stoddart, J.W., Jr. 1974. The potential for management of raptor populations in western grassland. In: Hamerstrom, F.N.; Harrell, B.N.; Olendorff, R.R., eds. Management of raptors: Proceedings of the conference on raptor conservation, part 4; 1973 March 22-24; Fort Collins, CO. Vermillion, SD: Raptor Research Foundation. Raptor Research Reports. 2: 44-48.
- Onderka, D.K.; Wishart, W.D. 1984. A major die-off from pneumonia in southern Alberta. In: Proceedings of the 4th biennial symposium. Northern Wild Sheep and Goat Council; 1984 Apr. 30-May 3; Whitehorse, Yukon. Thermopolis, WY: Northern Wild Sheep and Goat Council. 4: 356-363.
- Oregon Department of Fish and Wildlife. 1987. The herpetology of the Willamette Valley, Oregon. Nongame Wildlife Program Tech. Rep. 86-1-02. Portland, OR: Oregon Department of Fish and Wildlife. 79 p.
- Oregon Department of Fish and Wildlife. 1994. Juniper woodland management. Portland: Oregon Department of Fish and Wildlife, Habitat Conservation Division. 26 p.
- Orloff, S.; Hall, F.; Spiegel, L. 1986. Distribution and habitat requirements of the San Joaquin kit fox in the northern extreme of their range. Transactions of the Western Section of the Wildlife Society: 2260-2270.
- Ormsbee, Patricia C.; McComb, William C. 1998. Selection of day roosts by female long-legged myotis in the central Oregon Cascade Range. Journal of Wildlife Management. 62(2): 596-602.
- Owens, R.A.; Myres, M.T. 1973. Effects of agriculture upon populations of native passerine birds of an Alberta fescue grassland. Canadian Journal of Zoology. 51(7): 697-713.
- Page, J.L.; Dodd, N.; Osbornde, T.O.; Carson, J.A. 1978. The influence of livestock grazing on non-game wildlife. CALNEVA Wildlife. [Place of publication unknown]: The Wildlife Society, Western Section: 159-173.
- Parker, G.R.; Maxwell, J.W.; Morton, L.D. [and others]. 1983. The ecology of lynx (Lynx canadensis) on

- Cape Brenton Island. Canadian Journal of Zoology. 61: 770-786.
- Patla, S. 1990. Northern goshawk monitoring report no. 2, 1989. [Not paged]. On file with: U.S. Department of Agriculture, Targhee National Forest, P.O. Box 208, St. Anthony, ID 83445.
- Patterson, R.L. 1952. The sage grouse in Wyoming. Denver, CO: Sage Books. 341 p.
- Pedivillano, Cathy; Wright, Gerald R. 1987. The influence of visitors on mountain goat activities in Glacier National Park, Montana. Biological Conservation. 39(1): 1-11.
- Pekins, P.J.; Lindzey, F.G.; Gessaman, J.A. 1991. Physical characteristics of blue grouse winter use-trees and roost sites. Great Basin Naturalist. 51(3): 244-248.
- Pellant, Mike. 1994. History and applications of the Intermountain greenstripping program. In: Monsen, Stephen B.; Kitchen, Stanley G., eds. Proceedings-ecology and management of annual rangelands; 1992 May 18-21. Gen. Tech. Rep. INT-313. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 63-68.
- Perkins, J.M. 1992-1994. Results of summer bat surveys, Wallowa Valley Ranger District, Eagle Cap Ranger District, and the HCNRA of the Wallowa-Whitman National Forest, Wallowa County, OR. U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, Wallowa-Whitman National Forest. [Not paged]. Unpublished report. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.
- Perkins, J.M.; Cross, S.P. 1988. Differential use of some coniferous forest habitats by hoary and silver-haired bats in Oregon. Murrelet. 69(1): 21-24.
- Perkins, J.M.; Schommer, T. 1992. Survey protocol and an interim species conservation strategy for Plecotus townsendii in the Blue Mountains of Oregon and Washington. 23 p. Unpublished report. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.
- Perkins, J. Mark. 1987. Distribution, status, and habitat affinities of Townsend's big-eared bat (Plecotus townsendii) in Oregon. Tech. Rep. 85-5-01. [Portland, OR]: Oregon Department of Fish and Wildlife, Nongame Wildlife Program.
- Perkins, J. Mark. 1996. Does competition for roosts influence bat distribution in a managed forest? In: Barclay, Robert M.R.; Brigham, R. Mark, eds. Bats and forest symposium; 1995 October 19-21; [meeting location unavailable]. Victoria, BC: British Columbia Ministry of Forests Research Program: 164-172.
- Perkins, J. Mark; Peterson, Joshua R. 1997. Bat distribution in the juniper woodlands of the Idaho Owyhee Mountains, summer 1996. Tech. Bull. No. 97-4. U.S. Department of the Interior, Bureau of Land Management, Idaho State Office. 12 p.
- Perlmeier, S.I. 1995. Bats and bridges: a field study of the thermal conditions and social organization of night roosts in the Willamette National Forest. North York, ON: York University. 98 p. M.S. thesis.
- Perritt, J.E.; Best, L.B. 1989. Effects of weather on the breeding ecology of vesper sparrows in Iowa crop fields. American Midland Naturalist. 121(2): 355-360.

- Peterson, J.G. 1970. The food habits and summer distribution of juvenile sage grouse in central Montana. *Journal of Wildlife Management*. 34(1): 147-155.
- Pierson, Elizabeth D.; Rainey, William E.; Koontz, Dolora M. 1991. Bats and mines: experimental mitigation for Townsend's big-eared bat at the McLaughlin mine in California. In: Corner, R.D., Davis, P.R., Foster, S.Q. [and others], eds. *Proceedings V: issues and technology in the management of impacted wildlife*; 1991 April 8-10; Snowmass Resort, CO. Boulder, CO: Thorne Ecological Institute: 31-42.
- Pletscher, Daniel H.; Ream, Robert R.; Boyd, Diane K. [and others]. 1997. Population dynamics of a recolonizing wolf population. *Journal of Wildlife Management*. 61(2): 459-465.
- Ports, M.A.; Bradley, P.V. 1996. Habitat affinities of bats from northeastern Nevada. *Great Basin Naturalist*. 56(1): 48-53.
- Ports, M.A.; George, S.B. 1990. *Sorex preblei* in the northern Great Basin. *Great Basin Naturalist*. 50(1): 93-95.
- Post, G. 1951. Effects of toxaphene and chlordane on certain game birds. *Journal of Wildlife Management*. 15(4): 381-386.
- Pough, F.H. 1973. Lizard energetics and diet. *Ecology*. 54(4): 837-844.
- Poulton, C.E. 1955. Ecology of the non-forested vegetation in Umatilla and Morrow Counties, Oregon. Pullman, WA: Washington State University. 166 p. Ph.D. dissertation.
- Powell, Roger A.; Zielinski, William J. 1994. Fisher. In: Ruggiero, L.; Aubry, K.; Buskirk, S. [and others], tech. eds. *The scientific basis for conserving forest carnivores: American marten, fisher, lynx and wolverine in the western United States*. Gen. Tech. Rep. RM-254. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 38-73. Chapter 3.
- Powers, L.R.; Howard, R.R.; Trost, C.H. 1973. Population status of the ferruginous hawk in southeastern Idaho and northern Utah. In: Murphy, J.R.; White, C.M.; Harrell, B.E., eds. *Population status of raptors: Proceedings of the conference on raptor conservation techniques, part 6*; 1973 March 22-24; Fort Collins, CO. *Raptor Research Reports*. 3: 153-157.
- Pulliamin, E. 1968. Breeding biology of the wolverine (*Gulo gulo* L.) in Finland. *Annales Zoologica Fennici*. 5: 338-344.
- Pulliam, H.R. 1988. Sources, sinks, and population regulation. *American Naturalist*. 132(5): 652-661.
- Pulliam, H.R.; Danielson, B.J. 1991. Sources, sinks, and habitat selection: a landscape perspective on population dynamics. *American Naturalist*. 137:55-66.
- Pylypec, B. 1991. Impacts of fire on bird populations in a fescue prairie. *Canadian Field Naturalist*. 105(3): 346-349.
- Quick, H.F. 1956. Effects of exploitation on a marten population. *Journal of Wildlife Management*. 20(3): 267-274.
- Quigley, Thomas M.; Haynes, Richard W.; Graham, Russell T., tech eds. 1996. *Integrated scientific*

- assessment for ecosystem management in the interior Columbia basin and portions of the Klamath and Great Basins. Gen. Tech. Rep. PNW-GTR-382. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 303 p. (Quigley, Thomas M., ed. Interior Columbia Basin Ecosystem Management Project: scientific assessment).
- Quigley, Thomas M.; Lee, Danny C.; Haynes, Richard W. [and others]. 1997. Ecological integrity, socioeconomic resiliency, and trends in risk. In: Quigley, Thomas M.; Lee, Kristine M.; Arbelbide, Sylvia J., tech. eds. Evaluation of EIS alternatives by the Science Integration Team. Gen. Tech. Rep. PNW-GTR-406. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station: 835-896. Vol. 2.
- Rabe, Michael J.; Morrell, Thomas E.; Green, Heather [and others]. 1998. Characteristics of ponderosa pine snag roosts used by reproductive bats in northern Arizona. *Journal of Wildlife Management*. 62(2): 612-621.
- Ramakka, J.M.; Woyewodzic, R.T. 1993. Nesting ecology of ferruginous hawk in northwestern New Mexico. *Journal of Raptor Research*. 27(2): 97-101.
- Raphael, M.G.; Jones, L.C. 1997. Characteristics of resting and denning sites of American martens in central Oregon and western Washington. In: Proulx, G.; Bryant, H.N.; Woodard, P.M., eds. *Martes: taxonomy, ecology, techniques, and management*. Edmonton, AB: The Provincial Museum of Alberta: 146-165.
- Raphael, M.G.; White, M. 1984. Use of snags by cavity-nesting birds in the Sierra Nevada. *Wildlife Monographs*. 86: 1-66.
- Remington, T.E.; Hoffman, R.W. 1996. Food habits and preferences of blue grouse during winter. *Journal of Wildlife Management*. 60(4): 808-817.
- Reynolds, R.T. 1983. Management of western coniferous forest habitat for nesting accipiter hawks. Gen. Tech. Rep. RM-102. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 6 p.
- Reynolds, R.T.; Graham, R.T.; Reiser, M.H. [and others]. 1992. Management recommendations for the northern goshawk in the southwestern United States. Gen. Tech. Rep. RM-217. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 184 p.
- Reynolds, R.T.; Linkhart, B.D. 1990. Extra-pair copulation and extra-range movements in flammulated owls. *Ornis Scandinavica*. 21: 74-77.
- Reynolds, R.T.; Linkhart, B.D. 1992. Flammulated owls in ponderosa pine: evidence of preference for old growth. In: Old-growth forests in the Southwest and Rocky Mountain regions: Proceedings of a workshop. Gen. Tech. Rep. RM-213. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station: 166-169.
- Rich, T. 1981. Microgeographic variation in the song of the sage sparrow. *Condor*. 83(2): 113-119.
- Rich, T. 1984. Monitoring burrowing owl populations: implications of burrow re-use. *Wildlife Society Bulletin*. 12(2): 178-180.
- Rich, T. 1986. Habitat and nest-site selection by burrowing owls in the sagebrush steppe of Idaho.

- Journal of Wildlife Management. 50(4): 548-555.
- Rich, T.D. 1978. Cowbird parasitism of sage and Brewer's sparrows. *Condor*. 80: 348.
- Rich, T.D. 1985. Sage grouse population fluctuations: evidence for a 10-year cycle. *Tech. Bull.* 85-1. Boise, ID: U.S. Department of the Interior, Bureau of Land Management, Idaho State Office. 29 p.
- Rich, T.D.; Rothstein, S.I. 1985. Sage thrashers reject cowbird eggs. *Condor*. 87: 561-562.
- Rickard, William H.; Poole, Linda D. 1989. Terrestrial wildlife of the Hanford Site: past and future. *Northwest Science*. 63(4): 183-192.
- Rideout, Chester B. Mountain goat. 1978. In: Schmidt, John L.; Gilbert, Douglas L., eds. *Big game of North America: ecology and management*. Harrisburg, PA: Stackpole Books: 149-159.
- Risenhoover, Kenneth L.; Bailey, James A. 1985. Foraging ecology of mountain sheep: implications for habitat management. *Journal of Wildlife Management*. 49(3): 797-804.
- Roberson, Jay A. 1986. Sage grouse-sagebrush relationships: a review. In: McArthur, E. Durant; Welch, Bruce L., eds. *Proceedings, symposium on the biology of artemisia and chrysothamnus; 1984 July 9-13; Provo, UT. Gen. Tech. Rep. INT-200*. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 157-167.
- Robertson, M.D. 1989. Statewide survey of mountain quail, 1989: a report on the status of mountain quail in Idaho. Boise, ID: Idaho Department of Fish and Game. 27 p. Unpublished report. On file with Boise, ID: Idaho Department of Fish and Game, P.O. Box 25, Boise, ID 83707.
- Robertson, M.D. 1991. Winter ecology of migratory sage grouse and associated effects of prescribed fire in southeastern Idaho. Moscow, ID: University of Idaho. 88 p. M.S. thesis.
- Robinson, Scott K.; Rothstein, Stephen I.; Brittingham, Margaret C. [and others]. 1995. Ecology and behavior of cowbirds and their impact on host populations. In: Martin, Thomas E.; Finch, Deborah M., eds. *Ecology and management of neotropical migratory birds*. New York: Oxford University Press: 428-460.
- Rodenhouse, H.L.; Best, L.B.; O'Connor, R.J.; Bollinger, E. K. 1995. Effects of agricultural practices and farmland structures. In: Martin, T.E.; Finch, D., eds. *Ecology and management of neotropical migratory birds*. New York: Oxford University Press: 269-293.
- Rodgers, R.D. 1983. Reducing wildlife losses to tillage in fallow wheat fields. *Wildlife Society Bulletin*. 11: 31-28.
- Rogers, Lynn L.; Rogers, S.M. 1976. Parasites of bears: a review. In: Pelton, M.R; Lentfer, J.W.; Folk, G.E., eds. *Bears: their biology and management*. IUCN Publication New Ser. 40: 411-430.
- Roloff, G. 1995. Canadian lynx (*Felis lynx*) in Idaho: habitat conservation assessment and conservation strategy. Unpublished report. [Boise, ID: Idaho Department of Fish and Game]. 13 p. On file with: Idaho Department of Fish and Game, P.O. Box 25, Boise, ID 83707.
- Rolseth, S.L.; Koehler, C.E.; Barclay, R.M.R. 1994. Differences in the diets of juvenile and adult hoary bats, *Lasiurus cinereus*. *Journal of Mammalogy*. 75(2): 394-398.

- Rood, Stewart B.; Heinze-Milne, Sig. 1989. Abrupt downstream forest decline following river damming in southern Alberta. *Canadian Journal of Botany*. 67: 1744-1749.
- Rosenberg, K.V.; Raphael, M.G. 1986. Effects of forest fragmentation on vertebrates in Douglas-fir forests. In: Verner, J.; Morrison, M.L.; Ralph, C.J., eds. *Wildlife 2000: modeling habitat relationships of terrestrial vertebrates*. Madison, WI: University of Wisconsin Press: 263-272.
- Rotenberry, John T.; Wiens, John A. 1980. Habitat structure, patchiness, and avian communities in North American steppe vegetation: a multivariate analysis. *Ecology*. 61: 1228-1250.
- Rothstein, S.I. 1994. The cowbird's invasion of the Far West: history, causes and consequences experienced by host species. *Studies in Avian Biology*. 15: 301-315.
- Rothstein, S.I.; Verner, J.; Stevens, E.; Ritter, L.V. 1987. Behavioral differences among sex and age classes of the brown-headed cowbird and their relation to the efficacy of a control program. *Wilson Bulletin*. 99(3): 322-337.
- Rudolph, Karen M.; Cowan, Lisa M.; Hunter, David L. [and others]. [In prep.]. Sharing of *Pasteurella* spp. among free-ranging bighorn sheep, domestic sheep, and feral goats.
- Ruediger, Bill. 1996. The relationship between rare carnivores and highways. In: Evink, G; Garrett, P.; Berry, J., eds. *Transportation and wildlife: reducing wildlife mortality and improving wildlife passageways across transportation corridors*. Proceedings of a symposium; 1996 April 3-May 2; Orlando, FL. Orlando, FL: Department of Transportation, Federal Highway Administration: 24-38.
- Ruggiero, Leonard F.; Aubry, Keith B.; Buskirk, Steven W. [and others]. 1999. Ecology and conservation of lynx in the United States. Gen. Tech. Rep. RMRS-GTR-30WWW. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 480 p.
- Russell, R.W.; Carpenter, F.L.; Hixon, M.A.; Paton, David C. 1994. The impact of variation in stopover habitat quality on migrant rufous hummingbirds. *Conservation Biology*. 8(2): 483-490.
- Ryan, L.A.; Carey, A.B. 1995. Biology and management of the western gray squirrel and Oregon white oak woodlands: with emphasis on the Puget Trough. Gen. Tech. Rep. PNW-GTR-348. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 36 p.
- Saab, V.A. 1996. Influences of spatial scale and land-use practices on habitat relationships of breeding birds in cottonwood riparian forests. Boulder, CO: University of Colorado. 140 p. Ph.D. dissertation.
- Saab, V.A. 1998. Effects of recreational activity and livestock grazing on habitat use by breeding birds in cottonwood forests along the South Fork Snake River. Tech. Bull. No. 98-17. BLM-ID-PT99-004-1150. Boise, ID: U.S. Department of the Interior, Bureau of Land Management. 48 p.
- Saab, V.A. 1999. Importance of spatial scale to habitat use by breeding birds in riparian forests: a hierarchical analysis. *Ecological Applications*. 9(1): 135-151.
- Saab, V.A.; Bock, C.E.; Rich, T.D.; Dobkin, D.S. 1995. Livestock grazing effects in western North America. In: Martin, Thomas E.; Finch, Deborah M., eds. *Ecology and management of neotropical migratory birds*. New York: Oxford University Press: 311-353.
- Saab, V.; Dudley, J. 1995. Cavity use and nest usurpation by Lewis' woodpecker. On file with: U.S. Department of Agriculture, Forest Service, Intermountain Research Station, 316 E. Myrtle, Boise, ID

83702.

- Saab, V.A.; Marks, J.S. 1992. Summer habitat use by Columbian sharp-tailed grouse in western Idaho. *Great Basin Naturalist*. 52(2): 166-173.
- Saab, V.A.; Rich, T.D. 1997. Large-scale conservation assessment for neotropical migratory land birds in the interior Columbia River Basin. Gen. Tech. Rep. PNW-399. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 57 p. (Quigley, Thomas M., ed. Interior Columbia Basin Ecosystem Management Project: scientific assessment).
- Saab, Victoria A.; Dudley, Jonathan G. 1998. Responses of cavity-nesting birds to stand-replacement fire and salvage logging in ponderosa pine/Douglas-fir forests of southwestern Idaho. Res. Pap. RMRS-RP-11. Ogden, UT: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 17 p.
- Samson, F.B. 1980. Island biogeography and the conservation of prairie birds. In: Proceedings, 7th North American prairie conference; 1980 August 4-6; Springfield, MO. Springfield, MO: Southwest Missouri State University: 293-305.
- Sanborn, S.R.; Loomis, R.B. 1979. Systematics and behavior of collared lizards (*Crotaphytus*, *Iguanidae*) in southern California. *Herpetologica*. 35(2): 101-106.
- Sanders, S.D.; Flett, M.A. 1989. Montane riparian habitat and willow flycatchers: threats to a sensitive environment and species. Gen. Tech. Rep. PSW-110. Berkeley, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Forest and Range Experiment Station: 262-266.
- Sarell, M.J.; McGuinness, K.P. 1993. Rare bats of the shrub-steppe ecosystem of eastern Washington. Gen. Tech. Rep. Olympia, WA: Washington Department of Wildlife.
- Sauer, J.R.; Peterjohn, B.J.; Schwartz, S.; Hines, J.E. 1996. The North American breeding bird survey home page; version 95.1. Laurel, MD: Patuxent Wildlife Research Center. <http://www.mbr-pwrc.usgs.gov/bbs/bbsold.html>.
- Saunders, Jack K., Jr. 1955. Food habits and range use of the Rocky Mountain goat in the Crazy Mountains, Montana. *Journal of Wildlife Management*. 19(4): 429-437.
- Savage, D.E. 1969. The relationships of sage grouse to upland meadows in Nevada. *Transactions, The Wildlife Society, California-Nevada Section*. [Place of publication unknown]: The Wildlife Society. 16: 8-17.
- Schmutz, J.K. 1984. Ferruginous and Swainson's hawk abundance and distribution in relation to land use in southeastern Alberta. *Journal of Wildlife Management*. 48(4): 1180-1187.
- Schmutz, J.K.; Hungle, D.J. 1989. Populations of ferruginous and Swainson's hawks increase in synchrony with ground squirrels. *Canadian Journal of Zoology*. 67(10): 2596-2601.
- Schroeder, M.A. 1994. Productivity and habitat use of Columbian sharp-tailed grouse in north-central Washington. Yakima, WA: Washington Department of Fish and Wildlife; progress report.
- Schuler, Carol A.; Rickard; William H.; Sargeant, Glen A. 1993. Conservation of habitats for shrubsteppe birds. *Environmental Conservation*. 20 (1): 57-64.

- Schulz, T.T.; Leininger, W.C. 1991. Nongame wildlife communities in grazed and ungrazed montane riparian sites. *Great Basin Naturalist*. 51: 286-292.
- Scott, M.D.; Servheen, G. 1985. Caribou ecology. Boise, ID: Idaho Department of Fish and Game. Job completion rep., Proj. No. W-160-R-11. 136 p.
- Servheen, Christopher. 1983. Grizzly bear food habits, movements, and habitat selection in the Mission Mountains, Montana. *Journal of Wildlife Management*. 47(4): 1026-1035.
- Shaffer, Mark L. 1981. Minimum population sizes for species conservation. *BioScience*. 31(2): 131-134.
- Sharp, B.E. 1992. Neotropical migrants on National Forests in the Pacific Northwest: a compilation of existing information. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region. Unpublished report. On file with: U.S. Department of Agriculture, Forest Service, Pacific Northwest Region, P.O. Box 3623, Portland, OR 97208.
- Shriver, W.G.; Vickery, P.D.; Hedges, S.A. 1996. Effects of summer burns on Florida grasshopper sparrows. *Florida Field Naturalist*. 24(3): 68-73.
- Shump, K.A.; Shump, A.U. 1982. Lasiurus cinereus. *Mammalian Species*. No. 185. Baltimore, MD: American Society of Mammalogists. 5 p.
- Sibley, David. 1993. *Birds of Cape May*. 1st ed. Cape May Point, NJ: New Jersey Audubon Society, Cape May Point Bird Observatory. 145 p.
- Siddle, C.; Davidson, G. 1991. Status of the Lewis' woodpecker (Melanerpes lewis) in British Columbia. Report commissioned by Wildlife Branch, Ministry of Environment, Victoria, British Columbia, Canada. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, USDA Forest Service, 304 N. 8th Street, Boise, ID 83702.
- Simpson, K. 1987. The effects of snowmobiling on winter range use by mountain caribou. *Wildlife Working Rep.* WR-25. Nelson, BC: Ministry of Environment and Parks. 13 p.
- Singer, F.J. 1978. Behavior of mountain goats in relation to U.S. Highway 2, Glacier National Park, Montana. *Journal of Wildlife Management*. 42(3): 591-597.
- Singer, F.J. 1979. Status and history of timber wolves in Glacier National Park, Montana. In: Klinghammer, E., ed. *The behavior and ecology of wolves*. New York: Garland STPM Press: 19-42.
- Singer, F.J.; Doherty, J.L. 1985. Movements and habitat use in an un hunted population of mountain goats, Oreamnos americanus. *Canadian Field Naturalist*. 99: 205-217.
- Sirotnak, J.M.; Reese, K.P.; Connelly, J.; Radford, K. 1991. Characteristics of Conservation Reserve Program fields in southeastern Idaho associated with upland bird and big game habitat use. Federal Aid Wildlife Restoration Project W-160-R. Boise, ID: Idaho Department of Fish and Game.
- Smith, Christian A. 1986. Rates and causes of mortality in mountain goats in southeast Alaska. *Journal of Wildlife Management*. 50(4): 743-746.
- Smith, D.G.; Murphy, J.R. 1973. Breeding ecology of raptors in the eastern Great Basin of Utah. *Science Bulletin Biological Series*. Vol. 18, No. 3. Provo, UT: Brigham Young University. 76 p.

- Smith, Kirby G.; Urquhart, Martin A.; Festa-Bianchet, Marco. 1992. Preliminary observations of timing and causes of mountain goat kid mortality in west-central Alberta. In: Proceedings of the 8th biennial symposium. Northern Wild Sheep and Goat Council; 1992 Apr. 27-May 1; Cody, WY. Thermopolis, WY: Northern Wild Sheep and Goat Council. 8: 293-304.
- Smith, Michael Roger. 1994. Evaluating the conservation of avian diversity in eastern Washington: a geographic analysis of upland breeding birds. Seattle: University of Washington. 89 p. M.S. thesis.
- Smith, R.L. 1963. Some ecological notes on the grasshopper sparrow. *Wilson Bulletin*. 75(2): 159-165.
- Sorensen, E. 1986. A precipitous decline in Lewis' woodpecker in Salt Lake and Davis counties. *Utah Birds*. 2: 45-54.
- Spaulding, E.S. 1949. *The quails*. New York: The MacMillan Co. 123 p.
- Speiser, R. 1992. Notes on the natural history of the northern goshawk. *Kingbird*. 42: 133-137.
- Spraker, T.R.; Hibbler, C.P.; Schoonveld, G.G.; Adbetm, W.S. 1984. Pathological changes and micro-organisms found during a stress related die-off. *Journal of Wildlife Diseases*. 20: 319-327.
- Spring, Lowell W. 1965. Climbing and pecking adaptations in some North American woodpeckers. *Condor*. 67(6): 457-488.
- Squires, J.R.; Reynolds, R.T. 1997. Northern goshawk (*Accipiter gentilis*). In: Poole, A.; Gill, F., eds. *The birds of North America*. No. 298. Philadelphia, PA: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 32 p.
- Squires, J.R.; Ruggiero, L.F. 1996. Nest-site preference of northern goshawks in south central Wyoming. *Journal of Wildlife Management*. 60(1): 170-177.
- St. Clair, L.L.; Johansen, J.R. 1993. Introduction to the symposium on soil crust communities. *Great Basin Naturalist*. 53(1): 1-4.
- St. Clair, Larry L.; Johansen, Jeffrey R.; Rushforth, Samuel R. 1993. Lichens of soil crust communities in the Intermountain area of the western United States. *Great Basin Naturalist*. 53: 5-12.
- Stallcup, P.L. 1962. Contents of 103 stomachs of northern three-toed woodpeckers. *Journal of the Colorado-Wyoming Academy of Sciences*. 5: 39.
- Stebbins, Robert C. 1985. *Peterson field guides: western reptiles and amphibians*. Boston, MA: Houghton Mifflin Co. 336 p.
- Steenhof, K.; Kochert, M.N. 1985. Dietary shifts of sympatric buteos during a prey decline. *Oecologia*. 66(1): 6-16.
- Stewart, George; Hull, A.C. 1949. Cheatgrass (*Bromus tectorum* L.)-an ecologic intruder in southern Idaho. *Ecology*. 30: 58-74.
- Swenson, Jon E. 1985. Compensatory reproduction in an introduced mountain goat population in the Absaroka Mountains, Montana. *Journal of Wildlife Management*. 49(4): 837-843.
- Tashiro-Vierling, K.Y. 1994. Population trends and ecology of the Lewis' woodpecker (*Melanerpes lewis*)

- in southeastern Colorado. Boulder: CO: University of Colorado. 73 p. M.A. thesis.
- Terra-Berns, Mary; Call, Paula; Harris, Chuck [and others]. 1997. Canada lynx in Idaho: past, present, and future. Boise, ID: Idaho Department of Fish and Game. 79 p.
- Terres, John K. 1991. The Audubon Society encyclopedia of North American birds. New York: Wings Books. 1109 p.
- Thiel, R.P. 1985. Relationship between road densities and wolf habitat suitability in Wisconsin. *American Midland Naturalist*. 113: 404-407.
- Thomas, Jack Ward; Anderson, Ralph G.; Maser, Chris; Bull, Evelyn L. 1979. Snags. In: Thomas, Jack Ward, tech. ed. *Wildlife habitats in managed forests, the Blue Mountains of Oregon and Washington*. Agric. Handb. No. 553. Washington, DC: U.S. Department of Agriculture: 60-77.
- Thompson, I.D. 1994. Marten populations in uncut and logged boreal forests in Ontario. *Journal of Wildlife Management*. 58(2): 272-280.
- Thompson, I.D.; Colgan, P.W. 1987. Numerical responses of martens to a food shortage in north central Ontario. *Journal of Wildlife Management*. 51(4): 824-835.
- Thomsen, Lise. 1971. Behavior and ecology of burrowing owls on the Oakland Municipal Airport. *Condor*. 73(2): 177-192.
- Thorne, E. Tom; Hickey, William O.; Stewart, Shawn T. 1985. Status of California and Rocky Mountain bighorn sheep in the United States. In: Hoefs, Manfred, ed. *Wild sheep: distribution, abundance, management, and conservation of the sheep of the world and closely related mountain ungulates*. Northern Wild Sheep and Goat Council Special Report: 56-74.
- Thurber, Joanne M.; Peterson, Rolf O.; Drummer, Thomas D.; Thomasma, Scott A. 1994. Gray wolf response to refuge boundaries and roads in Alaska. *Wildlife Society Bulletin*. 22(1): 61-68.
- Thurow, T.L.; White, C.M. 1984. Nesting success and prey selection of long-eared owls along a juniper/sagebrush ecotone in south central Idaho. *Murrelet*. 65(1): 10-14.
- Thurow, T.L.; White, C.M.; Howard, R.P.; Sullivan, J.F. 1980. *Raptor ecology of Raft River Valley, Idaho*. Idaho Falls, ID: U.S. Department of Energy, Idaho Operations Office. 45 p.
- Tirhi, M.J. 1995. *Washington state management plan for sharp-tailed grouse*. Olympia, WA: Washington Department of Fish and Wildlife, Wildlife Management Program. 99 p.
- Tobalski, B.W. 1997. Lewis' woodpecker (*Melanerpes lewis*). In: Poole, A.; Gill, F., eds. *The birds of North America*. Bimonthly. No. 284: Washington, DC: American Ornithologists' Union. [page numbers unavailable]. ISSN: 1061-5466. Issued Jointly: Philadelphia, PA: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union.
- Tomich, P. Quentin. 1982. Ground squirrels (*Spermophilus beecheyi* and allies). In: Chapman, Joseph A.; Feldhamer, George A., eds. *Wild mammals of North America: biology, management, and economics*. Baltimore, MD: Johns Hopkins University Press: 192-208.
- Trulio, L.A. 1995. Passive relocation: a method to preserve burrowing owls on disturbed sites. *Journal of Field Ornithology*. 66(1): 99-106.

- Tuttle, M.D.; Taylor, D.A.R. 1994. Bats and mines. Resource Publ. No. 3. [Place of publication unknown]: Bat Conservation International, Incorporated.
- Ulliman, M.J. 1995. Rangewide conservation assessment for Columbian sharp-tailed grouse and its habitats. Moscow, ID: University of Idaho, Department of Fish and Wildlife Resources. 96 p.
- U.S. Department of Agriculture, Forest Service. 1996. Status of the interior Columbia basin: summary of scientific findings. Gen. Tech. Rep. PNW-GTR-385. Portland, OR: US Department of Agriculture, Forest Service, Pacific Northwest Research Station; U.S. Department of the Interior, Bureau of Land Management. 144 p.
- U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management. 1997a. Eastside draft environmental impact statement. Walla Walla, WA: Interior Columbia Basin Ecosystem Management Project. [Irregular pagination]. 2 vol.
- U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Bureau of Land Management. 1997b. Upper Columbia River basin draft environmental impact statement. Boise, ID: Interior Columbia Basin Ecosystem Management Project. [Irregular pagination]. 2 vol.
- U.S. Department of Agriculture, Forest Service; U.S. Department of the Interior, Fish and Wildlife Service. 1996. Conservation agreement: Spermophilus brunneus brunneus, northern Idaho ground squirrel. McCall, ID: U.S. Department of Agriculture, Forest Service, Payette National Forest; Boise, ID: U.S. Department of the Interior, Fish and Wildlife Service. 14 p.
- U.S. Department of the Interior, Bureau of Land Management. 1995. Mountain sheep ecosystem management strategy in the 11 western states and Alaska. BLM/SC/PL-95/001+6600. 79 p. [Place of publication unknown]: U.S. Department of the Interior, Bureau of Land Management.
- U.S. Department of the Interior, Bureau of Land Management. 1997. Proposed standards for rangeland health and guidelines for livestock grazing management. BLM/ID/PT-97/002+4120. Boise ID: U.S. Department of the Interior, Bureau of Land Management, Idaho State Office.
- U.S. Department of the Interior, Fish and Wildlife Service. 1987. Northern Rocky Mountain wolf recovery plan. Denver, CO: U.S. Department of the Interior, Fish and Wildlife Service. 119 p.
- U.S. Department of the Interior, Fish and Wildlife Service. 1993. Grizzly bear recovery plan. Missoula, MT. 181 p.
- U.S. Department of the Interior, Fish and Wildlife Service. 1994. Selkirk mountain woodland caribou recovery plan. Portland, OR. 59 p.
- U.S. Department of the Interior, Fish and Wildlife Service. 1997. The grizzly bear recovery in the Bitterroot ecosystem. Draft environmental impact statement, Lakewood, CO: Region 6, USDI Fish and Wildlife Service. [number of pages unknown].
- U.S. Government. 1994. Rules and regulations. Federal Register 59:224 (Nov. 22, 1994):60252-60281.
- Van Ballenberghe, Victor; Erickson, Albert W.; Byman, David. 1975. Ecology of the timber wolf in northeastern Minnesota. Wildlife Monographs. No. 43. 43 p.
- Van Dyke, Walter A.; Sands, Alan; Yoakum, Jim [and others]. 1983. Wildlife habitats in managed rangelands--the great basin of southeastern Oregon--bighorn sheep. Gen. Tech. Rep. PNW-GTR-159.

- Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station; U.S. Department of the Interior, Bureau of Land Management. 39 p.
- Van Horne, B.; Bader, A. 1990. Diet of nestling winter wrens in relationship to food availability. *Condor*. 92(2): 413-420.
- Vickery, P.D. 1996. Grasshopper sparrow (*Ammodramus savannarum*). In: Poole, A.; Gill, F., eds. *The birds of North America*. No. 239. Philadelphia, PA: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 24 p.
- Vickery, P.D.; Hunter, M.L., Jr.; Melvin, S.M. 1994. Effect of habitat area on the distribution of grassland birds in Maine. *Conservation Biology*. 8: 1087-1097.
- Vierling, K.T. 1997. Habitat selection of Lewis' woodpeckers in southeastern Colorado. *Wilson Bulletin*. 109: 121-130.
- Villard, P. 1994. Foraging behavior of black-backed and three-toed woodpeckers during spring and summer in a Canadian boreal forest. *Canadian Journal of Zoology*. 72(11): 1957-1959.
- Vogel, C.A.; Reese, K.P. 1995. Mountain quail status report: a document preliminary to a habitat conservation assessment for mountain quail. On file with: U.S. Department of Agriculture, Forest Service; U.S. Department of Interior, Bureau of Land Management; Interior Columbia Basin Ecosystem Management Project, 112 E. Poplar, Walla Walla, WA 99362. 33 p.
- Wai-Ping, V.; Fenton, M.B. 1989. Ecology of the spotted bat (*Euderma maculatum*) roosting and foraging behavior. *Journal of Mammalogy*. 70(3): 617-622.
- Wakelyn, Leslie A. 1987. Changing habitat conditions on bighorn sheep ranges in Colorado. *Journal of Wildlife Management*. 51(4): 904-912.
- Wakkinen, W.L. 1990. Nest site characteristics and spring-summer movements of migratory sage grouse in southeastern Idaho. Moscow, ID: University of Idaho. 57 p. M.S. thesis.
- Wallestad, Richard. 1975. Male sage grouse responses to sagebrush treatment. *Journal of Wildlife Management*. 39(3): 482-484.
- Wallestad, Richard; Peterson, Joel G.; Eng, Robert L. 1975. Foods of adult sage grouse in central Montana. *Journal of Wildlife Management*. 39(3): 628-630.
- Ward, J.C.; Martin, M.; Allred, W. 1942. The susceptibility of sage grouse to strychnine. *Journal of Wildlife Management*. 6(1): 55-57.
- Ward, J.M.; Kennedy, P. L. 1996. Effects of supplemental food on size and survival of juvenile northern goshawks. *Auk*. 113: 200-208.
- Ward, Richard M.P.; Krebs, Charles J. 1985. Behavioural responses of lynx to declining snowshoe hare abundance. *Canadian Journal of Zoology*. 63(12): 2817-2824.
- Washington Department of Wildlife. 1993a. Distribution status of the mountain quail (*Oreortyx pictus*) in Washington. Olympia, WA: Washington Department of Wildlife. Unpublished report. On file with: Olympia, WA: Washington Department of Wildlife, 600 Capitol Way North, Olympia, WA 98501-1091. 22 p.

- Washington Department of Wildlife. 1993b. Status of the western gray squirrel (*Sciurus griseus*) in Washington. Olympia, WA: Washington Department of Wildlife. Unpublished report. On file with: Olympia, WA: Washington Department of Wildlife, 600 Capitol Way North, Olympia, WA 98501-1091. 33 p.
- Waters, J.R.; McKelvey, K.S.; Zabel, C.J.; Oliver, W.W. 1994. The effects of thinning and broadcast burning on sporocarp production of hypogeous fungi. *Canadian Journal of Forest Research*. 24: 1516-1522.
- Waters, J.R.; Zabel, C.J. 1995. Northern flying squirrel densities in fir forests of northeastern California. *Journal of Wildlife Management*. 59(4): 858-866.
- Weaver, J. 1993. Lynx, wolverine and fisher in the western United States. Northern Rockies Conservation Cooperative. Unpublished report. On file with: Northern Rockies Conservation Cooperative, P.O. Box 8594, Missoula, MT 59807. 132 p.
- Weaver, J. 1994. Ecology of wolf predation amidst high ungulate diversity in Jasper National Park, Alberta. Missoula, MT: University of Montana. Ph.D. dissertation.
- Weaver, John L.; Paquet, Paul C.; Ruggiero, Leonard F. 1996. Resilience and conservation of large carnivores in the Rocky Mountains. *Conservation Biology*. 10(4): 964-976.
- Weckwerth, R.P.; Hawley, V.D. 1962. Marten food habits and population fluctuations in Montana. *Journal of Wildlife Management*. 26(1): 55-74.
- Weckwerth, R.P.; Wright, P.L. 1968. Results of transplanting fishers in Montana. *Journal of Wildlife Management*. 32(4): 977-980.
- Weiss, N.T.; Verts, B.J. 1984. Habitat and distribution of pygmy rabbits (*Sylvilagus idahoensis*) in Oregon. *Great Basin Naturalist*. 44(4): 563-571.
- Whisenant, Steven G. 1995. Landscape dynamics and arid land restoration. In: Roundy, Bruce A.; McArthur, E. Durant; Haley, Jennifer S.; Mann, David K., eds. Proceedings, wildland shrub and arid land restoration symposium; 1993 October 19-21; Las Vegas, NV. Gen. Tech. Rep. INT-315. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 26-34.
- Whitaker, J.O., Jr.; Maser, C. 1981. Food habits of seven species of lizards from Malheur County, southern Oregon. *Northwest Science*. 55(3):202-208.
- Whitaker, J.O., Jr.; Maser, C.; Cross, S.P. 1981. Food habits of eastern Oregon bats, based on stomach and scat analyses. *Northwest Science*. 55(4): 281-292.
- Whitaker, John O. Jr.; Maser, Chris; Keller, Laurel E. 1977. Food habits of bats of western Oregon. *Northwest Science*. 51(1): 46-55.
- White, C.M.; Thurow, T.L. 1985. Reproduction of ferruginous hawks exposed to controlled disturbance. *Condor*. 87(1): 14-22.
- White, P.J.; Ralls, Katherine; Garrott, Robert A. 1994. Coyote-kit fox interactions as revealed by telemetry. *Canadian Journal of Zoology*. 72: 1831-1836.
- Widen, P. 1987. Goshawk predation during winter, spring, and summer in a boreal forest area of central

- Sweden. *Holarctic Ecology*. 10: 104-109.
- Widen, P. 1989. The hunting habitats of goshawks, *Accipiter gentilis* in boreal forests of central Sweden. *Ibis*. 131: 205-231.
- Wielgus, Robert B.; Bunnell, Fred L. 1995. Tests of hypotheses for sexual segregation in grizzly bears. *Journal of Wildlife Management*. 59(3): 552-560.
- Wiens, John A.; Rotenberry, John T. 1981. Habitat associations and community structure of birds in shrubsteppe environments. *Ecological Monographs*. 5(1): 21-41.
- Willis, Edwin O.; Snow, David W.; Stotz, Douglas F. [and others]. 1993a. Olive-sided flycatchers in southeastern Brazil. *Wilson Bulletin*. 105: 193-194.
- Willis, M.J.; Keister, G.P., Jr.; Immell, D.A. [and others]. 1993b. Sage grouse in Oregon. *Wildl. Res. Rep.* 15. Portland: Oregon Department of Fish and Wildlife, Wildlife Research Section. 56 p.
- Wishart, William. 1978. Bighorn sheep. In: Schmidt, John L.; Gilbert, Douglas L., eds. *Big game of North America: ecology and management*. Harrisburg, PA: Stackpole Books: 161-171.
- Witmer, G.W.; Martin, S.K.; Saylor, R.D. 1998. Forest carnivore conservation and management in the interior Columbia basin: issues and environmental correlates. Gen. Tech. Rep. PNW-420. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 51 p. (Quigley, Thomas M., ed. *Interior Columbia Basin Ecosystem Management Project: scientific assessment*).
- Yensen, Eric. 1991. Taxonomy and distribution of the Idaho ground squirrel, *Spermophilus brunneus*. *Journal of Mammalogy*. 72(3): 583-600.
- Yoakum, J. 1980. Habitat management guides for the American pronghorn antelope. Tech. Note 347. Denver, CO: U.S. Department of the Interior, Bureau of Land Management. 77 p.
- Yoakum, James D. 1968. A review of distribution and abundance of American pronghorn antelope. *Biennial Antelope States Workshop Proceedings*. 3: 4-14.
- Yoakum, James D. 1978. Pronghorn. In: Schmidt, John L.; Gilbert, Douglas L., eds. *Big game of North America*. Harrisburg, PA: Stackpole Books: 103-121.
- Yoakum, James D. 1986a. Trends in pronghorn populations: 1800-1983. In: *Proceedings of the Twelfth Pronghorn Antelope Workshop*. [Meeting dates unknown]: [Meeting location unknown]. Jasper, AB: Alberta Recreation, Parks and Wildlife, Fish and Wildlife Division: 12: 77-85.
- Yoakum, James D. 1986b. Use of artemisia and chrysothamnus by pronghorns. In: McArthur, E. Durant; Welch, Bruce L., eds. *Proceedings of a symposium on the biology of Artemisia and Chrysothamnus*; 1984 July 9-13; Provo, UT. Gen. Tech. Rep. INT-200. Ogden, UT: U.S. Department of Agriculture, Forest Service, Intermountain Research Station: 176-185.
- Yoakum, Jim D.; O'Gara, Bart W. 1990. Pronghorn/livestock relationships. In: McCabe, R.E., ed. *Transactions of the 55th North American wildlife and natural resources conference*; 1990 March 16-20; Denver, CO. Washington, DC: Wildlife Management Institute: 475-487.
- Yosef, R. 1996. Loggerhead shrike (*Lanius ludovicianus*). In: Poole, A.; Gill, F., eds *The birds of North*

America. No. 231. Philadelphia: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 28 p.

Yosef, R.; Grubb, T.C., Jr. 1992. Territory size influences nutritional condition in nonbreeding loggerhead shrikes (Lanius ludovicianus): a ptilochronology approach. *Conservation Biology*. 6(3): 447-449.

Young, James V.; Bechard, Marc J. 1994. Breeding ecology of the northern goshawk in high-elevation aspen forests of northern Nevada. *The northern goshawk: ecology and management: Proceedings of a symposium of the Cooper Ornithological Society; 1993 April 14-15; Sacramento, CA*. In: *Studies in Avian Biology*. Cooper Ornithological Society; 16: 119-121.

Zager, P.L.; Mills, L.S.; Wakkinen, W.; Tallmon, D. 1995. Woodland caribou: a conservation dilemma. *Endangered Species Update*. Vol. 12 (10, 11): 1-4.

Zeveloff, Samuel I.; Collett, Farrell R. 1988. *Mammals of the intermountain west*. Salt Lake City, UT: University of Utah Press. 365 p.

Zwicker, F.C. 1972. Some effects of grazing on blue grouse during summer. *Journal of Wildlife Management*. 36: 631-634.

Zwicker, Fred C. 1992. Blue grouse, Dendragapus obscurus. In: Poole, Alan F.; Stettenheim, Peter; Gill, Frank B., eds. *The birds of North America*. No. 15. Philadelphia, PA: The Academy of Natural Sciences; Washington, DC: The American Ornithologists' Union. 28 p.

FOOTNOTES

¹ See "Abbreviations" table at end of text for definitions of abbreviated units of measure.

² See table 1, volume 1, for common and scientific names of the vertebrate broad-scale species of focus, and appendix 3, volume 3, for names of plants and animals not addressed as terrestrial vertebrate species of focus.

³ Personal communication. 1997. Wayne Wakkinen, regional wildlife biologist, Idaho Department of Fish and Game, HCR 85, Box 323-J, Bonners Ferry, ID 83805.

⁴ Personal communication. 1997. Robert Naney, wildlife biologist, Okanogan National Forest, 1240 South Second Avenue, Okanogan, WA 98840-9723.

⁵ Personal communication. 1998. David Mattson, U.S.G.S. Forest and Rangeland Ecosystem Science Center and Department of Fish and Wildlife Resources, University of Idaho, Moscow, ID 83844-1136.

⁶ The Interagency Grizzly Bear Committee is composed of top officials from the U.S.D.I. Fish and Wildlife Service, USDA Forest Service, USDI National Park Service, USDI Bureau of Land Management, USDI Bureau of Indian Affairs, state fish and game agencies of Montana, Wyoming, Idaho, and Washington, and management authorities from British Columbia and Alberta.

⁷ Personal communication. 1997. Rolf Johnson, manager - deer and elk section, Washington Department of Fish and Wildlife, 600 Capitol Way North, Olympia, WA 98501.

⁸ Personal communication. 1997. John McCarthy, special projects coordinator, Montana Department of Fish, Wildlife, and Parks, P.O. Box 200701, Helena, MT 59620-0701.

⁹ Personal communication. 1997. Lonk Kuck, wildlife game and research manager, Bureau of Wildlife, Idaho Department of Fish and Game, P.O. Box 25, Boise, ID 83707-0025.

¹⁰ Personal communication. 1998. Victor Coggins, regional wildlife biologist, Oregon Department of Fish and Wildlife, 65495 Alder Slope, Enterprise, OR 97828.

¹¹ Personal communication. 1998. Ron Garner, assistant district wildlife biologist, Oregon Department of Fish and Wildlife, P.O. Box 8, Hines, OR 97738.

¹² Personal communication. 1997. David Dobkin, wildlife biologist, High Desert Ecological Research Institute, 15 SW Colorado, Suite 300, Bend, OR 97702.

¹³ Personal communication. 1997. John Connelly, Upland Bird Research Coordinator, Idaho Department of Fish and Game, P.O. Box 25, Boise, ID 83707-0025.

¹⁴ Personal communication. 1997. Terrell D. Rich, National Avian Ecologist, U.S.D.I. Bureau of Land Management, 1387 S. Vinnell Way, Boise, ID 83709.

¹⁵ Personal communication. 1997. Tim Thier, wildlife biologist, Montana Department of Fish, Wildlife, and Parks, P.O. Box 507, Trego, MT 59934.