

Table 8--Percentage of ecological reporting units (ERUs) having various combinations of trend categories (TC_s) of relative change for each of 91 broad-scale species of focus^{a, b}

Family	Group	Common name	Number of ERUs	Percentage of ERUs in -1 or -2 category	Percentage of ERUs in 0 category	Percentage of ERUs in 1 or 2 category
1	2	Lewis' woodpecker (migrant population)	11	100	0	0
10	32	Ground snake	2	100	0	0
10	32	Washington ground squirrel	4	100	0	0
12	36	Columbian sharp-tailed grouse (summer)	11	91	9	0
12	37	Grasshopper sparrow	11	91	9	0
5	22	Rocky Mountain bighorn sheep (winter)	9	89	11	0
1	1	Pygmy nuthatch	13	85	15	0
10	31	Burrowing owl	13	85	15	0
10	31	Short-eared owl	13	85	15	0
10	31	Vesper sparrow	13	85	15	0
2	5	Flammulated owl	13	85	8	8
2	6	Williamson's sapsucker	13	85	0	15
8	29	Western bluebird	11	82	18	0
1	1	White-headed woodpecker	9	78	22	0
5	22	Rocky Mountain bighorn sheep (summer)	9	78	22	0
10	31	Western meadowlark	13	77	23	0
2	4	Blue grouse (winter)	13	77	15	8
10	31	Lark sparrow	13	77	15	8
2	6	Brown creeper	13	77	8	15
10	32	Longnose leopard lizard	8	75	25	0
10	32	Preble's shrew	12	75	25	0
10	32	Uinta ground squirrel	4	75	25	0
11	33	Lark bunting	4	75	25	0
12	37	Clay-colored sparrow	4	75	0	25
11	33	Sagebrush vole	11	73	18	9
11	34	Black-throated sparrow	7	71	29	0
5	22	California bighorn sheep	10	70	30	0
7	28	Pallid bat	10	70	30	0
10	31	Ferruginous hawk	10	70	30	0
10	31	Pronghorn	10	70	30	0
10	32	Striped whipsnake	10	70	30	0
2	7	Boreal owl	10	70	20	10
2	5	Northern goshawk (summer)	13	69	15	15
2	6	Hammond's flycatcher	13	69	15	15
2	6	Silver-haired bat	13	69	15	15
11	33	Brewer's sparrow	12	67	33	0

Table 8--Percentage of ecological reporting units (ERUs) having various combinations of trend categories (TC_s) of relative change for each of the 91 broad-scale species of focus^{a, b} (continued)

Family	Group	Common name	Number of ERUs	Percentage of ERUs in -1 or -2 category	Percentage of ERUs in 0 category	Percentage of ERUs in 1 or 2 category
12	37	Idaho ground squirrel	3	67	33	0
11	33	Sage grouse (summer)	12	67	25	8
11	33	Sage grouse (winter)	12	67	25	8
2	9	Black-backed woodpecker	12	67	8	25
11	33	Sage thrasher	11	64	36	0
11	33	Sage sparrow	11	64	36	0
1	1	White-breasted nuthatch	13	62	31	8
7	28	Western small-footed myotis	13	62	31	8
11	35	Loggerhead shrike	13	62	23	15
11	33	Pygmy rabbit	10	60	30	10
2	11	White-winged crossbill	10	60	10	30
2	12	Woodland caribou	5	60	0	40
6	23	Broad-tailed hummingbird	7	57	14	29
6	23	Rufous hummingbird	13	54	31	15
2	13	Northern flying squirrel	13	54	23	23
2	5	American marten	13	54	8	38
6	25	Northern goshawk (winter)	13	54	8	38
2	5	Fisher	13	54	0	46
11	34	Kit fox	4	50	50	0
2	6	Chestnut-backed chickadee	10	50	20	30
3	17	Blue grouse (summer)	13	46	15	38
4	18	Lazuli bunting	13	46	15	38
2	6	Golden-crowned kinglet	13	46	8	46
2	11	Three-toed woodpecker	13	46	0	54
1	3	Western gray squirrel	7	43	43	14
2	8	Great gray owl	12	42	17	42
2	6	Winter wren	12	42	0	58
10	32	Wyoming ground squirrel	5	40	60	0
7	27	Townsend's big-eared bat	13	38	46	15
2	6	Varied thrush	11	36	9	55
5	19	Grizzly bear	13	31	62	8
2	6	Hoary bat	13	31	54	15
5	21	Long-eared owl	13	31	54	15
2	10	Olive-sided flycatcher	13	31	15	54
2	6	Pileated woodpecker	10	30	10	60
2	6	Vaux's swift	11	27	0	73
7	28	Spotted bat	12	25	58	17

Table 8--Percentage of ecological reporting units (ERUs) having various combinations of trend categories (TC_s) of relative change for each of the 91 broad-scale species of focus^{a, b} (continued)

Family	Group	Common name	Number of ERUs	Percentage of ERUs in -1 or -2 category	Percentage of ERUs in 0 category	Percentage of ERUs in 1 or 2 category
5	20	Mountain goat	8	25	38	38
6	24	California mountain kingsnake	4	25	25	50
5	19	Gray wolf	13	23	77	0
6	24	Black-chinned hummingbird	13	23	23	54
7	27	Pine siskin	13	23	8	69
10	32	White-tailed antelope squirrel	5	20	80	0
7	26	Long-eared myotis	13	15	85	0
3	15	Wolverine	13	15	15	69
3	16	Lynx	9	11	67	22
3	17	Mountain quail (summer)	9	11	33	56
7	26	Yuma myotis	11	9	91	0
10	32	Mojave black-collared lizard	2	0	100	0
10	32	Longnose snake	1	0	100	0
NA	38	Black rosy finch	7	0	100	0
NA	38	Gray-crowned rosy finch	11	0	100	0
NA	39	Lewis' woodpecker (resident population)	1	0	100	0
7	26	Fringed myotis	11	0	73	27
3	15	Pygmy shrew	5	0	60	40
7	26	Long-legged myotis	13	0	54	46
9	30	Bushtit	9	0	22	78
9	30	Ash-throated flycatcher	9	0	11	89
3	14	Hermit warbler	4	0	0	100
6	24	Sharptail snake	3	0	0	100
NA	40	Brown-headed cowbird	13	0	0	100

NA = not applicable; these species not assigned to families.

^a Trend categories were defined such that -2 equals a decrease ≥ 60 percent; -1 equals a decrease ≥ 20 percent and < 60 percent; 0 equals a decrease or increase of < 20 percent; 1 equals an increase ≥ 20 percent and < 60 percent; and 2 equals an increase ≥ 60 percent.

^b Species are listed in descending order by percentage of ERUs with a trend category of -1 or -2. Percentages were calculated only for ERUs where the species occurred.